



## LATE WINTER 2018 MUSKOX PHOTO COMPOSITION SURVEY, EAST ARM OF GREAT SLAVE LAKE

---

Jan Adamczewski<sup>1</sup>, Kristen Olesen<sup>2</sup>, Dave Olesen<sup>2</sup>,  
Judy Williams<sup>1</sup>, Dean Cluff<sup>1</sup> and John Boulanger<sup>3</sup>

<sup>1</sup> Environment and Natural Resources, Government of Northwest Territories

<sup>2</sup> Hoarfrost River Huskies Ltd., Yellowknife, NWT

<sup>3</sup> Integrated Ecological Research, Nelson, BC

2021

Manuscript Report No. 296

The contents of this report are the sole responsibility of the authors.



## ABSTRACT

A strip-transect fixed-wing survey of muskoxen in the East Arm region of the Northwest Territories in late winter 2018 (Cluff et al. 2019) demonstrated that muskox densities had increased by approximately six-fold since the last previous muskox survey in this area in 2010. This represented an exceptionally high annual growth rate of about 25%/year over this period. A large portion of the survey area was above treeline and a substantial proportion was below treeline, either in transition boreal-tundra zones or well into the boreal forest. To add to our understanding of this expanding population, additional flying was carried out to obtain a demographic profile through a photographic composition survey.

An Aviat Husky, a small fixed-wing aircraft, was used with a crew of D. Olesen (pilot) and K. Olesen (photographer) in late winter 2018 for this composition survey. Muskox groups found during the larger regional muskox survey were photographed, with the emphasis on obtaining frontal views of the muskoxen. The Husky is a small, quiet and slow-flying aircraft and alarm responses of muskoxen appeared limited. Multiple photos and photo passes were taken to improve the chances of clear images showing the heads of the muskoxen. The photos were later reviewed to identify calves (approaching one year of age), two-year-old females, two-year-old males, three-year-old females, three-year-old males, females four-years-old or older and males four-years-old or older. In most cases, one main photo often provided a clear view of most of the muskoxen in the group. Having a sequence of photos made it possible to identify all or nearly all individuals in a group, where more than one photo allowed additional views of some individuals. These were compared to an Alaska Fish and Game photo guide to sex and age classes of muskoxen (Alaska Department of Fish and Game 2010).

Classification of muskoxen was limited to groups where at least 80% of the animals could be classified. Some muskox groups were in thickly forested areas where most of the animals could not be identified. In some cases it was not possible to record the frontal views needed to classify individual muskoxen. Overall, classification of muskoxen was possible in 56 of 75 (75%) of the groups photographed.

Within the 56 groups, 891 muskoxen were classified. Totals of each category of muskox classified were: calves 215, two-year-old females 87, three-year-old females 68, four-year-old+ females 148, unknown females 34 (identified as females but the age class based on horns was not quite clear in photos), total females 337, two-year-old males 88, three-year-old males 72, four-year-old+ males 122, total males 282 and unknown 57 (6.4%). These results yielded a male-female ratio of 83.7 bulls:100 cows, a calf-cow ratio of 63.8 calves:100 cows when all cows were included and a calf-cow ratio of 86 calves:100

cows if the 87 two-year-old females were omitted. Females approaching two years of age would have had to breed at about six months of age to have a calf nearly a year old at the time of the survey, which is unlikely in muskoxen. This demographic profile is consistent with a population growing at a near-maximum rate and suggests that pregnancy rates were very high along with very high survival rates of calves and adults.

Of the classified groups there were 17 bull-only groups, which ranged in size from single bulls (four) to pairs (six), three bulls (four) and single instances of five, seven and eight bulls. There were 39 mixed groups, which were generally larger and ranged in size from five to 40, with groups of ten to 15 (seven), 16-20 (eight) and 21-25 (six) occurring most frequently.

Extensive feeding sign and trails in the snow around several groups suggested that muskoxen often remained at the same site for several days or weeks.

Habitats in which muskox groups were found were defined at a broad scale as tundra, transition (tundra-boreal) and forest. Of the 56 classified groups, 39 (69.6%) were in the boreal forest, ten (17.9%) were in transition areas and seven (12.5%) were on the tundra. The high proportion of groups photographed south of treeline compared to tundra is largely a reflection of the flying effort, which was primarily in the boreal forest. Habitats south of treeline were further divided on a finer scale and the numbers and proportions of groups found in these habitats from most common to least common were: rocky ridge 17 (34.7%), sparse forest 14 (28.6%), lake edge nine (18.4%), thick forest eight (16.3%) and old burn one (2.0%). It is important to note that these habitat types are not based on any rigorous analysis of vegetation/terrain types; rather they are designations of convenience from the photos.

## TABLE OF CONTENTS

|  |     |
|--|-----|
| ABSTRACT .....   | iii |
| LIST OF FIGURES .....  | vi  |
| LIST OF TABLES .....   | vii |
| INTRODUCTION .....   | 1   |
| METHODS .....  | 3   |
| Survey Flying .....  | 3   |
| Photography of Muskox Groups .....   | 4   |
| Classification of Muskoxen from Photos .....   | 5   |
| RESULTS .....  | 8   |
| Overall Classification Results .....   | 8   |
| Distribution and Composition of Muskox Groups Classified .....   | 10  |
| Muskox Group Sizes .....   | 12  |
| Wolves and Wolf Kills .....  | 13  |
| Habitats used by Muskoxen .....  | 13  |
| DISCUSSION .....   | 16  |
| Demographic Profile of the Muskox Population in the East Arm Area of Great Slave Lake .....  | 16  |
| Utility of a Small Fixed-wing Photographic Survey to Classify Muskoxen .....   | 18  |
| Habitat use by Muskoxen in the East Arm Area .....   | 19  |
| ACKNOWLEDGEMENTS .....   | 20  |
| LITERATURE CITED .....   | 21  |
| APPENDIX 1. COMPOSITION, HABITAT TYPES AND LOCATIONS OF MUSKOX GROUPS CLASSIFIED IN EAST ARM AREA OF GREAT SLAVE LAKE, NWT, MARCH-APRIL 2018 ..... | 23  |
| APPENDIX 2. SELECTED MUSKOX PHOTOS (MIXED GROUPS) .....  | 26  |
| APPENDIX 3: SELECTED MUSKOX PHOTOS (BULL-ONLY GROUPS) .....  | 33  |

## LIST OF FIGURES

|  |    |
|--|----|
| <b>Figure 1.</b> Muskox fixed-wing survey area in late winter 2018 in the East Arm area of Great Slave Lake.....   | 3  |
| <b>Figure 2.</b> A group of muskoxen photographed March 4, 2018 on a small ridge in the boreal forest in the East Arm area of Great Slave Lake.....  | 4  |
| <b>Figure 3.</b> Part of a group of muskoxen photographed April 9, 2018 on a small ridge in the boreal forest in the East Arm area of Great Slave Lake.....  | 6  |
| <b>Figure 4.</b> A group of muskoxen photographed March 12, 2018 in the boreal forest in the East Arm area of Great Slave Lake.....  | 7  |
| <b>Figure 5.</b> Two muskox bulls photographed during March-April 2018 in the East Arm area of Great Slave Lake.....   | 10 |
| <b>Figure 6a.</b> Locations, group size and composition of muskox groups classified in the East Arm area of Great Slave Lake in March-April 2018.....  | 11 |
| <b>Figure 6b.</b> Locations, group size and composition of muskox groups classified in the East Arm area of Great Slave Lake in March-April 2018.....  | 12 |
| <b>Figure 7.</b> (a) Group size distribution of bull-only (left) and (b) mixed (right) muskox groups classified in the East Arm area of Great Slave Lake in March-April 2018.....                              | 13 |
| <b>Figure 8.</b> (a) A black wolf photographed March 16, 2018 in the East Arm area of Great Slave Lake during a muskox survey (left); (b) nearby kill site on the same day (right).....                        | 13 |
| <b>Figure 9a.</b> A group of muskoxen photographed south of treeline on April 7, 2019 in rugged terrain with multiple ridges and relatively sparse forest cover, in the East Arm area of Great Slave Lake..... | 14 |
| <b>Figure 9b.</b> A group of muskoxen photographed on the tundra on April 7, 2018 in the East Arm area of Great Slave Lake.....  | 15 |

## LIST OF TABLES

|  |   |
|--|---|
| <b>Table 1.</b> Numbers of muskoxen classified in each category in March-April 2018 in the East Arm area of Great Slave Lake and resulting bull:cow and calf:cow ratios..... | 9 |
|--|---|

## INTRODUCTION

Free-ranging muskoxen (*Ovibos moschatus*) have shown a highly variable productivity of calves, depending primarily on whether the population was growing, stable or declining. Gray (1987) observed muskoxen on the ground in the Polar Bear Pass region of Bathurst Island between 1968 and 1980 and in the first three years (1968-1970) did not see a single calf and little evidence of rutting behaviour. Tener (1965) noted that “the evidence gathered by early travellers in muskox country suggested that cows produced calves in alternate years.” Percentages of calves recorded by observers in the 1950s and early 1960s in Tener (1965) varied “from 0 to as high as 18%”. Introduced populations in Alaska showed very rapid growth in some cases; the calf-cow ratio in the Sadlerochit River muskox herd was 89 calves:100 cows of reproductive age in 1979 (Jingfors and Klein 1982). A rapidly expanding population in northern Quebec (Le Hénaff and Crête 1989) had similar calf productivity and grew at 25% per year. In an Alaskan introduced population in the Arctic National Wildlife Refuge, some cows (early 1980s) initially produced calves in several successive years, but by 1991-1993 most females successfully reproduced at intervals of two-three years (Reynolds 2001). Larter and Nagy (2001) documented substantial variation in calf production and yearling recruitment in Banks Island muskoxen between 1986 and 1999 while this population increased from 1986-1994 and then declined to 1999.

Muskoxen can be identified to sex and age class by their growth and horn development. A number of guides to age and sex classification have been published: Henrichsen and Grue (1980) developed one of the first guides in Greenland; Gray (1987) included a guide to muskox classification on Bathurst Island in his book; Olesen and Thing (1989) developed a muskox photo guide in West Greenland; and Alaska Department of Fish and Game (2010) developed a photo guide for Alaskan muskoxen. Comparison of these guides suggests that the rates at which muskoxen reach mature size and horn growth vary, depending on the population phase that the animals are in. The guide in Gray (1987) included males identified as one, two, three, four, five and six+ years old, Henrichsen and Grue (1980) recognized males one, two, three, four and five+ years old, while the Alaska Department of Fish and Game guide (2010) included one, two, three and four+ year old male classes. These differences most likely reflect more rapid growth and better nutrition in expanding populations (e.g. Olesen and Thing 1989, Alaska Fish and Game 2010) than in stable or declining populations (e.g. Gray 1987).

Classification surveys to identify muskox sex and age classes have been primarily conducted on the ground with binoculars and spotting scopes (Jingfors and Klein 1982, Larter and Nagy 1999, 2001), usually with air support on large remote ranges (Larter and Nagy 2001). A photographic approach was used in muskox surveys of the Nelson and



Nunivak Islands in Alaska (e.g. Jones 2015a, b). These were conducted by small, quiet slow-flying aircraft such as Piper PA-18 Supercubs, which tend to disturb the animals relatively little and allow high-resolution photos to be taken of the front ends of the muskoxen. Muskoxen can later be classified to age and sex classes from the photos (see Jones 2015a, b).


A large-scale muskox survey in an area around the East Arm of Great Slave Lake in the Northwest Territories (NWT) by Cluff et al. (2019) in February and March 2018 showed an increase of approximately six-fold in abundance from a previous similar survey in 2010. This translates to an annual growth rate of about 25%, similar to the 25% annual growth rate documented in an expanding colonizing muskox population in northern Quebec (Le Hénaff and Crête 1989).

To more fully understand the demography of this rapidly growing muskox population in the East Arm area, we flew a photographic composition survey in March and early April 2018 with an Aviat Husky, a modern-day Supercub. The purpose was to photograph a sample of the muskox groups found during the survey by Cluff et al. (2019) and identify proportions of muskox sex and age classes. Unlike the muskoxen on Nelson and Nunivak Islands, which are on open tundra, much of the study area around the East Arm of Great Slave Lake is below treeline in the boreal forest, thus our survey was in part a pilot project to assess whether this photographic method could also be used for muskoxen in forested country.

## METHODS

### Survey Flying

A fixed-wing muskox survey was flown between February 26 and March 2, 2018 in an area around the East Arm of Great Slave Lake in the NWT (Cluff et al. 2019; Figure 1).



**Figure 1.** Muskox fixed-wing survey area in late winter 2018 in the East Arm area of Great Slave Lake. Grey lines were part of the large regional survey flown by Cluff et al. (2019). Red dots show locations of muskox groups. Pink lines are the additional lines flown in the southern part of the survey area with the Husky and the additional flying to photograph muskox groups. Black dots show muskox groups seen during the Husky flying.

The area surveyed in 2018 was larger and included an area east of the survey area from 2010, to provide information on muskox abundance in an area that had some reported muskox sightings in recent years. Further flight lines were added to the south of the main survey area using the Husky in March and April 2018, to provide further information on muskox abundance in that area. Information from local observers and particularly D. Olesen who is a pilot based at the Hoarfrost River and has flown this area extensively, indicated that muskoxen were abundant in that area and had increased substantially in recent years. The overall survey area was mostly on the tundra above treeline, but also included a substantial proportion below treeline. The highest densities of muskox groups were at the south end of the survey area south of Łutsel K'e, well below treeline.

Overall, muskox density (number per km<sup>2</sup>) in the East Arm area had increased about six-fold over the period between 2010 and 2018 (Cluff et al. 2019).

### Photography of Muskox Groups

The additional flying that was used to photograph muskox groups for classification, which this report describes, was carried out by D. and K. Olesen in March and April, 2018 in the Aviat Husky. The aircraft is small and suited to one pilot and one passenger and no Department of Environment and Natural Resource (ENR) biologists were on board during these flights. K. Olesen is an experienced photographer and we were interested in obtaining clear high-resolution photos from which muskoxen could be classified. Some of the photography was taken while flying the southern-most flight lines in Figure 1 and during opportunistic flights in following weeks when suitable weather occurred and when the aircraft and pilot were available. Waypoints recorded on GPS units for muskox groups from the fixed-wing survey lines in Figure 1 were also used a few days or a few weeks after the fixed-wing survey lines had been completed. Muskoxen in late winter are fairly localized in their movements, thus GPS locations recorded a few days earlier or even a week or two previously almost always meant that the animals would not be far away. Photos were taken on eight days: March 4, 12, 13, 14, 16 and 19, and April 7 and 19.



**Figure 2.** A group of muskoxen photographed March 4, 2018 on a small ridge in the boreal forest in the East Arm area of Great Slave Lake. Most are standing facing the aircraft and can readily be identified to age and sex.

A Nikon D3X and a Nikon D800 camera were used for the photography, in combination with a number of zoom lenses and primarily a Nikkor AF-S 28-300mm lens. For most groups, a number of photo passes (two-three) were made. The emphasis was on photos

that showed most of the muskoxen in a group from a frontal view where their horns were clearly visible. In some cases, it appeared that the muskoxen were not much disturbed by the aircraft and they lined up facing the aircraft and the camera (Figure 2).

For all groups photographed, multiple photos were taken in sequence, to increase the chances that most or all of the animals would be positioned so that their front ends could be clearly seen. All photos of muskoxen were stored on an external hard drive, together with information about the group's location and habitat they were found in. All photos were taken by K. Olesen, including all examples in this report.

### **Classification of Muskoxen from Photos**

All the photos of muskox groups were reviewed on a computer monitor. For most groups, one photo pass generally had the most suitable photos with the animals largely facing the camera. Within that photo pass, it was often possible to select the one best photo where most of the muskoxen were clearly identifiable. A program was used to digitally mark each muskox and each animal was then classified using the photo guide from Alaska Department of Fish and Game (2010). We used this guide in preference to the other guides listed earlier, as it appeared that the Alaskan guide was from a muskox population growing rapidly, thus most appropriate for a population in this area. Photos in the sequence before and after the main photo were also used as this sometimes meant that muskoxen that were obscured by other animals or behind trees or were turned away from the camera could in a further photo be seen clearly and classified.

For most groups, it appeared that the muskoxen were not greatly alarmed by the aircraft, which is relatively small and quiet and can fly relatively slowly. On several photos there were bedded muskoxen that did not stand up. In several cases the best photos were taken on the first pass when the animals lined up to face the aircraft and the camera. An example of a small group of muskoxen that included calves and two and three year old males is shown in Figure 3.

In some groups photographed, most or many of the muskoxen could not be classified, either because they were obscured in thick forest (Figure 4), or photos of the front ends were not feasible, or in a few cases they were tightly bunched so that some individuals were behind others, or they were on the run. We used only classification of groups where we could classify at least 80% of the animals, as we were concerned that in groups where we classified lower percentages of the animals, the counts might be biased towards fairly obvious animals like large bulls or calves. Although the bulk of the muskox groups were photographed south of treeline, many were in relatively open areas, such as ridges or small hills, lake edges with few trees, or relatively open, sparsely forested areas. Due to visibility,

muskox groups on the tundra were relatively easy to photograph and muskox groups in thickly forested locations tended to be more difficult to photograph.

Bootstrap methods (Manly 1977) were used to obtain percentile-based confidence limits on bull-cow and calf-cow ratios. The *boot* package in R statistical software (R Development Core Team, 2009) was used for bootstrap estimates. Pie charts maps were generated using the QGIS software package (QGIS Foundation 2015).



**Figure 3.** Part of a group of muskoxen photographed April 9, 2018 on a small ridge in the boreal forest in the East Arm area of Great Slave Lake. From left to right, the muskoxen are: a calf, a two-year-old male, a calf, a calf and a three-year-old male.



**Figure 4.** A group of muskoxen photographed March 12, 2018 in the boreal forest in the East Arm area of Great Slave Lake.

## RESULTS

### Overall Classification Results

We classified 869 muskoxen in 56 groups and were unable to obtain adequate classification in another 19 groups, hence we classified 75% of the groups photographed. The muskox classes we identified based on the Alaskan guide included calves (approaching one year), two-year-old females, three-year-old females, females at least four-years-old, two-year-old males, three-year-old males and males at least four-years-old (Table 1). Calves are difficult to identify as males and females as their body size is similar and there are no obvious differences yet in the horns. Of the 869 muskoxen, 57 (6.5%) were classified as unknown. In addition, 34 muskoxen (out of 337 total females, or 10.1% of the females) were identified as females based on thin horn bases, but could not be definitively identified to age class. Numbers of each sex and age class in each group classified are listed in Appendix 1. A selection of photos of muskox groups is provided in Appendix 2 (mixed groups) and Appendix 3 (bull-only groups).

Calves accounted for nearly a quarter (24.7%) of the 891 total muskoxen classified. If all the females are included, then the calf:cow ratio was 63.9 calves:100 cows. However, the survey was carried out in March, when all the muskoxen would have been approaching their birth dates (calving likely occurred in late April and early May). Two-year-old cows would have had to breed at five months of age to have a newborn calf in April/May 2017, which is unlikely. If the 87 two-year-old females are excluded, then the calf:cow ratio was 86.0 calves:100 cows.

**Table 1.** Numbers of muskoxen classified in each category in March-April 2018 in the East Arm area of Great Slave Lake and resulting bull:cow and calf:cow ratios. Bootstrap-based 95% confidence limits are given for ratios. Bull:cow ratio includes bull-only and mixed groups. Upper confidence limit on calf:cow ratio is truncated at 100 calves:100 cows.

| Category   | Number Classified              | % of Total | Notes  |
|--|--------------------------------|------------|--|
| <b>Calves (nearly one year old)</b>                      | 215                            | 24.7       | % of 891   |
| <b>Females two years old</b>                             | 87                             | 10.0       | % of 891   |
| <b>Females three years old</b>                           | 68                             | 7.8        | % of 891   |
| <b>Females at least four years old</b>                   | 148                            | 17.0       | % of 891   |
| <b>Unknown Females</b>                                   | 34                             | 3.9        | % of 891   |
| <b>Total Females</b>                                     | 337                            | 54.4       | <i>% of Total Males and Total Females (619)</i>  |
| <b>Males two years old</b>                               | 88                             | 10.1       | % of 891   |
| <b>Males three years old</b>                             | 72                             | 8.3        | % of 891   |
| <b>Males at least four years old</b>                     | 122                            | 14.0       | % of 891   |
| <b>Total Males</b>                                       | 282                            | 45.6       | <i>% of Total Males and Total Females (619)</i>  |
| <b>Unknown</b>   | 57                             | 6.6        | % of 891   |
| <b>Total</b>   | 869                            | 99.9       |  |
| <b>Ratios</b>  |                                |            |  |
| <b>Bulls:100 Cows</b>                                    | 83.7:100, (SE=9, CI=69-106).   |            | 282 Males and 337 Females                        |
| <b>Calves:100 Cows</b>                                   | 63.8:100 (SE=5, CI=55-74:100)  |            | 215 Calves and 337 Females                       |
| <b>Calves:100 Cows (excluding females two years old)</b> | 86.0:100 (SE=8, CI=71-100:100) |            | 215 Calves and 250 Cows three years old or older |

The representation of two and three year old males and females was high in this survey, consistent with the high representation of calves in the population. The total of two and three year old females was 155, larger than the total of females classified as four years old or older (148) and the total of two and three year old males was 160, substantially more than the total of males classified as four years old or older (122).

Close assessment of the males classified as at least four years old suggested that many of these males were no more than four or five years old, as their bosses appeared light-coloured and did not show the full central development and darker colour often seen in older males (Figure 5).







**Figure 5.** Two muskox bulls photographed during March-April 2018 in the East Arm area of Great Slave Lake. Both were identified as at least four years old when compared to the Alaska Department of Fish and Game (2010) guide, however the one on the left has a very light-coloured boss and lacks the full development of the boss compared to the male on the right. The bull on the left is likely just four years old while the one on the right is likely at minimum five or six years old.

#### **Distribution and Composition of Muskox Groups Classified**

Muskox groups classified in March and April 2018 were mapped in Figure 6a and 7b. Group size and composition were generally similar throughout the survey area. Bull-only groups tended to be more common at the southern end of the distribution.




**Figure 6a.** Locations, group size and composition of muskox groups classified in the East Arm area of Great Slave Lake in March-April 2018. Circles are sized in proportion to group size. Calves, males, females and unknown muskoxen are shown in the pie charts.



**Figure 6b.** Locations, group size and composition of muskox groups classified in the East Arm area of Great Slave Lake in March-April 2018. Circles are sized in proportion to group size. Calves, males and females two, three and four+ plus years old and unknown muskoxen are shown in the pie charts.

### Muskox Group Sizes

There were 17 bull-only muskox groups among the 56 groups classified. Bull-only groups ranged in size from lone bulls to a group of eight bulls, with the most common being pairs (Figure 7a). Mixed groups were generally larger and ranged in size from five to 40, with groups of 11-15, 16-20 and 21-25 being most frequent (Figure 7b).



**Figure 7.** (a) Group size distribution of bull-only (left). (b) Mixed (right) muskox groups classified in the East Arm area of Great Slave Lake in March-April 2018.

### Wolves and Wolf Kills

In the flying carried out by the Aviat Husky, two wolves (one black, one grey) were seen at a kill site location, which appeared to be the remains of a muskox (Figure 8). One additional muskox kill site was found during the flying. As bears would not likely have been active at this time of year, this was likely also a wolf kill.



**Figure 8.** (a) A black wolf photographed March 16, 2018 in the East Arm area of Great Slave Lake during a muskox survey (left); (b) nearby kill site on the same day (right). The remains appear to be a muskox.

### Habitats used by Muskoxen

Please note that the following habitat classes do not derive from any formal habitat definition; they were rather units of convenience based on the photos.

Of the 56 muskox groups photographed and classified, 39 (71.9%) were in the boreal forest, ten (17.2%) were in transition areas and seven (12.1%) were on the tundra. The high proportion of groups photographed south of treeline compared to tundra is largely a reflection of the flying effort of the Aviat Husky, which was primarily in the boreal forest. Habitats were further divided on a finer scale below treeline and the numbers and proportions of groups found in these habitats were as follows: rocky ridge 17 (29.3%), sparse forest 14 (24.1%), thick forest nine (17.2%), lake edge eight (15.5%) and old burn one (1.7%). Continuous snow cover on the tundra did not allow identification of specific vegetation or habitat types. Some muskox groups were found in relatively thick boreal forest but most were in fairly rugged terrain with at least some openings.

A common observation during the survey flying was that muskox groups observed on one day could often be found nearby (from the GPS waypoint) a few days or even a few weeks later. Areas used by muskoxen often showed extensive feeding craters, trails and signs of use consistent with multiple days of use (or longer) in the area (Figures 10a and 10b).



**Figure 9a.** A group of muskoxen photographed south of treeline on April 7, 2019 in rugged terrain with multiple ridges and relatively sparse forest cover, in the East Arm area of Great Slave Lake. Extensive sign of feeding, bedding sites and trails in the snow suggested that the muskoxen had been in the area for several days and possibly longer.



**Figure 9b.** A group of muskoxen photographed on the tundra on April 7, 2018 in the East Arm area of Great Slave Lake. As in Figure 9a, extensive sign of feeding and bedding sites suggested that the muskoxen had been in the area for several days and possibly longer.

## DISCUSSION

### Demographic Profile of the Muskox Population in the East Arm Area of Great Slave Lake

The results of the composition survey in the East Arm area of Great Slave Lake in March/April 2018 are consistent with the exceptionally rapid growth of this population documented by Cluff et al. (2019) between 2010 and 2018. The annual growth rate of about 25% for this population matches the 25% annual growth rate documented by Le Hénaff and Crête (1989) between 1983 and 1986 in a muskox population that was colonizing new range near Baie D'Ungava in northern Quebec. This muskox population originated from a captive herd that was released north of Kuujuaq in an area with no historic or archaeological record of muskox presence. There were 26 calves (23% of the population) in June 1983 and 1986, very similar to the 24.7% we recorded in 2018. The numbers are not directly comparable as the Quebec surveys were carried out in June when calves would be two months old or less, whereas our survey was in late winter when calves of the previous year were just under one year of age.

Growth rates of 23 and 24% per year were also recorded in small introduced muskox populations in northeast Alaska and the Seward Peninsula (Jingfors and Klein 1982). A calf:cow ratio of 89 calves:100 cows of reproductive age was observed in one of these Alaskan introduced herds on the Sadlerochit River in 1979, albeit the entire herd in this case was 58 animals. We recorded a very similar calf:cow ratio of 86 calves:100 cows of reproductive age in March/April 2018, although again the surveys are not directly comparable as the Alaskan surveys were in May and June when new-born calves had been born just a few weeks earlier, whereas our survey was in late winter when calves of the previous year were approaching one year of age.

A further example of a muskox population increasing at near-maximum rates between 1977 and 1990 was described by Olesen (1993) in the Angujaartorfiup Nunaa range in West Greenland. In this population, the average calf percentage was 24.0% and the average annual population increment averaged 30% (Olesen 1993). Representation of sex and age classes in this population followed Olesen (1989) and included the same categories as the Alaska Fish and Game (2010) guide. The high representation of muskoxen that were one, two and three years old observed by Olesen (1993) was similar to our results, although ground-based surveys of the West Greenland population varied in seasonal timing through the year.

The Banks Island muskox population increased from an estimated 3,800 in 1972 to 34,225 in 1989 (Gunn et al. 1991) and then further to 64,608 in 1994 (Larter and Nagy 2001). Over the period 1972-1989, the annual rate of increase was 13% (Gunn et al. 1991), a rapid

increase although well below the more rapid rates of increase in the Alaskan, West Greenland and Quebec populations cited above and in the East Arm area (Cluff et al. 2019). The highest muskox calf percentages recorded on Banks Island during aerial surveys occurred in June 1971 (25.4%) and 1972 (23.1%) but they were generally lower in the 1980s (Gunn et al. 1991).

Overall, these comparisons suggest that the muskox population in the East Arm area of Great Slave Lake was increasing between 2010 and 2018 at rates similar to the highest population growth rates observed in Alaska (Jingfors and Klein 1982), northern Quebec (Le Hénaff and Crête 1989) and West Greenland (Olesen 1993). The authors of these three studies indicated that mortality rates of all sex and age classes appeared to be minimal. The very high calf:cow ratio we documented in late winter (86.0 calves:100 cows of reproductive age), together with the 25% annual growth rate 2010-2018, could only have happened if the pregnancy rate was very high and the mortality rates of calf, young and mature muskoxen were very low in the East Arm area.

Wolves are present in the East Arm area and two muskox kill sites were recorded in the flying summarized in this report, one of them with two wolves present. Grizzly bears are also present in the region and they can be effective predators of calf and adult muskoxen (Reynolds et al. 2002). We assume that as of 2018, muskox predation rates by wolves and bears in the East Arm area were very low, either due to low numbers of wolves and bears or due to the predators not being used to hunting muskoxen as a “new” potential prey animal.

We suggest regular muskox monitoring of the East Arm area via population surveys augmented by composition surveys like the one we carried out in 2018. The near-maximal rate of increase documented between 2010 and 2018 is unlikely to last more than a few years. Rapid increases of endemic muskox populations have sometimes been followed by large declines: the muskox population on Banks Island increased rapidly in the 1970s and 1980s but then declined, with a very rapid decline between 36,676 muskoxen estimated in 2010 and 13,767 muskoxen estimated in 2014 (Davison et al. 2017). Die-offs attributed at least in part to the bacterial pathogen *Erysipelothrix* were implicated in the rapid decline of muskoxen on Banks Island (Kutz et al. 2015). Introduced populations in Alaska have sometimes increased rapidly during an initial phase (Jingfors and Klein 1982), but later predation (Reynolds et al. 2002), reduced productivity (Reynolds 2001) and a complex of health issues (Afema et al. 2017) have contributed to decline.

It is possible to use population survey and composition data to model the demography of muskox populations. A study with bison (Boulanger et al. 2021) found that it was possible to obtain survival rate estimates based on composition surveys in the absence of collar data. The demographic approach might allow further inference into trends in overall



productivity as well as calf and sub-adult survival. This general approach would be most applicable when there is a time series of population and composition surveys available.

### **Utility of a Small Fixed-wing Photographic Survey to Classify Muskoxen**

Muskoxen have been classified to sex and age classes in many locations across their circumpolar range; in most studies ground-based methods have been used for classification. In Canada the Banks Island population has had relatively frequent monitoring of population size as well as composition (Gunn et al. 1991, Larter and Nagy 1999, 2001). Given the large ranges to be covered, aerial support (helicopter) has been used and classification has been mostly carried out from the ground with binoculars and spotting scopes between June and August (Larter and Nagy 1999, 2001). For small populations in Alaska, ground-based observation has been used (e.g. Jingfors and Klein 1982, Reynolds 2001). A colonizing muskox population in northern Quebec was surveyed using a combination of a helicopter to find groups and ground-based methods to count and classify the animals (Le Hénaff and Crête 1989). Photography to count and classify muskoxen from the air using a small, quiet fixed-wing aircraft has been used for several years for the Nelson and Nunivak Island populations in Alaska (e.g. Jones 2015a and b).

The survey described here was in part undertaken to test whether photography from a small fixed-wing aircraft could be used to classify sex and age classes in an increasing muskox population in the East Arm area of Great Slave Lake. Overall, we believe that the photographic method developed in Alaska was successfully adapted to the study area around the East Arm of Great Slave Lake. The area is remote and has no roads and a substantial part of the muskox population is below treeline. We were concerned that muskox groups in the forest would be difficult to photograph in such a way as to allow most animals to be classified, with a clear view of their heads and horns. Classification was unsuccessful in about a quarter of the muskox groups photographed, either because of thick cover or because we did not obtain clear photos of the heads of the muskoxen. In a few cases muskoxen were so closely bunched that some animals were obscured, or were on the run. However, most muskox groups below treeline were on ridges or small hills, in sparsely forested areas, or on edges of lakes or ponds, thus photos were possible.

A photographic survey with a small fixed-wing aircraft to classify muskoxen has some advantages over methods that rely on a helicopter, as would be necessary otherwise because of the large, remote survey area. The hourly rate of the Husky is about  $\frac{1}{3}$  to  $\frac{1}{4}$  the hourly rate of a helicopter. The aircraft did not appear to greatly alarm the animals, as evidenced by many photos of muskoxen lined up standing to face the aircraft. In some photos some or most of the animals were bedded. The results form a long-term record of high-resolution photos which can be examined carefully and repeatedly on a computer

monitor. Classification can be repeated by more than one observer, where results of visual-only surveys are only visible to the survey crew.

We recognize that not being able to classify all muskox groups from photos is a drawback, however we have no reason to think that the unclassified groups (25% of the groups photographed) were different in composition from the ones classified successfully. Of the 869 muskoxen classified in 56 groups, 57 animals (6.6%) remained unclassified, which introduces a degree of variability to the data. However, it is unlikely that classification of those 57 muskoxen would have substantively changed the results of the survey.

### **Habitat use by Muskoxen in the East Arm Area**

This survey was not designed to assess habitat selection by muskoxen and the habitat types described are units of convenience based on the photos of muskox groups, not rigorously chosen habitat types. However, the photos did provide an indication of the kinds of terrain and vegetation that muskoxen south of treeline in the East Arm area were using. Habitats in which muskox groups were found were defined at a broad scale as tundra, transition (tundra-boreal) and forest. Of the 56 groups, 39 (69.6%) were in the boreal forest, ten (17.9%) were in transition areas and seven (12.5%) were on the tundra. The high proportion of groups photographed south of treeline compared to tundra is largely a reflection of the flying effort, which was primarily in the boreal forest.

Habitats south of treeline were further subjectively divided on a finer scale and the numbers and proportions of groups (49 total) found in these habitats from most common to least common were: rocky ridge 17 (34.7%), sparse forest 14 (28.6%), lake edge nine (18.4%), thick forest eight (16.3%) and old burn one (2.0%). Overall, 83.7% of the muskox groups were in areas with either relatively open forests or openings due to rugged terrain (ridges or small hills). Ridges or small hills tended to be wind-swept with shallow snow cover, which likely offered good feeding conditions. We assume that muskoxen found near the edges of lakes or ponds used these areas to forage on sedges or grasses. Extensive feeding sign and trails in the snow around several groups suggested that muskoxen often remained at the same site for several days or weeks. A few muskox groups were in relatively thick boreal forest, but the most common type of terrain used by muskoxen was rugged areas – ridges or small hills – that were wind-swept with limited snow cover.

## **ACKNOWLEDGEMENTS**

The survey flying in this report was funded by the Government of the Northwest Territories. We would like to thank Patrick Jones, biologist with Alaska Department of Fish and Game, for the methodology of photographing muskoxen from Supercubs and classifying them in the Nelson and Nunivak Island populations (see Jones 2015a and b). We adapted these methods to the muskox population in the East Arm area of Great Slave Lake.

## LITERATURE CITED

- Afema, J.A., K.B. Beckmen, S.M. Arthur, K.B. Huntington and J.A.K. Mazet. 2017. Disease complexity in a declining Alaskan muskox (*Ovibos moschatus*) population. *Journal of Wildlife Diseases*, 53(2):311-329.
- Alaska Department of Fish and Game. 2010. Muskox: a guide to identification, hunting and viewing. Alaska Department of Fish and Game, Nome, AK.
- Boulanger, J., K.G. Poole and C. DeMars. 2021. Review of bison monitoring program for the Northwest Territories. Environment and Natural Resources, Government of the Northwest Territories. Manuscript Report 290.
- Cluff, D. and B. Croft. 2010. A muskox survey in the East Arm area of Great Slave Lake, Northwest Territories, Canada, in late winter 2010. Environment and Natural Resources. Manuscript Report In Prep.
- Cluff, D., S. Goodman and B. Croft. 2019. A muskox survey in the East Arm area of Great Slave Lake, Northwest Territories, Canada, in late winter 2018. Department of Environment and Natural Resources, Government of Northwest Territories, Yellowknife, Northwest Territories, Canada. Manuscript Report In Prep.
- Davison, T., J. Williams and J. Adamczewski. 2017. Aerial survey of Peary caribou (*Rangifer tarandus pearyi*) and muskoxen (*Ovibos moschatus*) on Banks Island, July 2014. Environment and Natural Resources, Government of the Northwest Territories. Manuscript Report 270.
- Gray, D.R. 1987. The muskoxen of Polar Bear Pass. Markham, Ontario: Fitzhenry and Whiteside.
- Gunn, A., C. Shank and B. McLean. 1991. The history, status and management of muskox on Banks Island. *Arctic* 44(3):188-195.
- Henrichsen, P. and H. Grue. 1980. Age criteria in the muskox (*Ovibos moschatus*) from Greenland. *Danish Review of Game Biology* 11:3-15.
- Jingfors, K.T. and D.R. Klein. 1982. Productivity in recently established muskox populations in Alaska. *Journal of Wildlife Management* 46:1,092-1,096.
- Jones, P. 2015a. 2015 Nelson Island Muskox Survey Results. Alaska Department of Fish and Game, Division of Wildlife Conservation Northwest, Bethel, AK. Unpublished report. 7pp.
- Jones, P. 2015b. 2015 Nunivakand Island Muskox Survey Results. Alaska Department of Fish and Game, Division of Wildlife Conservation Northwest, Bethel, AK. Unpublished report. 4pp.

- Kutz, S.J., T. Bollinger, M. Branigan, S. Checkley, T. Davison, M. Dumond, B. Elkin, T. Forde, W. Hutchins, A. Niptanatiak and K. Orsel. 2015. *Erysipelothrix rhusiopathiae* associated with recent widespread muskox mortalities in the Canadian Arctic. *Canadian Veterinary Journal* 56:560–563.
- Larter, N.C. and J.A. Nagy. 1999. Sex and age classification surveys of muskoxen on Banks Island, 1985–1998: A review. Resources, Wildlife and Economic Development, Government of Northwest Territories. Manuscript Report No.113.
- Larter, N.C. and J.A. Nagy. 2001. Calf production, calf survival and recruitment of muskoxen on Banks Island during a period of changing population density from 1986–99. *Arctic* 54:394-406.
- Le Hénaff, D. and M. Crête. 1989. Introduction of muskoxen in northern Quebec: the demographic explosion of a colonizing herbivore. *Canadian Journal of Zoology* 67:1,102–1,105.
- Manly, B.F.J. 1997. *Randomization and Monte Carlo Methods in Biology*. 2nd edition. Chapman and Hall, NY.
- Olesen, C.R. and H. Thing. 1989. Guide to field classification by sex and age of the muskox. *Canadian Journal of Zoology* 67:1,116–1,119.
- Olesen, C.R. 1993. Rapid population increase in an introduced muskox population, West Greenland. *Rangifer* 13:27-32.
- QGIS Foundation. 2015. QGIS: A free and open geographic information system ([www.qgis.org](http://www.qgis.org)).
- R Development Core Team. 2009. R Foundation for Statistical Computing, Vienna, AT.
- Reynolds, P.E. 2001. Reproductive patterns of female muskoxen in northeastern Alaska. *Alces* 37(2):403-410.
- Reynolds, P.E., H.V. Reynolds and R.T. Shideler. 2002. Predation and multiple kills of muskoxen by grizzly bears. *Ursus* 13:79-84.
- Tener, J.S. 1965. Muskoxen in Canada: a biological and taxonomic review. *Canadian Wildlife Service Monograph Series 2*. Queen's Printer, Ottawa, ON.

## APPENDIX 1. COMPOSITION, HABITAT TYPES AND LOCATIONS OF MUSKOX GROUPS CLASSIFIED IN EAST ARM AREA OF GREAT SLAVE LAKE, NWT, MARCH-APRIL 2018

C=Calf (born previous year), F2=Female two years old, F3=Female three years old, F4=Female at least four years old; UF= Female Unknown Age, F All=All Females, M2=Male two years old, M3=Male three years old, M4=Male at least four years old, M All=All Males, U=Unknown.

| Date          | Group # | C  | F2 | F3 | F4 | UF | F All | M2 | M3 | M4 | M All | U  | Total | Habitat    | Fine Habitat  | Latitude              | Longitude  |
|---------------|---------|----|----|----|----|----|-------|----|----|----|-------|----|-------|------------|---------------|-----------------------|------------|
| March 4 2018  | 1       | 8  | 0  | 4  | 5  | 1  | 10    | 0  | 2  | 2  | 4     | 0  | 22    | transition | Sparse forest | 62.899722             | -109.18333 |
| March 4 2018  | 2       | 5  | 1  | 2  | 0  | 0  | 3     | 0  | 4  | 4  | 8     | 1  | 17    | forest     | Sparse forest | 62.973333             | -108.99778 |
| March 4 2018  | 3       | 4  | 0  | 2  | 7  | 3  | 12    | 4  | 2  | 5  | 11    | 2  | 29    | tundra     | Tundra        | 62.971111             | -109.00278 |
| March 4 2018  | 4       | 7  | 2  | 0  | 6  | 3  | 11    | 1  | 1  | 2  | 4     | 1  | 23    | transition | Sparse forest | 62.9075               | -108.81167 |
| March 4 2018  | 5       | 12 | 1  | 0  | 5  | 7  | 13    | 5  | 2  | 0  | 7     | 8  | 40    | transition | Sparse forest | 62.935                | -109.27389 |
| March 4 2018  | 6       | 9  | 0  | 0  | 5  | 4  | 9     | 2  | 2  | 0  | 4     | 14 | 36    | transition | Sparse forest | 62.92                 | -109.30722 |
| March 4 2018  | 7       | 6  | 2  | 1  | 6  | 0  | 9     | 4  | 2  | 0  | 6     | 1  | 22    | transition | Sparse forest | 62.890833             | -109.38194 |
| March 12 2018 | 8       | 6  | 2  | 2  | 4  | 0  | 8     | 2  | 2  | 1  | 5     | 0  | 19    | forest     | Lake edge     | 62.65148              | -108.88193 |
| March 12 2018 | 9       | 5  | 3  | 1  | 3  | 0  | 7     | 4  | 2  | 1  | 7     | 1  | 20    | forest     | Lake edge     | 62.502838             | -108.71008 |
| March 12 2018 | 10      | 0  | 0  | 0  | 0  | 0  | 0     | 0  | 0  | 5  | 5     | 0  | 5     | forest     | Lake edge     | 62.314269             | -108.41204 |
| March 12 2018 | 11      | 0  | 0  | 0  | 0  | 0  | 0     | 0  | 0  | 2  | 2     | 0  | 2     | forest     | Sparse forest | 62.025556             | -109.15778 |
| March 12 2018 | 12      | 0  | 0  | 0  | 0  | 0  | 0     | 0  | 0  | 2  | 2     | 0  | 2     | forest     | Sparse forest | 61.947906             | -109.18696 |
| March 12 2018 | 13      | 0  | 0  | 0  | 0  | 0  | 0     | 0  | 0  | 3  | 3     | 0  | 3     | forest     | Lake edge     | 61.934167             | -108.69389 |
| March 12 2018 | 14      | 2  | 1  | 0  | 3  | 0  | 4     | 3  | 1  | 2  | 6     | 1  | 13    | forest     | Rocky ridge   | No wpt; near group 13 |            |
| March 12 2018 | 15      | 0  | 0  | 0  | 0  | 0  | 0     | 0  | 0  | 3  | 3     | 0  | 3     | forest     | Rocky ridge   | No wpt; near group 13 |            |
| March 13 2018 | 16      | 4  | 1  | 0  | 3  | 0  | 4     | 3  | 0  | 5  | 8     | 0  | 16    | forest     | Rocky ridge   | 62.675245             | -109.75054 |
| March 13 2018 | 17      | 0  | 0  | 0  | 0  | 0  | 0     | 0  | 0  | 2  | 2     | 0  | 2     | forest     | Rocky ridge   | 62.551328             | -110.18407 |
| March 13 2018 | 18      | 2  | 0  | 0  | 2  | 0  | 2     | 0  | 4  | 1  | 5     | 0  | 9     | forest     | Thick forest  | 62.087986             | -111.41419 |
| March 13 2018 | 19      | 6  | 3  | 3  | 5  | 1  | 12    | 3  | 3  | 3  | 9     | 1  | 28    | forest     | Rocky ridge   | 62.036803             | -111.40877 |
| March 13 2018 | 20      | 1  | 1  | 0  | 2  | 0  | 3     | 0  | 0  | 1  | 1     | 0  | 5     | forest     | Thick forest  | 61.97215              | -109.26825 |
| March 13 2018 | 21      | 0  | 0  | 0  | 0  | 0  | 0     | 0  | 0  | 2  | 2     | 0  | 2     | forest     | Thick forest  | 61.934436             | -109.35495 |
| March 13 2018 | 22      | 5  | 2  | 4  | 3  | 1  | 10    | 1  | 0  | 0  | 1     | 0  | 16    | forest     | Old burn      | 61.94115              | -110.09775 |

| Date          | Group # | C  | F2 | F3 | F4 | UF | F All | M2 | M3 | M4 | M All | U | Total | Habitat    | Fine Habitat  | Latitude              | Longitude  |
|---------------|---------|----|----|----|----|----|-------|----|----|----|-------|---|-------|------------|---------------|-----------------------|------------|
| March 13 2018 | 23      | 2  | 2  | 2  | 3  | 1  | 8     | 1  | 3  | 0  | 4     | 1 | 15    | forest     | Thick forest  | 61.94069              | -110.22019 |
| March 13 2018 | 24      | 0  | 0  | 0  | 0  | 0  | 0     | 0  | 0  | 1  | 1     | 0 | 1     | forest     | Thick forest  | 62.16946              | -109.80263 |
| March 13 2018 | 25      | 0  | 0  | 0  | 0  | 0  | 0     | 0  | 0  | 3  | 3     | 0 | 3     | forest     | Lake edge     | 62.66915              | -109.42112 |
| March 14 2018 | 26      | 0  | 0  | 0  | 0  | 0  | 0     | 0  | 0  | 3  | 3     | 0 | 3     | forest     | Rocky ridge   | 62.35855              | -108.78146 |
| March 14 2018 | 27      | 4  | 2  | 0  | 2  | 2  | 6     | 2  | 0  | 3  | 5     | 2 | 17    | forest     | Sparse forest | 62.36656              | -108.77311 |
| March 14 2018 | 28      | 5  | 3  | 2  | 3  | 0  | 8     | 2  | 1  | 3  | 6     | 1 | 20    | transition | Rocky ridge   | 62.16912              | -108.57919 |
| March 14 2018 | 29      | 0  | 0  | 0  | 0  | 0  | 0     | 0  | 0  | 2  | 2     | 0 | 2     | forest     | Thick forest  | 61.56935              | -109.25533 |
| March 14 2018 | 30      | 0  | 0  | 0  | 0  | 0  | 0     | 0  | 0  | 2  | 2     | 0 | 2     | forest     | Thick forest  | 61.67321              | -109.15426 |
| March 14 2018 | 31      | 2  | 2  | 0  | 2  | 0  | 4     | 3  | 1  | 1  | 5     | 1 | 12    | transition | Sparse forest | 61.76283              | -108.83995 |
| March 14 2018 | 32      | 2  | 0  | 2  | 1  | 1  | 4     | 0  | 0  | 3  | 3     | 0 | 9     | transition | Sparse forest | 61.74727              | -109.24565 |
| March 14 2018 | 33      | 4  | 4  | 1  | 8  | 2  | 15    | 3  | 2  | 1  | 6     | 1 | 26    | forest     | Thick forest  | 62.21467              | -110.25206 |
| March 16 2018 | 34      | 5  | 3  | 1  | 2  | 0  | 6     | 3  | 0  | 1  | 4     | 0 | 15    | forest     | Rocky ridge   | 61.95261              | -111.60884 |
| March 16 2018 | 35      | 3  | 1  | 1  | 2  | 1  | 5     | 2  | 2  | 0  | 4     | 0 | 12    | forest     | Rocky ridge   | No wpt; near group 36 |            |
| March 16 2018 | 36      | 8  | 3  | 6  | 3  | 0  | 12    | 4  | 5  | 2  | 11    | 6 | 37    | forest     | Rocky ridge   | 61.91464              | -111.74888 |
| March 16 2018 | 37      | 6  | 1  | 1  | 0  | 0  | 2     | 1  | 2  | 0  | 3     | 0 | 11    | forest     | Rocky ridge   | 61.92572              | -111.72383 |
| March 16 2018 | 38      | 3  | 2  | 0  | 0  | 1  | 3     | 2  | 4  | 3  | 9     | 1 | 16    | forest     | Rocky ridge   | 61.91447              | -111.66896 |
| March 16 2018 | 39      | 0  | 0  | 0  | 0  | 0  | 0     | 0  | 0  | 1  | 1     | 0 | 1     | forest     | Rocky ridge   | 61.76171              | -111.81396 |
| March 16 2018 | 40      | 3  | 2  | 0  | 5  | 1  | 8     | 2  | 4  | 3  | 9     | 3 | 23    | forest     | Lake edge     | 61.74919              | -111.268   |
| March 16 2018 | 41      | 0  | 0  | 0  | 0  | 0  | 0     | 0  | 0  | 1  | 1     | 0 | 1     | forest     | Sparse forest | 61.75221              | -111.10798 |
| March 16 2018 | 42      | 0  | 0  | 0  | 1  | 0  | 1     | 0  | 0  | 5  | 5     | 0 | 6     | forest     | Lake edge     | 61.56901              | -110.9265  |
| March 16 2018 | 43      | 0  | 0  | 0  | 0  | 0  | 0     | 0  | 0  | 1  | 1     | 0 | 1     | forest     | Rocky ridge   | 61.63697              | -112.1234  |
| March 16 2018 | 44      | 3  | 2  | 0  | 1  | 1  | 4     | 3  | 2  | 0  | 5     | 3 | 15    | forest     | Lake edge     | 61.68465              | -110.84001 |
| March 16 2018 | 45      | 8  | 6  | 1  | 4  | 0  | 11    | 2  | 6  | 0  | 8     | 3 | 30    | forest     | Rocky ridge   | No wpt; near group 38 |            |
| March 19 2018 | 46      | 10 | 5  | 4  | 8  | 0  | 17    | 0  | 4  | 5  | 9     | 0 | 36    | forest     | Lake edge     | 61.83735              | -109.27361 |
| March 19 2018 | 47      | 5  | 4  | 5  | 5  | 0  | 14    | 1  | 0  | 3  | 4     | 1 | 24    | forest     | Rocky ridge   | 61.82033              | -109.72696 |
| April 7 2018  | 48      | 12 | 5  | 7  | 4  | 0  | 16    | 6  | 1  | 4  | 11    | 0 | 39    | tundra     | Tundra        | 63.21617              | -109.88883 |
| April 7 2018  | 49      | 13 | 5  | 2  | 3  | 1  | 11    | 6  | 2  | 0  | 8     | 1 | 33    | tundra     | Tundra        | 63.46267              | -109.71757 |
| April 7 2018  | 50      | 12 | 7  | 2  | 4  | 1  | 14    | 5  | 3  | 0  | 8     | 0 | 34    | tundra     | Tundra        | No wpt; near group 49 |            |
| April 7 2018  | 51      | 0  | 0  | 0  | 0  | 0  | 0     | 0  | 0  | 8  | 8     | 0 | 8     | tundra     | Tundra        | 63.47253              | -108.76001 |

| <b>Date</b>   | <b>Group #</b> | <b>C</b> | <b>F2</b> | <b>F3</b> | <b>F4</b> | <b>UF</b> | <b>F All</b> | <b>M2</b> | <b>M3</b> | <b>M4</b> | <b>M All</b> | <b>U</b> | <b>Total</b> | <b>Habitat</b> | <b>Fine Habitat</b> | <b>Latitude</b> | <b>Longitude</b> |
|---------------|----------------|----------|-----------|-----------|-----------|-----------|--------------|-----------|-----------|-----------|--------------|----------|--------------|----------------|---------------------|-----------------|------------------|
| April 7 2018  | 52             | 10       | 0         | 5         | 10        | 0         | 15           | 2         | 2         | 4         | 8            | 0        | 33           | tundra         | Tundra              | 63.34973        | -108.34719       |
| April 7 2018  | 53             | 2        | 3         | 5         | 7         | 0         | 15           | 0         | 0         | 5         | 5            | 0        | 22           | tundra         | Tundra              | 62.8285         | -107.13645       |
| April 7 2018  | 54             | 1        | 3         | 1         | 3         | 0         | 7            | 1         | 0         | 0         | 1            | 0        | 9            | transition     | Sparse forest       | 62.80488        | -107.96432       |
| April 7 2018  | 55             | 10       | 3         | 1         | 8         | 2         | 14           | 5         | 1         | 1         | 7            | 3        | 34           | transition     | Rocky ridge         | 62.84955        | -109.01676       |
| April 19 2018 | 56             | 0        | 0         | 0         | 0         | 0         | 0            | 0         | 0         | 7         | 7            | 0        | 7            | forest         | Sparse forest       | 61.91782        | -109.80638       |
| <b>Totals</b> |                | 215      | 87        | 68        | 148       | 34        | 337          | 88        | 72        | 122       | 282          | 57       | 869          |                |                     |                 |                  |



**APPENDIX 2. SELECTED MUSKOX PHOTOS (MIXED GROUPS)**

Photo credit: K. Olesen















### APPENDIX 3: SELECTED MUSKOX PHOTOS (BULL-ONLY GROUPS)

Photo credit: K. Olesen





