Compendium of Research in the Northwest Territories **2011**




This publication is a collaboration between the Aurora Research Institute, the Department of Environment and Natural Resources, Fisheries and Oceans Canada and the Prince of Wales Northern Heritage Centre. Thank you to all who submitted a summary of research or photographs, and helped make this publication possible.

Editor: Wayne Condon, Jolie Gareis, Niccole Hammer, Erika Hille, Ashley Mercer, Jonathon Michel, Sarah Rosolen, and Pippa Seccombe-Hett

Copyright © 2014 ISSN: 1205-3910

Printed in Fort Smith through the Aurora Research Institute





Pêches et Océans





FOREWORD

The 2011-2012 Compendium of Research in the Northwest Territories provides a brief summary of most research that has taken place in the Northwest Territories. The rich and continuing tradition of research in the NWT is showcased in this publication. Over the past two years, researchers traveled from near and far to observe, record and analyse our dynamic region. The results of this work have contributed to an ever-growing body of knowledge on topics ranging from aboriginal languages evolution to climate change impacts, from ancestral ways of life to regional health management. The breadth and depth of this research is truly extraordinary.

The NWT has long been an important location for research. This year, however, I am struck by the growth in research from local community researchers and NWT citizens on their own regions and heritage. This promising trend will benefit this territory as we move towards increasingly more independent governance. The NWT's growing research capacity will be important to help direct the understanding of our home and inform our future decision-making.

In September 2011, the NWT celebrated the opening of the Western Arctic Research Centre in Inuvik. This new facility was supported through the Arctic Research Infrastructure Fund, administered by Aboriginal Affairs and Northern Development Canada, and was one of twenty funded projects in northern Canada. The Western Arctic Research Centre replaces the old Inuvik Research Centre after 47 years of continuous operation. The new state-of-the-art facility will expand research programming and research opportunities for the NWT. This renewed investment in research infrastructure will continue to support and build upon the rich and diverse range of research activity in the NWT.

The summaries in this publication are only a brief outline of the rich findings and scientific advancements researchers have made over the past year. In many cases, more in-depth reports and publication are available. I encourage you to contact the researchers if there is a project that interests you. Additional information can be found at: http://data.nwtresearch.com/.

Pippa Seccombe-Hett
Director, Aurora Research Institute

TABLE OF CONTENTS

Introduction	iii
Aurora Research Institute	vi
The Department of Environment & Natural Resources	vii
Department of Fisheries and Oceans	viii
Prince of Wales Northern Heritage Centre	ix
2011 Licenced Research Projects	
Biology	
Contaminants	14
Engineering	18
Health	20
Physical Sciences	24
Social Sciences	56
Traditional Knowledge	73
Archaeology	78
Wildlife	86
Fisheries Permits	97
2012 Licenced Research Projects	
Biology Licences	
Contaminants Licences	
Engineering Licences	
Health Licences	
Physical Sciences Licences	
Social Sciences Licences	
Traditional Knowledge Licences	
Archaeology Permits	196
Wildlife Permits	206
Fisheries Permits	
Glossary	
Author Index	241

INTRODUCTION

This compendium offers a summary of research licences/permits that were issued in the Northwest Territories during 2011 and 2012. The information contained in this book is a product of a collaboration between the Aurora Research Institute (ARI), the Prince of Wales Northern Heritage Centre (PWNHC), the Department of Environment and Natural Resources (ENR) and the Department of Fisheries and Oceans (DFO). The Compendium series began in 1986.

Licensing in the NWT

Under territorial legislation, all research in the NWT requires a licence/permit from one of four agencies, depending on the type of research being conducted:

- Prince of Wales Northern Heritage Centre Archaeology;
- Department of Environment and Natural Resources, Government of the Northwest Territories - Wildlife;
- Department of Fisheries and Oceans Fisheries; or
- Aurora Research Institute all other research in the NWT.

Through the licensing process, researchers are informed of appropriate organizations, communities and other licensing/permitting agencies that should be contacted prior to conducting studies. Licensing ensures research activities are communicated to interested parties and provides opportunities for the exchange of information.

The Compendium provides a summary of all licences/permits issued in the NWT by all four licensing/permitting bodies. As each research project is represented by a short abstract, the reader is encouraged to contact the researcher for additional information and results.

How to Use This Book

This book has four main sections. Each of these sections reflects a specific licensing agency and type of licence/permit issued. Within each section, research descriptions have been grouped by subject and listed alphanumerically by the principal researcher's last name. Refer to the Table of Contents for the specific page on which each section and/or subject begins. An index is included at the end of the compendium listing all researchers in each section.

1. File Number

The file numbers shown in each of the Aurora Research Institute's subject areas refer to the file number issued to a particular researcher. It allows cross referencing with research material that may be available on file or in the ARI library. The reference numbers of the other three agencies refer directly to the permit numbers given to each researcher. When requesting information from any of these agencies on specific research outlined in the compendium, please refer to the reference number in your correspondence.

2. Regional Abbreviations

Throughout the book, reference is given to the specific land claim regions in which the research took place. The regions are shown on the following page. Some of the land claim regions are still under negotiation and the boundaries shown are only approximations. The abbreviations shown for each region are as follows:

DC	Dehcho	SS	South Slave
NS	North Slave	SA	Sahtú Settlement Area
IN	Inuvialuit Settlement	GW	Gwich'in Settlement Area
	Region		

3. Glossary

A glossary of terms has been added to the Compendium. The intent of the glossary is to allow the reader to better appreciate the research descriptions.

Available in Print or Free Download

This compendium is available as a printed publication or can be downloaded from the Aurora Research Institute's website (www.nwtresearch.com). Copies can also be requested by contacting the Aurora Research Institute.

Send Us Your Comments

Whether you are a researcher or an interested member of the public, the Aurora Research Institute welcomes your comments and suggestions concerning this publication. Contact us by mail, fax, email or telephone (see address on page vi).




Figure 1. Land claim regions in the Northwest Territories

AURORA RESEARCH INSTITUTE

The Aurora Research Institute's mandate is to improve the quality of life for NWT residents by applying scientific, technological and indigenous knowledge to solve northern problems and advance social and economic goals.

ARI is responsible for:

- licencing and coordinating research in accordance with the NWT Scientists Act: this
 covers all disciplines including the physical, social, biological sciences and traditional
 knowledge;
- promoting communication between researchers and the people of the communities in which they work;
- promoting public awareness of the importance of science, technology and indigenous knowledge;
- fostering a scientific community within the NWT which recognizes and uses the traditional knowledge of northern aboriginal people;
- making scientific and indigenous knowledge available to the people of the NWT;
- supporting or conducting research and technological developments which contribute to the social, cultural and economic prosperity of the people of the NWT.

For more information, contact ARI at:



Aurora Research Institute

PO Box 1450 Inuvik, NT X0E 0T0 Tel: 867-777-3298 Fax: 867-777-4264

E-mail: <u>licence@nwtresearch.com</u>
Website:www.nwtresearch.com

DEPARTMENT OF ENVIRONMENT & NATURAL RESOURCES

The Government of the Northwest Territories' Department of Environment and Natural Resources (ENR) has a mandate to promote sustainable development through the management and protection of the quality, diversity and abundance of natural resources and the integrity of the environment.

With respect to permitting for research and monitoring, ENR is responsible for issuing Wildlife Research Permits under the Wildlife Act (Section 24) for all studies on wildlife or wildlife habitat in the Northwest Territories. Wildlife includes all vertebrates, except fish and marine mammals.

For more information, contact ENR at:

Wildlife Division

Environment and Natural Resources Government of the Northwest Territories PO Box 1320 Yellowknife. NT X1A 2L9

Fax: 867-873-0293

Email: wildliferesearch permit@gov.nt.ca

Website: www.nwtwildlife.com/ResearchPermits/



DEPARTMENT OF FISHERIES AND OCEANS

The Department of Fisheries and Oceans Canada (DFO) is responsible for developing and implementing policies and programs in support of Canada's scientific, ecological, social and economic interests in oceans and fresh waters. Some Fisheries management responsibilities have been delegated or transferred to other federal agencies (e.g. Parks Canada), provinces/territories and co-management groups under Land Claim agreements.

DFO Fisheries Management is responsible for issuing Commercial, Domestic, Licence to Fish for Scientific Purposes (LFSP), Exploratory, Public Display and Educational licences in the NWT. Subject to Land Claim agreements, a Commercial licence is required to sell or barter fish

All individuals fishing for scientific purposes or participating in the acts described below are required to obtain a Licence to Fish for Scientific Purposes (LFSP):

- activities involving fishing, catching or attempting to catch fish;
- activities where the potential exists for the incidental capture of fish;
- sampling or possessing fish caught in a subsistence fishery.

For further information about licensing, contact DFO at:

Licensing Officer

Central & Arctic Region Government of Canada Fisheries and Oceans Canada PO Box 1871

Inuvik, NT X0E 0T0 Tel: (867) 777-7500 Fax: (867) 777-7501

Email: xca-inuvikpermit@dfo-mpo.gc.ca
Website: www.dfo-mpo.gc.ca/index-eng.htm

PRINCE OF WALES NORTHERN HERITAGE CENTRE

The Prince of Wales Northern Heritage Centre (PWNHC), a division of the Department of Education, Culture and Employment, Government of the Northwest Territories, is responsible for managing and protecting the archaeological resources of the NWT. Representing a continuous human occupation stretching back over 7000 years, archaeological sites are fragile and non-renewable and are protected from disturbance by legislation, regulation and policy in the NWT. There are currently about 6000 archaeological sites recorded in the NWT, though this number represents only a fraction of the existing sites as large areas remain unexplored for archaeological resources. A large part of the work done at the PWNHC involves reviewing land use and development permit applications. On average, 300 permits are reviewed per year, with recommendations being proffered to nine land management authorities.

With respect to permitting for research and monitoring, PWNHC is responsible for issuing NWT Archaeology Research Permits.

For more information, contact the Prince of Wales Northern Heritage Centre at:

NWT Cultural Places Program
Prince of Wales Northern Heritage Centre
4750 48th Street
PO Box 1320

Yellowknife, NT X1A 2L9 Phone: (867) 873-7551 Fax: (867) 873-0205

Email: archaeology@gov.nt.ca

Website: www.pwnhc.ca





BIOLOGY 2011

Blaschuk, Katherine

Imperial Oil Limited Calgary, AB katherine.a.blaschuk@esso.ca

File No: 12 402 842 **Licence No:** 14847

Region: SA **Location:** Bosworth Creek; Norman Wells

Bosworth Creek aquatics and fisheries monitoring program Gahcho Kué

The objective of the Bosworth Creek aquatics and fisheries monitoring program was to determine current water and aquatic habitat quality, and to establish baseline conditions for evaluation of future monitoring data. In order to achieve the project objectives, sampling stations within and downstream of the active Imperial Oil Norman Wells lease area were compared to an upstream reference station, located beyond the possible influence of the development. Field monitoring was conducted on August 31 and September 1, 2011. The 2011 program included the following components: (1) collection and analysis of surface water quality samples from four upstream and three downstream locations; (2) Assessment of the benthic invertebrate community at six locations coincident with the surface water quality sampling; and (3) assessment of fish habitat quality at two upstream and two downstream locations. Please note that fish sampling was not conducted in 2011. Analysis of data collected in 2011, and preparation of a summary report, is currently underway.

Buddle, Chris

McGill University Ste-Anne-de-Bellevue, QB chris.buddle@mcgill.ca

File No: 12 402 861 Licence No: 14914

Region: IN, SA, NS **Location:** Norman Wells; Yellowknife; Aulavik National

Park on North East Banks Island

Ecological structure of northern arthropods: adaptation to a changing environment

The research team visited three areas in the Northwest Territories: Yellowknife (June 6 to 21), Norman Wells (June 6 to 19), and Aulavik National Park on Banks Island (July 6 to 22). In all three locations they conducted their standardized sampling of insects and spiders at six sites, as well as some opportunistic collection in a variety of both terrestrial and aquatic habitats. Collections in all three locations were highly productive, especially for biting flies, higher flies, spiders, and parasitoid wasps. Team members are currently processing the samples in their labs at McGill University and the University of Toronto. Specimens in groups of interest are being sorted, dried, pinned, labeled, and identified. Data will be analyzed to test how, and to what degree, the structure

of arthropod communities change between boreal, subarctic and arctic ecoclimatic zones, and to assess how northern arthropod communities have adapted to recent (50-60 year) changes to their environment. Cutting-edge molecular genetic techniques will aid species identification and help elucidate longer-term (i.e., phylogeographical) patterns. In addition to their collection-based activities, the research team also participated in public education activities as bug experts at "Bugfest at the Museum" in Yellowknife.

Budziak, Jerry

Seaway Energy Services Inc.
Calgary, AB
jerry.budziak@seawayenergy.com

File No: 12 402 475 **Licence No:** 14890

Region: SA **Location:** Nota Creek C-17 well site

Phytoremediation study on the Canadian forest et al. Nota Creek C-17 wellsite

Phytoremediation is a remediation strategy involving the use of plants to remove contaminants. In theory, plants take up the contaminant from the soil, are harvested and then removed from the site. This process is repeated until the impacted soil is remediated to applicable guidelines. Phytoremediation activities on the Nota Creek C-17 well site progressed to full site planting in 2009 and 2010. Remediation results were encouraging enough to support excavating impacted soil still buried on the site and integrating it into the phytoremediation process. Personnel and equipment were mobilized to the site, approximately 43 km southeast of Norman Wells and 38 km northwest of Tulít'a, in February/March 2011 under frozen ground conditions. Some of the remaining impacted soil that was still buried was excavated and stockpiled. In early July 2011, a portion of the excavated impacted material was incorporated into the phytoremediation process. The soil was spread and conditioned, and the site fertilized and seeded. Plant health and vigor was assessed in mid-August during a monitoring trip. In late September personnel were mobilized to the site to collect plant and soil samples, to harvest the growth from the impacted areas and to remove that growth from the site. Initial laboratory results from the collected samples were encouraging and support continuation of the application of phytoremediation technology to the well site.

Carriere, Suzanne

Environment and Natural Resources Yellowknife, NT suzanne carriere@gov.nt.ca

File No: 12 402 858 Licence No: 14901

Region: IN Location: The Baillie Islands and northwestern portions of

Cape Bathurst

Population inventory of hairy rockcress (*Braya pilosa*): NWT's rarest species of global importance

Hairy rockcress, a plant first collected by Sir Richardson in the mid-1800s, is known from only one area in the world; Cape Bathurst on the Beaufort Sea coast. Specimens from that site were collected more than 200 years ago, and then thought to be lost until 2004 when the exact location was re-discovered. The objectives of the 2011 field trip were to determine how many populations of this rare plant grow on Cape Bathurst, and to describe and evaluate the threats to its survival. Part of the team arrived by airplane on July 25, set up camp and walked to and along the coast. New locations for this plant were found. The major threat to this rare plant is shoreline erosion. For example, the 2004 site was re-examined and found to be almost all gone due to ground

slumping into the sea and salt kills. On August 1, additional team members arrived with a helicopter and a large area of Cape Bathurst was searched for one day. The whole team left the Cape on August 2, 2011. In all, about 15,000-20,000 plants were observed in 10 locations, some of which are protected from coastal erosion for now. The results of this survey are being used to assess the biological status of the rare plant in reports for the Committee on Endangered Wildlife in Canada and for the NWT Species at Risk Committee.

Cote, Jason

Cambria Marshall Cote Ltd. Yellowknife, NT jcote@cambria-nwt.com

File No: 12 402 263 **Licence No:** 14975

Region: NS **Location:** The La Martre River, extending from the La

Martre River Falls to the confluence with the Marian River

La Martre River fall fisheries program

In 2009, the Tłycho Investment Corporation proceeded with a pre-feasibility investigation of the La Martre River Hydroelectric Project. The project is a 13.2 MW run-of-river power generation facility that proposes to divert 30 m³/s, or approximately 89% of the mean annual discharge, from a headworks facility on the La Martre River. In order to understand the potential effects of a sudden increase/decrease in flow on fish and fish habitat within the river, Cambria Marshall Côté consultants completed a preliminary assessment in 2011. The results indicated that there is a risk of stranding fish and/or dewatering incubating eggs; however, the study was conducted at a desktop level and no fisheries information was available downstream of the proposed tailrace. It was recommended that a fish and fish habitat study be conducted downstream from the proposed tailrace. Fieldwork was completed in September 2011 by Cambria Marshall Côté and a local technician from the community of Whatì. The study area was broken into two distinct river reaches: upper (tailrace to 8 km downstream) and lower (8 km to 34 km downstream of tailrace). A total of 42 sample sites were visited within the two reaches, and four habitat types were identified in each. A total of 159 fish from 9 species -- stickleback, sculpin, arctic grayling, northern pike, sucker spp., lake chub, burbot, emerald shiner, and trout perch – were captured within the upper section. A braided side-channel habitat was found to provide high value habitat to a number of species and potentially critical habitat for arctic grayling. A total of 129 fish from 7 species stickleback, sculpin, northern pike, sucker spp., lake chub, inconnu, and emerald shiner -- were captured within the lower section. Run/glide and deep pool habitats in this section were documented to provide excellent rearing potential, and based on the depth and velocity characteristics, likely provide excellent overwintering potential.

Guthrie, Glen

Sahtú Renewable Resources Board Norman Wells, NT rrco@srrb.nt.ca

File No: 12 402 780 **Licence No:** 14841

Region: SA Location: Bosworth Creek

Bosworth Creek monitoring project

This Bosworth Creek Monitoring Project (BCMP) is a high resolution, long-term study of a 125 square kilometer watershed at Norman Wells, NT. The project was initiated after the Sahtú Renewable Resources Board received a request from local residents for information on fish stocks in Bosworth Creek, following the removal of a weir in 2005 by Imperil Oil Resources. The

project has focused on studying benthic insects and creating a chemical inventory over the past three years. The benthic work will be completed within one year and will compare Bosworth insects with three other nearby streams. The water quality analysis will be completed by September 2011. This information will be invaluable for local Renewable Resources Council stewardship. The project will continue to monitor potential or existing impacts by climate change and industry. The BCMP has become a permanent component of Mackenzie Mountain School's high school curriculum program through the NWT Experiential Science Program. The BCMP will continue to provide professional development for local youth through associations with academic and industrial institutions.

Guthrie, Glen

Sahtú Renewable Resources Board Norman Wells, NT rrco@srrb.nt.ca

File No: 12 402 780 **Licence No:** 14873

Region: SA **Location:** Lac St. Therese; Kelly Lake; Lennie Lake;

Stewart Lake; Tate Lake; Hodgson Lake

Baseline mercury levels in predatory fish in the Sahtú Settlement Area

No research was conducted under this licence in 2011.

Johnson, Mary Ann

AMEC Earth and Environmental Calgary, AB maryann.johnson@amec.com

File No: 12 404 771 Licence No: 14938

Region: SA **Location:** Along the proposed Highway Route within the

K'ahsho Got'ine District from Gibson Gap to the Thunder

River

Mackenzie Valley Highway project description report

Environmental fieldwork was conducted along the proposed K'ahsho Got'ine Highway (KGH). This fieldwork was undertaken by a fisheries biologist and ecologist with AMEC Environment and Infrastructure. The proposed highway would upgrade the winter road from Norman Wells to Fort Good Hope into a year round road and construct a new road northwards to the Gwich'in Settlement Area boundary. The goal of this fieldwork was to verify the environmental information that was available for potential fish and wildlife habitat, as well as plant communities along the proposed highway. The fieldwork was conducted from July 12th to 16th, 2011. The field work was planned to use a helicopter to assess watercourse crossings with potential fish habitat and portions of the route with limited environmental information. However, due to issues with the helicopter, part of the proposed highway near Fort Good Hope was accessed by hiking and by boat. Notes and photographs were taken on the presence or absence of fish habitat and plant communities at each watercourse crossing. In addition, notes were made on the suitability of wildlife habitat. The final project description report was delivered to the K'ahsho Development Foundation in January 2012.

Jones, Paul

University of Saskatchewan paul.jones@usak.ca

File No: 12 402 867 Licence No: 14970

Region: SS **Location:** The mouth of the Slave River Delta near Fort

Resolution; the Slave River within the municipal

boundaries of Fort Smith

Fish health study in the Slave River and the Slave River Delta

The aim of this study is to investigate the contamination of the Athabasca/Slave river system with chemicals potentially derived from oilsands activities in northern Alberta. Contamination effects on fish health, and the quality of those fish as food, were examined. The potential for contaminants effects on the health of fishes in the Athabasca River was identified previously from measures of chemicals in water and snowmelt, and there have been reports in the press of the occurrence of fish with lesions. Furthermore, observations from local harvesters suggest that similar health impacts may be happening to fish from the Slave River system, which is immediately downstream of the Athabasca system. Potential impacts on fisheries were assessed by collecting four species of fishes from several locations in the Athabasca/Slave River system. Samples were collected at Ft. McMurray AB (upstream of oilsands activities), Ft. McKay AB (immediately downstream of oilsands activities), Ft. Chipewyan AB (where the Athabasca River enters Lake Athabasca), Ft. Smith NT (on the Slave River downstream of Lake Athabasca) and Ft. Resolution NT (where the Slave River empties into Great Slave Lake). Fish were collected in collaboration with local harvesters and other community members, aboriginal organizations, provincial or territorial authorities and Federal agencies as appropriate. The health assessment did not indicate any major observable differences in fish health between sites, however this data is still being compiled for statistical analysis, which cannot be conducted until all sampling is complete in the spring of 2012. Samples have been submitted to laboratories for analysis of metal and organic chemical content. While preliminary results for metals have been provided they have not yet been subjected to final quality assurance/quality control (QA/QC) assessment, and so cannot be released.

Krizan, Julia

IMG-Golder Corporation Inuvik, NT jkrizan@golder.com

File No: 12 402 664 **Licence No:** 14960

Region: IN Location: Along the proposed Tuktoyaktuk to Inuvik

highway

Archaeological and fish habitat assessment for the Tuktoyaktuk to Inuvik highway

IMG-Golder Corporation completed an archaeological and fish habitat assessment on behalf of the Government of the Northwest Territories Department of Transportation, as required for the proposed Inuvik to Tuktoyaktuk highway and two potential realignments (Alternative 1 and Alternative 3) in the Inuvialuit Settlement Region. The objectives of the archaeological assessment were to identify, record and assess cultural heritage resources that might be impacted by the proposed highway project, and to devise mitigation strategies should any be found in conflict with the proposed highway. A field investigation was completed in September 2011 over a six day period, along the planned highway right-of-way and at several proposed borrow source locations. No artifacts were found and no new sites were recorded. The objective of the fish habitat assessment was to assess the 36 watercourse crossing locations of the proposed highway and potential realignments that remained following the 2009 and 2010 surveys. Site assessments on the proposed watercourse crossings were conducted in September 2011. A total of 11 watercourses were assessed as being ephemeral with no defined channel, 10 were

classified as intermittent, and 15 were classified as perennial with a defined channel. One crossing could not be located.

Langhorne, Amy

Golder Associates Ltd.
Saskatoon, SK
amy_langhorne@golder.com

File No: 12 404 733 **Licence No:** 14962

Region: NS Location: The Kennady Lake watershed and adjacent

watershed areas

De Beers - Gahcho Kué ecological risk assessment program

Collections of soil, berries, and vegetation were completed in Sept 2011 at the Gahcho Kué project. Up to 10 samples each of lichen, berries (cranberries), leaves (dwarf birch), and grass were collected among the 23 soil sample sites. Samples were sent to a lab to measure metal and polycyclic aromatic hydrocarbon levels. Results will be used to refine the environmental assessment for the project.

Lantz, Trevor University of Victoria Victoria, BC

tlantz@uvic.ca

File No: 12 404 758 **Licence No:** 14929

Region: IN **Location:** Peel Plateau; Aklavik Area; Delta Uplands;

Outer Delta; Aklavik / Western Delta

Vegetation monitoring and science training in the Mackenzie Delta region

From 2010-2012, AANDC scientists have been working with researchers at the University of Victoria, and Hunters and Trappers Committees (HTCs) in the Mackenzie Delta, to develop a vegetation monitoring protocol that can be implemented by a range of users. The long-term goal of this program is to establish and maintain a network of sites to characterize regional environmental variability, and serve as a baseline against which to measure changes resulting from the cumulative impacts of multiple natural and anthropogenic disturbances. At all sites in the network, the researchers measure vegetation structure, plant community composition, tree density, the productivity of edible berries, active layer depth, and near surface ground temperatures. At core sites they also maintain meteorological stations, frost tubes, and deep ground temperature cables. Since 2010, they have established 35 community-based monitoring sites in 6 terrain types. Statistical power simulations, using data from 2010-2012, also show that the protocol is capable of detecting small changes in vegetation structure. Through a partnership with Environment Canada and the Canadian Forest Service, they are working to identify areas of overlap among this and other programs and, where possible, develop common monitoring techniques. By selecting monitoring sites that are regionally representative, and including disturbances that communities have identified as priorities, this monitoring program also provides a platform for directed research and hypotheses-driven investigations, that will contribute to local decision making. The researchers are in the final stages of production of a user-friendly protocol guidebook. They anticipate completion and first printing in advance of the 2013 field season. Baseline data on vegetation, active layer, and snow is being added to the NWT Discovery Portal on an ongoing basis. In February 2012, a presentation describing project outcomes was given to

their community partners in Inuvik, as well as several project presentations were also delivered in Yellowknife.

Lawson, Nick
Det'on Cho Stantec
Yellowknife, NT
nick.lawson@stantec.com

File No: 12 402 685 **Licence No:** 14945

Region: NS,SS **Location:** Within Avalon's Thor Lake Property; Redemption Lake, just northeast of Thor Lake

2011 baseline studies for Avalon Rare Metals Inc. proposed Thor Lake rare earth element project - aquatics and fisheries

In 2011, Deton'Cho Stantec conducted three field programs for the aquatics component of this project. This included: water and plankton (phytoplankton and zooplankton) sampling in June; water, sediment, plankton and benthic invertebrate sampling in September; and water sampling in October. Sampling was carried out at 8 lake stations. Results from the 2009-2011 field programs indicate neutral to basic water and very low nutrient levels at all stations. There were large fluctuations in some general and metal parameters, primarily during winter in small, shallow lakes that developed highly reducing, anoxic conditions under ice. Sediment characteristics varied, though generally lake sediment had high phosphorus, nitrogen and organic carbon content; metal levels in sediment ranged from less than detection to higher than the guidelines set by the Canadian Council of Ministers of the Environment. Chlorophyll levels varied among lakes and seasons, and most of the lakes were oligotrophic. From 2009-2011, phytoplankton and zooplankton abundance, richness and diversity varied between lakes and years. Predominant phytoplankton species included filamentous and coccoid blue-green algae, colonial yellow-brown algae and small cryptoflagellates. One rotifer species was the predominant zooplankton taxa in most lakes from 2009-2011. Similarly, benthic invertebrate abundance, richness, diversity and evenness varied between lakes and years. One dipteran family (Chironomidae), one crustacean order (Amphipoda) and one clam family (Sphaeriidae) were found in most lakes. The fisheries component of the 2011 Thor Lake baseline study included a fish sampling program at five lakes (Ring, Buck, Drizzle, Ball and Murky). A total of three juvenile northern pike were captured; two from Drizzle Lake and one from Murky Lake. No fish were caught in Ring, Ball or Buck Lakes. These results support the conclusions from field studies in 2009-2010: Drizzle and Murky Lakes are fish-bearing while Ring, Ball and Buck Lakes are non-fish bearing.

Lennie-Misgeld, Peter NWT Hydro Corporation Yellowknife, NT plennie-misgeld@ntpc.com

File No: 12 402 856 **Licence No:** 14887

Region: SA **Location:** The Great Bear River from Great Bear Lake to

the Mackenzie River, near Déline and Tulít'a

Great Bear River environmental and traditional knowledge baseline program

A field program was conducted in August 2011 with participation from both Tulít'a and DélĮnę. Five sites were identified on the Great Bear River, and data were collected on the channel structure, sediment, water chemistry, fish community and habitat, benthic invertebrates, and other environmental disciplines. Data is still being analyzed, but the following points provide a brief summary of field program results: Fourteen species of fish were identified; the average age of the

fish caught using gillnets was 8 years; the oldest fish caught was a 15 year old walleye; the mercury level in the water at all sites was very low; the creeks flowing into the Great Bear River have different water chemistry. The traditional knowledge (TK) portion of this study is underway at this time, and residents of both Tulít'a and DélĮnę are involved. The focus of the TK portion of the study is to better understand conditions of the river, including how the river is used, freeze/thaw patterns, locations of important cultural sites and key hunting and fishing areas. TK information will be documented by conducting workshops and interviews with elders, renewable resources councils, land users and residents from Tulít'a and DélĮnę.

Leski, Michael Buffalo Grove, IL United States peterlep28@yahoo.com

File No: 12 402 859 **Licence No:** 14902

Region: GW **Location:** Richardson Mountains near Husky Channel;

area south of Inuvik to Fort McPherson

Reassessment of the *Polygonia faunus* complex

The primary objective of this project was to collect a species of butterfly (*Polygonia faunus arcticus*) from sites in the Yukon and Northwest Territories, with a particular emphasis placed on obtaining specimens from Black Mountain in the Richardson Mountains, the type locality of this subspecies. These specimens supported a taxonomic assessment of the *Polygonia faunus arcticus*. Genetic analysis indicates that subspecies *arcticus* is synonymous with subspecies *faunus*, while subspecies *hylas* contains two additional as yet undescribed subspecies, and at least two independent expansion phases of *Polygonia faunus* into eastern North America have occurred. The results of this research have been submitted to BMC Evolutionary Biology, a peerreviewed scientific journal of entymology, and a trip report was published in the News of the Lepidopterist Society. A secondary objective of this study was to assess other butterfly species from this region. Initial evaluation of *Polygonia gracilis* from the Richardson Mountains demonstrates that the dorsal surface of these individuals resembles that of the eastern subspecies *Polygonia gracilis gracilis*. Thus, the Richardson Mountains are a blend area for *Polygonia gracilis*.

Low, George

Dehcho First Nations Hay River, NT geobarbgeo@hotmail.com

File No: 12 402 857 **Licence No:** 14875

Region: DC **Location:** Fort Simpson - Cli Lake, Little Doctor, Sibbeston Lake, Tsesto lake; Wrigley - Blackwater Lake, Fish Lake;

Jean Marie River - McGill Lake, Deep Lake; Trout Lake -

Trout Lake

Updating data on mercury levels in food fish species in lakes used by Dehcho communities In 2011, the Dehcho Aboriginal Aquatic Resources and Ocean Management (AAROM) program carried out the first year of this multiyear project. Work was done on five lakes with the assistance of local community monitors, who were critical to completing work in the field. Fish were collected and sampled, and water quality samples were taken. In the Jean Marie River First Nation area, fish and data were collected from Deep, McGill and Ekali Lakes. In total 5 yellow walleye, 4

northern pike, 3 burbot and one lake whitefish were caught in 13 sets at Deep Lake; 33 yellow walleye, 19 northern pike, 12 white sucker and 3 lake whitefish in 5 sets at McGill Lake; and 18 yellow walleye, 16 northern pike and 35 lake whitefish in 6 sets at Ekali Lake. In the Pehdzeh Ki First Nation (Wrigley) area, fish and data were collected from Fish Lake. In total 5 yellow walleye, 20 northern pike, 10 lake trout, 7 burbot and 25 lake whitefish were caught in 20 sets. Twenty-five lake trout were also collected from Trout Lake by residents of Sambaa K'e, and fish were sampled in Hay River. All of the fish collected during this project were biologically sampled in the field for fork length, total length, round weight, ageing structures, stomach contents, sex/maturity and gonad weight. If it was possible, fish were distributed to each local community afterwards.

Maier, Kris

Gwich'in Renewable Resources Board Inuvik, NT kmaier@grrb.nt.ca

File No: 12 402 851 **Licence No:** 14952

Region: GW Location: Stony Creek (NT); Vittrekwa River (Yukon)

Dolly Varden char assessment in the Peel Plateau

This project had two parts. The first was to re-assess the population size of the Vittrekwa River Dolly Varden char stock. The second was to determine if Dolly Varden char exist in the Stony Creek watershed near Fort McPherson. The project was carried out in August 2011. A team of three flew to the upper Vittrekwa River by helicopter and installed a fish weir on the spawning creek used by Vittrekwa River char. The weir was in place for 19 days and a total of 44 mature adult char were captured. Another 19 mature adult char were captured by angling. All char were sampled for length and weight, while only 50 had a small piece of fin removed for genetic analysis. It is important to note that only adult, sexually mature char were sampled. The actual population size of this stock is larger than this number (63 char sampled) because immature char were not counted in this assessment, as they typically do not migrate to the spawning tributary. Also, during the early period after the weir was installed, it was not operating efficiently and missed capturing a large number of mature male char. Because of this, the number of char counted through the weir can only indicate an absolute minimum population size. Visual counts of the spawning area estimated approximately 140 mature adult char. In addition to fish sampling, 9 CABIN (Canadian Aquatic Biomonitoring Network) reference sites were established for comparison with streams affected by permafrost slumping in the lower Vittrekwa and Stony watersheds. Sampling in the Stony Creek watershed was not completed because of high water and unfavorable conditions. One burbot was accidentally captured in an invertebrate kick net and was released. Juvenile grayling were observed in off-channel habitat but were not sampled.

Panayi, Damian

Golder Associates Ltd. Yellowknife, NT damian_panayi@golder.com

File No: 12 402 848 Licence No: 14849

Region: NS Location: Yellowknife River between Prosperous Lake and

Bluefish Lake

NTPC Bluefish Hydro repairs

The objective of this study was to describe the aquatic environment in the Yellowknife River between Prosperous Lake and Bluefish Lake. In particular, efforts were made to document current fish use of the existing dam. The results will be used to determine the most suitable means of fish

habitat compensation, and to guide mitigation strategies during construction of the new dam for Bluefish Hydro. In October 2011, a four-day fisheries assessment was completed in Bluefish Lake. Gillnets and Gee minnow traps were deployed, both near the existing dam and at the inflow to Bluefish Lake. Most fish were released live, although some were sacrificed to document age, sex, and reproductive status. Additional bathymetry information was also collected in the vicinity of the existing dam, and lake shore substrate and water quality was documented in Bluefish Lake.

Osawa, Akira

Kyoto University, Graduate School of Agriculture Kyoto, Kyoto Japan aosawa@kais.kyoto-u.ac.jp

Region: GW, SS **Location:** Adjacent to and along Highway #5, between the

boundary of Wood Buffalo National Park west of Fort Smith and Angus Tower; Along Highway #5, between the Park boundary west of Fort Smith and the intersection with the

road leading to Thebacha Camp; near Inuvik

Structure, carbon dynamics, and silvichronology of boreal forests

Forest landscape may be going through changes in growth under a warming climate. The researchers are trying to evaluate that thought (or hypothesis) by measuring forest growth at present, and by estimating its history during the past 100 years. Tree rings have been extensively used. They have also measured the movement of carbon in forest ecosystems, including growth of fine roots in the soil. Previous research they did in the Northwest Territories indicated that the proportion of growth in fine roots to the whole amount of forest growth is large, so root growth needs to be studied seriously. The researchers continued to collect data in small (several square kilometers) forest plots near Inuvik and in about a dozen forests near Fort Smith this year to estimate forest growth. Remote sensing was also used. They have almost finished measuring tree rings from several hundred stem samples they collected last year. They will use this data to calculate forest growth history. They will continue to study the forest plots next year to improve the accuracy of their reconstructed forest history. Thanks to a new development this year, a research partnership between Japan and the USA for investigating arctic environments, the researchers are now considering extending their experience in the Canadian north to a study of the entire arctic environment by starting similar projects in Scandinavia, and by integrating old and new results of their research activities.

Robb. Tonia

Rescan Environmental Services Yellowknife, NT trobb@rescan.com

File No: 12 402 766 **Licence No:** 14850

Region: NS Location: Waterbodies located within the EKATI claim

block

EKATI aquatic monitoring program, 2009-2013

In 2011, seven monitoring projects were ongoing in the lakes and streams of the Koala, King-Cujo and Pigeon watersheds, where EKATI mine infrastructure are located. The objectives of the aquatic effects monitoring program (AEMP) and the Fay Lake monitoring program were to assess the current conditions in the lakes and streams of the Koala, King-Cujo and Pigeon watersheds,

in order to determine whether there have been any mine effects. The objective of the surveillance network monitoring program was to confirm EKATI's compliance with its water licenses. The assessments incorporate some or all of the following: meteorology, hydrology, water quality and physical limnology, sediment quality, phytoplankton, zooplankton, benthos and fish data. Data analyses for 2011 are currently being completed. A detailed review of the AEMP plan for 2010-2012 was submitted to the Wek'eezhii Land and Water Board in 2011. Fish populations in the Panda diversion were monitored for the 13th consecutive year. A compilation and analysis of this work is also in progress. Assessment of fish habitat created in Nero-Nema Stream was ongoing in 2011. The nitrate in situ treatment study is a mitigation strategy developed and implemented in 2010 to reduce nitrate concentrations in the Long Lake containment facility (LLCF). In 2011, the study continued to closely monitor the physical, chemical and biological environment in Cell D and Cell E of the LLCF, with minor modifications to fertilizer timing additions. Air quality was monitored using high volume air sampling, continuous ambient monitors and dust fall measurements as a part of the air quality monitoring program.

Tonn, William

University of Alberta Edmonton, AB bill.tonn@ualberta.ca

File No: 12 402 724 Licence No: 14891

Region: NS **Location:** Seven lakes, and their six outlet streams,

draining into Lac de Gras

Improving habitat connectivity to enhance productive capacity of arctic freshwater ecosystems

The Diavik Diamond Mine, located on Lac de Gras in the Northwest Territories, has proposed a habitat compensation project for nearby lake and stream systems. Headwater lake outlet streams at two sites will be modified to improve fish passage, and thus ecological "connectivity" among these headwater lakes and with Lac de Gras. One set of habitat manipulations occurred in fall 2011, while the second habitat project will occur during summer 2012. Sampling during the 2011 summer field season (Year 3 of the "before" period) was conducted according to plan. Combined with the two lake-stream ecosystems designated for habitat compensation, several other lakestream reference systems were studied to provide information on the spatial variability of aquatic ecosystems in the barren lands, and to incorporate natural temporal patterns in these ecosystems. The researchers sampled each lake and stream for hydrology, water quality, habitat characteristics, primary producers, invertebrates, and fish to establish pre-manipulation baseline conditions for these ecosystems. Electrofishing and hoop netting in streams revealed very low abundances of slimy sculpin and juvenile burbot. They collected Surber and drift samples of macroinvertebrates, in both riffle and pool habitats, from seven streams in the Lac de Gras watershed (4 slated for modification, 3 controls) during the ice-free season. Post-winter recolonization of streams by macroinvertebrates was measured by placing different colonization boxes, open to only one type of colonization (upstream, downstream, aerial, and vertical), in each stream. Samples are currently being analyzed. Lake fish assemblages were surveyed by gill netting, angling, and electrofishing. Fish communities and species abundances vary among lakes, but consist mainly of arctic grayling, lake trout, round whitefish, burbot, longnose sucker, and slimy sculpin. During this "before" period, the researchers are comparing the ecology (diet composition, condition, growth) of lake trout in lakes with other fishes, to that of lake trout in lakes without other fishes.

Trimble, Annika

Aurora Research Institute Inuvik, NT atrimble@auroracollege.nt.ca

File No: 12 402 842 Licence No: 14853 Region: IN, GW Location: Inuvik

Northern native seed development field trials

The objective of this project is to continue assessing the performance of native plants which were seeded and transplanted into field plots in 2006 and 2007. Data regarding winter survival, seedling emergence, and overall plant vigor and productivity will be collected in order to assess each species' and collection's suitability for use in land reclamation in the Northwest Territories (NT). The results of this study will help to determine which plant species are best suited for restoring disturbed sites to their natural condition in different habitats across the NT. This year data was recorded at all three sites. As in previous years, plants which had been transplanted had higher survival than those that were direct seeded. Survival was also higher at the gravel and clay sites than at the peat site. Data analysis is ongoing, and results will be made available at www.nwtresearch.com.

Turetsky, Merritt University of Guelph Guelph, ON

mrt@uoquelph.ca

File No: 12 402 864 **Licence No:** 14958

Region: DC, SS **Location:** The sites of six recent fires that occured south

of Yellowknife since 2004

Effects of wildfire on biomass combustion in boreal peatlands and forests

Study plots were established in several recent fire scars south of Yellowknife. Data were collected to estimate the depth of burning of fuels on the forest floor (moss and peat). These data will be used to explore ecosystem type and weather as controls on burn severity.

Venables, Chandra

University of Calgary Calgary, AB cvenable@ucalgary.ca

File No: 12 402 860 **Licence No:** 14903

Region: NS, SS **Location:** Daring Lake; ponds, lakes, and creeks in the

area of Fort Smith

A phylogenetic and phylogeographic study of predaceous diving beetles in the Nearctic with a focus on the tribe Agabini (Coleoptera, Dytiscidae, Colymbetinae)

The main purpose of this project is to look for information about how water beetles respond to a changing environment, including: how have they re-colonized northern Canada and Alaska after the retreat of glaciers at the end of the last ice age? Another goal is to examine where different types of beetles are found in the north, and to see if these beetles are expanding their ranges northward. During the summer of 2011 (from July to early August), beetles were collected within 5 km of Daring Lake. To date, the beetles have not been fully identified; however, so far there are between 5 and 7 species that, in comparison to where they were found 30-40 years ago, appear

to have shifted their ranges to the north. This information may be important because it tells the researchers how animals in the north may be responding to a changing environment. These, and future results, will be presented at scientific conferences and in scientific journals. They will also be provided to the GNWT, to be placed in the library so the public can access them.

Wilcockson, John

Hatfield Consultants North Vancouver, BC jwilcockson@hatfieldgroup.com

File No: 12 402 865

Region: DC

Licence No: 14961

Location: Prairie Creek

Exfiltration fence habitat assessment

A fisheries field assessment was conducted on August 10 and 11, 2011. The objective of the study was to assess fish utilization of a section of Prairie Creek that will contain the proposed mine effluent exfiltration trench (diffuser) and effluent mixing (dilution) zone. The study area consisted of a 100 m long zone of the creek. Electrofishing was conducted along both banks of the 100 m zone. Caught fish were identified, weighed, and measured for length. Other habitat assessment work included an assessment of substrate, flow rate, and stream morphology along five transects distributed evenly within the 100 m assessment reach. Slimy sculpin (*Cottus cognatus*) were relatively abundant (23 were caught), and a variety of life stage and sizes/weights were noted, indicating that the proposed effluent mixing zone represents potential spawning and rearing habitat for slimy sculpin. A single bull trout (*Salvelinus confluentus*) was captured in the channel margin of the creek in shallow, slower moving water. Given the lack of instream cover or rearing habitat available at the point of capture, it is likely that this individual trout was utilizing these slower flows as an opportunity to rest during a migration past the effluent site.

CONTAMINANTS 2011

Blowes, David University of Waterloo Waterloo, ON blowes@uwaterloo.ca

File No: 12 402 843 **Licence No:** 14846

Region: NS Location: Diavik Diamond Mines Inc. - Lac de Gras mine

site

Waste rock studies at a diamond mine site

The objective of this research was to investigate the processes related to water quality and quantity draining from experimental waste rock piles that are located in areas of continuous permafrost. Waste rock piles are mounds of rock removed from open pit and underground mines. The quality of water draining from a waste rock pile is determined by the combined effects of oxygen transport in the air, biogeochemical processes that control mineral weathering rates, the release of heat and dissolved constituents due to sulfide mineral oxidation, and hydrologic processes that control unsaturated water flow. The transport of dissolved constituents is further affected by the formation and subsequent dissolution of secondary minerals. Three instrumented experimental waste rock piles were constructed from 2004 to early 2007 at the Diavik diamond mine. In 2010, three 40 m deep boreholes were drilled into the operational waste dump and a series of instruments similar to those in the test piles were installed. In 2011, two additional 40 m boreholes and one 80 m borehole were installed and instrumented with similar instruments. Data from these instruments will be compared to data from the test piles to evaluate differences in measurement scale. The study is undertaken entirely as a graduate research program through the University of Waterloo, University of British Columbia, and University of Alberta. At least one presentation on the Diavik test piles project will be presented at the Geoscience Forum in November, 2011 in Yellowknife.

Evans, Marlene

Environment Canada Saskatoon, SK marlene.evans@ec.gc.ca

File No: 12 402 681 **Licence No:** 14950

Region: SS Location: Great Slave Lake: West Basin- near Fort

Resolution; and West Basin- near Hay River

Spatial and long-term trends in persistent organic contaminants and metals in lake trout and burbot from the Northwest Territories

This is an ongoing study under the Northern Contaminants Program (NCP); some support was provided by the Cumulative Impacts Monitoring Program. The researchers are investigating whether contaminant levels are changing in Great Slave Lake fish, which they have been studying since the early 1990s. Twenty lake trout from Hay River, 20 pike and 20 burbot from Fort Resolution, and 20 lake trout and 20 burbot from Łutsel K'e were shipped, frozen and whole, by community members to a lab in Saskatoon. In the lab, length, weight, and age were determined and samples were submitted for metals (including mercury) and persistent organic contaminant analyses. Pike and Łutsel K'e burbot were analyzed only for mercury. Mercury has been showing an increasing trend, although levels remain below the 0.5 ppm guideline. Organic contaminant concentrations have not been increasing; some such as HCH and DDT have decreased in concentration due to their decreased usage. The results of their 2011 findings will be presented in a 2011 NCP report, with those findings shared with their community partners.

Gantner, Nikolaus (Klaus)

University of Victoria Victoria, BC gantnern@uvic.ca

File No: 12 402 868 Licence No: 14974

Region: IN Location: Inuvik; Tuktoyaktuk; Yaya Lake; Noell Lake; Big

Lake; Husky Lakes system; 16 lakes along the 177 road; 4

lakes along the Tuk-Inuvik road corridor

Evaluation of hydro-climatic drivers of contaminant transfer in aquatic food webs in the Husky Lakes Watershed (Inuvialuit Settlement Region, NWT)

This fall field campaign was a multi-purpose trip that included some helicopter-supported work, land-based work, and more outreach/consultation. The timing for the late fall/winter work was suggested to the researchers by the Tuktoyaktuk Hunters' and Trappers' Committee (Tuk HTC) and during the review by the Regional Contaminants Committee in the proposal stage. The researchers were asked to come conduct sampling of fish and other parameters when fishers set fishing nets under the ice, thereby limiting helicopter use and their scientific fishing efforts in this ecologically and culturally sensitive ecosystem. Prior to arranging flights to Inuvik, the researchers communicated with the Aurora Research Institute (ARI) in Inuvik and the Tuk HTC to ensure the ice on the lakes was safe for travel. They spent 3 weeks in the Inuvialuit Settlement Region, split between Inuvik and Tuktoyaktuk. They met with local partners (ARI, Fisheries Joint Management Committee, Fisheries and Oceans Canada, Tuk HTC) and the public in Tuktoyaktuk to discuss project details and future directions. They went to all study lakes to develop the methods related to sample collection for two graduate projects (at U Victoria and U Manitoba). The field crew at all sites consisted of three people, with 2-4 local people helping in Tuktoyaktuk. They used Spot devices to map their field trips via GPS Satellite tracking.

Geddes, Robert

AMEC Earth & Environmental Calgary, AB brian.geddes@amec.com

File No: 12 402 862 Licence No: 12940

Region: SA Location: Along the south shore of the Great Bear River,

from the source of the river on Great Bear Lake to Tulít'a

Great Bear River site assessments

No research was conducted under this licence in 2011.

Krizan, Julia IMG-Golder Corporation Inuvik, NT jkrizan@golder.com

File No: 12 402 664 **Licence No:** 14953

Region: IN **Location:** The abandoned Panarctic Satellite F-68 well

site, located at Satellite Bay on the north-western coast of

Prince Patrick Island

Detailed site description, remediation feasibility and risk assessment of the Panarctic Satellite F-68 Wellsite, Satellite Bay, Prince Patrick Island, NWT

Talisman Energy Inc. conducted a supplemental Phase II Environmental Site Assessment (ESA) program at the abandoned Panarctic Satellite F-68 wellsite area at Satellite Bay on Prince Patrick Island, Northwest Territories . Panarctic Satellite F-68 was a dry exploratory petroleum well drilled in the 1970s. In 2010, several Areas of Potential Environmental Concern were identified around the wellsite. These included the well site area itself, a small pond containing old steel fuel drums, a landfill area, surficial debris, surface stains, potential buried (likely detonated) explosives, and an area of suspected buried debris adjacent to the landfill. During the 2011 program, soil and water sampling was conducted to delineate contaminated areas that were identified during the 2010 program. The debris that had been consolidated in 2010 remained as it was left in 2010. The supplemental Phase II ESA included a geophysical survey covering areas not included in 2010, testing of surface water from the creeks/rivers near the Site, and collection of groundwater and soil samples. Samples were submitted to an accredited laboratory for analysis and compared to applicable territorial and federal guideline criteria. Exceedances for metals and petroleum hydrocarbons were identified in both 2010 and 2011. Delineation of known contamination was conducted in 2011. Information was collected to define the extents of contaminated soil and groundwater as well as waste materials. Some geotechnical data was collected to support the potential design and construction of a landfill on-site. An assessment of the airstrip was also conducted to determine what aircraft could safely land/take off, as well as what maintenance would be required to support larger aircraft.

McLachlan, Stephane

University of Manitoba Winnipeg, MB mclachla@cc.umanitoba.ca

File No: 12 402 863 **Licence No:** 14954

Region: SS **Location:** Nagel Channel Boat Landing, Fort Resolution;

Fort Smith Boat Launch

Multi-scale environmental health implications of the Athabasca oil sands for Aboriginal communities in Alberta and Northwest Territories

From June 4-20 2012, three members of the University of Manitoba's Environmental Conservation Laboratory conducted field research contributing to this project. During this period the researchers conducted video interviews with community members in Fort Chipewyan, Alberta and Fort Smith and Fort Resolution, Northwest Territories, about environmental change and community concerns about contamination arising from industrial activities affecting the Slave River and Peace Athabasca river deltas. The researchers shared community newsletters created to publicize the

work they had completed between June 2011 and spring 2012 with community collaborators. These newsletters are an attempt to create a communication network for communities to share their knowledge and concerns with each other, and for the scientific community to communicate their research approaches and results. Two thousand copies of these newsletters have been printed for distribution to First Nation and Métis Nation research partners in Fort Chipewyan, Fort Smith and Fort Resolution. The researchers anticipate return visits to Fort Chipewyan in fall 2012 and production of a second newsletter for distribution to all community partners in spring of 2013.

Wiatzka, Gerd

SENES Consultants Ltd. Richmond Hill, ON gwiatzka@senes.ca

File No: 12 402 778 **Licence No:** 14872

Region: NS, SS **Location:** Blanchet Island Mine; Outpost Island Mine;

Copper Pass Mine; DeStaffany Mine

Great Slave Lake area mines: site assessment and remediation planning

No research was conducted under this licence in 2011.

Wolfe, Brent

Wilfrid Laurier University Waterloo, ON bwolfe@wlu.ca

File No: 12 402 866 **Licence No:** 14966

Region: SS Location: Slave River Delta

Sediment core sampling to assess contaminant deposition to the Slave River Delta (NWT) over time

The perceived negative downstream effects of oil sands development is a major environmental issue of international concern. In the Slave River Delta, a key unknown is whether industrial activity is enhancing the delivery of natural sources of oil sands-derived contaminants. In this study, sediment cores were taken from a flood-dominated lake in the Slave River Delta. The historical record is necessary for determining baseline concentrations of polycyclic aromatic compounds (PACs). Using a gravity corer, lake sediment cores were collected in September 2011 from a small (~1.2 km²), shallow (maximum depth ~1.5 m) flood-dominated lake in the active Slave River Delta. Sediments in this lake contain a record of spring break-up flooding for at least the past century. Sediment cores were sectioned into 1 -cm intervals, placed in sample bags and shipped to the University of Waterloo. Samples are currently being analyzed for 1) radiometric isotope (137Cs, 210Pb) concentration to develop the sediment core chronology, 2) loss-onignition to characterize physical properties of the sediment core, 3) organic carbon and nitrogen elemental and isotopic composition to reconstruct past hydrological conditions, and 4) PACs to examine their depositional patterns and trends over time. Research is being conducted as an initiative of the Slave River and Delta Partnership. Fieldwork in September 2011 was conducted with the assistance of a community member. Also, while conducting fieldwork, presentations were made to classes of school children at the Fort Resolution Deninu School and an open-house was held in Fort Resolution to inform the community of the project. Analyses are in progress but results are expected to contribute new knowledge of how and if PAC deposition in the Slave River Delta has changed over time. Research will contribute to addressing concerns of local residents over the perceived impacts of upstream oil sands development, and will help prioritize future research and monitoring needs.

ENGINEERING 2011

Patterson, R. Tim
Carleton University
Ottawa, ON
tpatters@earthsci.carleton.ca

File No: 12 406 054 **Licence No:** 14949

Region: NS **Location:** Along the length of the Tibbitt to Contwoyto

Winter Road

Paleoclimatological assessment of the central Northwest Territories: implications for the long-term viability of the Tibbitt to Contwoyto winter ice road

In support of the researchers' three-year multi-disciplinary research project, 80 sediment/water interface samples from 43 lakes along the route of the Tibbitt to Contwoyto Winter Road have thus far been analyzed for water property data, substrate characteristics, nutrient loading, water geochemistry, isotopes and environmentally available metals. This data is being used to develop training sets and transfer functions based on micropaleontological proxies; the camoebians, diatoms and chironomids. Twenty-one Glew cores and 16 freeze cores have also been collected from these lakes. Use of a freeze core microtome has permitted subsampling of freeze cores to mm-resolution (2-5 years). Preliminary time series analyses results indicate that throughout the late Holocene there has been considerable climate variability, with winter and summer signals often becoming decoupled. The Pacific Decadal Oscillation and North Atlantic Oscillation have contributed to step-wise temperature changes as these phenomena vary between positive and negative phases. There is also a correspondence between solar cycles and seasonal climate variability, with solar cycle peaks corresponding to cooler summers and warmer winters, and troughs corresponding to warmer summers and colder winters.

Trimble. Annika

Aurora Research Institute Inuvik, NT atrimble@auroracollege.nt.ca

File No: 12 406 058 **Licence No:** 14910

Region: NS **Location:** Thor Lake near the Avalon Rare Metals Inc.

development

Wind energy monitoring at Thor Lake 2010-2011

In 2009/10, one year of wind data was analyzed from a 50 m meteorological station installed near Thor Lake on a ridge overlooking the Hearne Channel. The projected long-term average wind speed at 48 m above the ground is estimated to be 5.7 m/s. At 80 m above the ground, the long-term wind speed is estimated to be 6.5 m/s. Data collection continued through 2010/11 with similar

results, though winter wind speed measurements were not recorded due to mechanical issues. This will be resolved by extending the monitoring stage of the project. All reports related to this project are available at www.nwtresearch.com

Trimble. Annika

Aurora Research Institute Inuvik, NT atrimble@auroracollege.nt.ca

File No: 12 406 058 Licence No: 14852 Region: NS Location: Wekweètì

Wind energy monitoring in Wekweètì: 2010-2012

Two wind monitoring towers were installed in Wekweètì in October, 2010, following a prefeasibility study and community consultation. The first tower installed is 34 meters tall, and the second is 10 meters tall. Each tower is equipped with anemometers to measure wind speed, a wind vane for wind direction, a meteorological station to record weather, and a data logger to store the information. The 34 m tower is now monitoring wind speed and direction for potential wind energy to serve the entire community, and the 10 m tower collects data for potential wind energy to directly serve the new complex. After the towers were installed, a local resident was hired and trained to be the wind monitor. His duties include monthly site visits, the collection of data, and maintenance of the towers. These towers will collect data for up to two years, after which the data will be analyzed and a feasibility report written. Recommendations will be presented to the community for review and discussion. In 2011, data collection was ongoing. The project technician returned to Wekweètì to check the towers and provide additional training. For more information on wind energy project activities, please visit the researchers' website at www.nwtresearch.com

Trimble. Annika

Aurora Research Institute Inuvik, NT atrimble@auroracollege.nt.ca

File No: 12 404 720 **Licence No:** 14963

Region: DC Location: Jean Marie River; Fort Providence

Solar irradiance monitoring in Jean Marie River and Fort Providence

The majority of Canada's northern communities are dependent on fossil fuels for electricity generation. Due to their remoteness, the cost of transporting diesel fuel to these communities is a large financial burden on the territory's government and utility companies. Renewable energy, particularly in the north, offers many potential benefits to northern communities. Using wind or solar power in place of diesel can help to reduce particulate emissions and the greenhouse gas emissions which contribute to climate change. Solar energy is of particular interest in the southern parts of the Northwest Territories, though little irradiance data has been collected in the territory to date. The objective of this project, therefore, is to measure solar irradiance levels in Jean Marie River and Ft. Providence, in order to support pre-feasibility studies on the use of solar energy in those communities. In August, 2011, the project engineer traveled to Jean Marie River and Fort Providence to install solar irradiance monitoring equipment in the two communities. The sensors will collect solar irradiance data for at least one year, depending on the success of data capture and annual variation in weather (cloud cover, etc.). Once the study is complete, the sensors will be removed and pre-feasibility studies will be produced and distributed in both plain language and technical reports. These reports will be made available at www.nwtresearch.com.

HEALTH 2011

Badry, Dorothy
CanNorthwest FASD Research
Calgary, AB
badry@ucalgary.ca

File No: 12 410 901 **Licence No:** 14969

Region: IN, NS, SS Location: Łutsel K'e; Behchokò; Ulukhaktok; Yellowknife

Brightening Our Home Fires: Women and wellness project program report

Brightening Our Home Fires was a qualitative research project, which used Photovoice to explore women's perceptions, attitudes and experiences related to the prevention of Fetal Alcohol Spectrum Disorder (FASD) in the Northwest Territories (NT). Four communities participated: Behchokò, Ulukhaktok, Yellowknife and Łutsel K'e. Yellowknife was also included as women from remote northern communities often migrate to this urban location for reasons such as homelessness, domestic violence, substance abuse treatment, financial need, employment opportunities and a stronger resource network. Women participants were given digital cameras, some hands on training by research team members and an introduction to the research question: What does health and healing look like for you in your community? About 30 women participated from the NT. The women provided beautiful images and captions from their communities about their understanding of health and healing – a foundation to FASD prevention. The photos and words that were shared with the research team provided a deep understanding of the need for culturally based health support resources. Key findings relate to the importance of children, relationship to elders and participation in traditional activities as important parts of health and healing. Photovoice helps women voice life experiences. FASD prevention requires involvement of women, men and communities.

Brennan, Jodi

Aurora College Yellowknife, NT jbrennan@auroracollege.nt.ca

Formative evaluation study of the BSN program

The formative evaluation study of the Bachelor of Science in Nursing (BSN) Program has collected some data from student and faculty focus groups and student questionnaires over the past six months. Unfortunately, this collaborative study across sites has been stopped by the Collaboration for Academic Education in Nursing (CAEN) Steering Committee. Information can

still be retrieved specifically for Aurora College, until this point in the study. Therefore, there is no need to continue this research licence.

Goodman, Karen

University of Alberta Edmonton, AB karen.goodman@ualberta.ca

File No: 12 408 149 **Licence No:** 14886

Region: IN, GW Location: Aklavik; Tuktoyaktuk

The Aklavik *H. pylori* project

In February 2011, the İnuvialuit Settlement Region (ISR) *H. pylori* pilot project was launched in Tuktoyaktuk, NT. Of the 93 participants who enrolled in the pilot, 86 had a breath test for *H. pylori* infection, 35 provided health data, and 23 provided individual-level and household-level socio-environmental data. Using feedback from this pilot, planning is underway for the full ISR *H. pylori* project to launch in 2012. One component of the researchers' community *H. pylori* projects is the collection of interview data on health problems related to *H. pylori* infection, along with relevant data from participants' medical records. In May 2011, the researchers completed a project to assess the completeness of information obtained from medical records in Aklavik, and to improve their chart review tool to collect more accurate information. In September 2011, a project examining ways to foster effective communication between researchers, knowledge users, and other community members involved in the Aklavik *H. pylori* project was initiated. This project aims to help promote the direct application of research knowledge to effective *H. pylori* screening practices, both in Aklavik and other remote northern communities throughout Canada.

Hammond, Merryl

Consultancy for Alternative Education Baie d'Urfé, QB merryl.hammond@videotron.ca

File No: 12 408 148 **Licence No:** 14885

Region: IN Location: Aklavik; Ulukhaktok

Changing the "culture of smoking": Community-based participatory research to empower Inuvialuit communities

This community-based participatory research (CBPR) project began in Aklavik and Ulukhaktok in 2007. CBPR team members completed baseline surveys in late 2008, and completed data entry in 2009. Data were checked and analyzed. A summary of findings was shared with communities in late December 2010. The response rate was 50% in Ulukhaktok and only 12% in Aklavik; in Ulukhaktok, 60% said they smoke every day and 25% are non-smokers, while in Aklavik these figures are 38% and 45% respectively. In both communities, peer pressure was the biggest reason to start smoking. Smokers in Aklavik expressed a much greater willingness to quit. In Aklavik, "health concerns" were mentioned by 56% of ex-smokers, but were mentioned by only 29% of ex-smokers in Ulukhaktok. A total of 52% of women in Ulukhaktok reported smoking during their most recent pregnancy, compared to only 30% in Aklavik. Aklavik smokers smoked more than Ulukhaktok smokers. The teams launched a second smoking cessation challenge, the Be Smoke-free Challenge, in November 2010. Community responses were excellent, with 33% of the total population of Aklavik signing up, and 24% in Ulukhaktok. More non-smokers than smokers entered, and slightly more women and girls than men and boys.

Hannon, Judy

Canadian Blood Services Edmonton, AB judy.hannon@blood.ca

File No: 12 408 142 Licence No: 14982

Region: IN, GW, SA, DC, NS, SS Location: Community health centres; Hospitals; Clinic

laboratories

RHD alleles in prenatal patients from northern Canada

No research was conducted under this license in 2011.

Kuhn, Karen

University of Bath Victoria, BC karen.kuhn@telus.net

File No: 12 408 184 Licence No: 14981

Region: DC, NS **Location:** Yellowknife Primary Care Clinic and Stanton

Territorial Hospital; Fort Simpson Clinic; Behchokò Clinic;

Dettah Clinic

Evaluation of the electronic health record (EHR) system used in the Northwest Territories

No research was conducted under this licence in 2011.

Martin, Jim

Tłįcho Community Services Agency Behchokò, NT imartin@tlicho.net

File No: 12 408 143 **Licence No:** 14932

Region: NS Location: Behchokò; Gamètì; Wekweètì; Whatì;

Yellowknife

Tłicho Natsedzi Nihtsi: Tłicho healing wind project / promoting sexual health

Despite many well-intentioned programs and policies, the burden of illness among rural Aboriginal communities continues to grow, particularly in the North. In 2006, the rate of sexually transmitted infections (STIs) in the Northwest Territories was eight times the Canadian rate, at 20.4 per 1,000. In the same time frame, the rates in the Tłycho Region were three times the Northwest Territories' rate, at 67.5 per 1000. This summary describes how four rural and remote Aboriginal communities reduced their rates of syphilis and improved sexual health in the relevant age groups. The Tłicho Community Services Agency (TCSA) Board and the Tłycho Government strongly supported the development of community research skills to provide the base for programs and policies that fit with regional priorities, including sexual health. Traditional clinically based models were ineffective in reducing the STI rates in the region, so the TCSA designed a community-based strategy, which began with visits by community leaders to every household to provide information for families. Next, a region-wide survey conducted by trained community-based researchers showed how people learned about sexual health, and what their attitudes were towards related issues. The follow-up activities were hard to implement; people who were expected to do the knowledge translation were already busy in their full-time TCSA positions, and thus communication and program development were uneven. To fill this gap, a team of young Tłycho adults was recruited in May of 2009 to be trained as researchers and communicators. They worked with health

professionals and others to develop, conduct and evaluate research to ensure that programming and policies continued to respond to community needs and priorities. The Community Action Research Team (CART) was trained by local people in cultural values, and by academics in research and communication skills. CART developed a series of programs and resources based on the survey findings. Examples include resource materials (e.g. pamphlets, booklets, manuals, posters, etc.), media (e.g. radio, DVD, web, blog, etc.), and community events (e.g. workshops in the community, youth conferences, puberty camps for boys and for girls, focus group discussions). Research and communications training and mentoring were provided by CIETcanada, an international non-government research organization. A second survey was conducted in 2010 to measure progress. That study revealed that condom use was making a difference, as was participation in CART-related activities. This was an integrated program; at the health centre, the public health nurse trained one of the community health representatives to trace and visit contacts in the communities and encourage people to come to the clinic for testing. The education system welcomed classroom presentations. The CART members created a video production for YouTube, featuring local actors and elders. The integration of community, public health, education and social programs and academic perspectives created a collaborative approach that led to a series of activities to sustain the low rates of infection, and to support behavior change regarding sexual health. This is an on-going program, and activities will continue. with collection of biomedical data, frequent focus groups with affected age groups, targeted activities and regular evaluations to monitor the effectiveness of programs.

Mitton, Craig

University of British Columbia Vancouver, BC craig.mitton@ubc.ca

File No: 12 408 183 **Licence No:** 14899

Region: IN, GW, SA, DC, NS, SS Location: Health Authorities of the NWT

Achieving high performance in health care priority setting

Between January and April 2011, the researchers conducted an on-line survey of senior executive team members at the Regional Health Authorities across Canada. The researchers received 92 responses, with 4% of total respondents from the Northwest Territories, representing four Health and Social Services Authorities. The survey inquired about structures, process features and behaviors related to organization-wide resource allocation decisions in the health sector. Decision making rules and procedures, enabling and constraining factors, criteria, participation, and internal and external communications were among the topics addressed. Among key findings were the following: about one-half of respondents indicated that their organization used primarily a formal process for resource allocation, while the others reported that political or historical factors held sway. Seventy percent of respondents agreed or strongly agreed with the statement, "our resource allocation process is fair". Just over one-half assessed their own process as 'good' or 'very good'. Complete results are available upon request to the investigators.

PHYSICAL SCIENCES 2011

Armstrong, Terry

Envronment and Natural Resources, GNWT Fort Smith, NT terry armstrong@gov.nt.ca

File No: 12 404 750 **Licence No:** 14964

Region: DC, SS **Location:** From the Mackenzie River to approximately

Birch Lake

Landscape scale flooding in the Great Slave Lake Plain

A comparison of tree growth records from sampling in 2010 and temperature and precipitation records from Fort Providence and Hay River suggested growth of trees in this region was influenced by a combination of temperature and precipitation. The relationship is convoluted in recent decades as many of the sampled trees became submerged by rising water, which affected their growth. In 2011, 14 new sites and multiple species (white spruce, tamarack, jack pine) were sampled in areas thought to be unaffected by rising lake levels, which may provide a more dependable climatic relationship and strengthen confidence in results. In addition to the tree core samples, two lake-core samples from Caen Lake were obtained, which will aid in the understanding of past climates of the region. Tree- and lake-core samples are currently being analyzed. Aerial photographs and satellite imagery of 12 lakes in the study area showed that some had dramatic changes in size between the late 1940's and 2010, while others had changed little. Most of the study lakes had flooded large areas in the late 1940's, up to 1971, were smaller by 1984, and then increased. Some grew to 8 to 10 times their earlier size.

Arrigo, Kevin

Stanford University Stanford, CA United States arrigo@stanford.edu

File No: 12 404 768 Licence No: 14916 Location: Arctic Ocean

ICESCAPE - Impacts of Climate on EcoSystems and Chemistry of the Arctic Pacific Environment

No research was conducted under this licence in 2011.

Aubet, Natalie
University of Alberta
Edmonton, AB

aubet@ualberta.ca

File No: 12 404 756 **Licence No:** 14980

Region: GW, NS **Location:** Point Lake; Russell Lake; Damoti Lake; Bell

Lake; Rapitan; Yellowknife

Precambrian banded iron-formations: palaeoceanographic, palaeoclimatic, and palaeobiologic implications

During 2011, the researchers reviewed the sampling conducted in 2010 in collaboration with staff from the Northwest Territories Geoscience Office. Activities were partially funded through a grant from the Canadian Circumpolar Institute (CCI Research Grants Program), and the Natural Sciences and Engineering Research Council of Canada (NSERC). Traditionally, investigations about banded iron formations (BIF's) and associated sediments have focused on sequences from South Africa, Australia, India, and Brazil. Geochemical studies on BIF's from Canada, especially from the Northwest Territories, however, are still scarce. Geochemical characterization of BIF's from the Northwest Territories provides relevant information to the researchers' understanding of the atmospheric evolution of Earth, as well as the chemical composition of the oceans during the Archean-Paleoproterozoic (2500 my ago), which ultimately lead to the appearance of animal life on Earth (~750 my ago). As a result of this work, two communications will be presented at (1) the 22nd Goldschmidt Geochemistry Conference (Montreal, June 2012), and (2) the 39th Annual Yellowknife Geoscience Forum (November, 2012). In addition, one paper about the geochemistry and geochronology of BIF's from the Northwest Territories is under preparation.

Bédard, Jean

Geological Survey of Canada Québec, QB jbedard@nrcan.gc.ca

File No: 12 404 735 Licence No: 14924

Region: IN Location: Ulukhaktok base camp

Northern base and precious metal potential, Victoria Island (NWT) and Nunavut

Major discoveries and achievements in the field (so far) include: (1) completion of the planned mapping block in the Proterozoic rocks; (2) completion of the follow up to the stream sediment survey; (3) completion of the mapping and subdivision of the Paleozoic succession; (4) documentation of the chemical heterogeneity of the volcanic succession, which indicates the existence of a complex feeder system with multiple vents; (5) discovery of unconsolidated scoria, spatter and fumarolic deposits, and of pillow-hyaloclastite deposits, allowing the researchers to localize vent complexes; (6) the association of native copper deposits with these fumarolic vent deposits. This provides, for the 1st time, a context for the occurrence of the native copper in the region; (7) lateral facies and thickness variations of lahar-like deposits implying infilling of prevolcanic topography; (8) discovery of volcaniclastics and exhalites at the Killian-Kujjua transition, marking the awakening of the volcanic system at that time: (9) recognition of pepperitic deposits on the fluviatile sands of the Kujjua; (10) recognition of coarse volcanic deposits (possible agglutinate); (11) Discovery of two feeder complexes, that include sulphide-rich propagator tips. This suggests that wallrock assimilation did indeed trigger immiscibility in places: (12) discovery of a sub-sill gossan zone, including rheomorphic breccias, hybrid magma bodies, graphitized black shales, and extensive sulphide stringers in the footwall; (13) discovery of two more oxidesulphide skarn systems associated with sills; (14) discovery of ankaramitic lavas, which implies that primitive magmas were capable of reaching the surface, and so increases the size of the exploration target. Community Engagement: Two graduating high-school students were involved as geological assistants for 2 weeks. Three Inuvialuit archaeological sites were discovered and

reported to the ILA. The two OHTC wildlife monitors were extremely helpful, and the researchers provided wildlife sighting logs to them.

Bhatti, Jagtar

Natural Resources Canada Edmonton, AB jbhatti@nrcan.gc.ca

File No: 12 404 679 **Licence No:** 14870

Region: DC, GW, SA **Location:** Inuvik; Norman Wells; Fort Simpson; Nahanni

Butte; Wrigley

Recent changes in carbon source-sink relationships and greenhouse gas emissions in forest and peatland ecosystems along the Mackenzie Valley region of Canada

Summary of research not provided for this 2011 licence.

Bird. Sam

WorleyParsons Canada Calgary, AB sam.bird@worleyparsons.com

File No: 12 404 777 Licence No: 14957

Region: IN **Location:** Esso Tuk Base 800 m east across Tuktoyaktuk

Harbour from Tuktoyaktuk

Tuk Base demolition supplementary sampling

During the 1970's and 1980's Imperial Oil built a logistic base on the east side of Tuktoyaktuk Harbour called Esso Tuk Base. The land is Inuvialuit Private 1 (a) Land and is managed by the Inuvialuit Land Administration. Since about 2006, Imperial Oil has been cleaning up the site by removing old drilling supplies and equipment that is not required. Buildings and fuel tanks were demolished at the site during the summer of 2011. The debris from the demolition was sent to the south for disposal. During past clean-up activities at the site, soil and water samples were collected to look at areas that may be contaminated by substances such as hydrocarbons, metals and salt. The past results found soil contaminated by hydrocarbons. In 2011, soil samples were collected to better understand the size of contaminated soil areas. Groundwater samples were collected from monitoring wells to continue the existing ground water monitoring program. Soil and water samples were sent for analysis at a laboratory in Edmonton. The soil results have helped better determine where the boundary between contaminated soil and clean soil is located. The water results showed that contamination is not moving through the groundwater.

Bottenheim, Jan

Environment Canada Toronto, ON jan.bottenheim@ec.gc.ca

File No: 12 404 729 Licence No: 14834 Region: IN Location: Arctic Ocean

O-buoy measurements of ozone, carbon dioxide and bromine oxide over frozen surface of Hudson Bay and Arctic Ocean

O-buoy instrumentation package is capable of long term measurements of the concentrations of bromine monoxide and two important greenhouse gases (ozone and carbon dioxide), along with

a full suite of meteorological parameters, ice drift, sky and ice conditions. One of the two units that belongs to Air Quality Research Division of Environment Canada was deployed with the help of the Canadian Extended Continental Shelf Program at latitude 88.15° N and longitude 157.49° W on September 5, 2011. Data collected are transmitted on daily schedule via satellite and displayed on the public web site. Seven weeks of data have been collected, since the deployment. Data are in the process of being integrated into the Marine Weather Forecasting and International Artic Buoy Programs. O-buoy measured carbon dioxide and ozone concentrations are within the concentration range recorded by an identical unit over the ice of Beaufort Gyre. The project results were reported at the NWT IPY conference held in Inuvik in January 2011. Data is currently being analyzed and the most interesting findings will be presented at IPY conference in Montreal in April 2012.

Burgess, David

Natural Resources Canada Ottawa, ON david.burgess@nrcan.gc.ca

File No: 12 404 707 Licence No: 14859

Region: IN **Location:** South Melville Ice Cap

Melville Island South Ice Cap mass balance and snow pollution

Measurements of snow accumulation and ice melt were performed at 21 pole locations on the Melville South Ice Cap by D. Burgess and J. Zheng on April 18 and 19, 2011. The Melville Island South Ice Cap is a small plateau ice cap (76 km2 in size) that is located on the western portion of Melville Island in the Canadian high Arctic, Northwest Territories. Pole measurements indicate that the ice cap, as a whole, has thinned by 94 cm, which equates to a loss of 0.072 km3 (or 1 km x 1 km x 0.072 Km) of water to the ocean over the past year, as a result of warm temperatures during the summer of 2010. These melt rates are second only to 2007, which was the year of greatest mass loss experienced by the ice cap, since the record began in 1963. Temperature data downloaded from the automatic weather station on the ice cap indicate that the 2010 summer melt season extended from early June to late August, with periods of sub-zero temperatures in early July and early August. Recent thinning has resulted in large areas of bedrock being exposed within the interior sections of the Melville South Ice Cap. These areas of bedrock have likely been covered by ice for several thousand years. Further investigation of these sites may offer insight into the age of the ice cap, and possibly the climatic conditions that prevailed at that time. Continued monitoring of the Melville South Ice Cap is important, as the rapid changes experienced by this ice cap are valuable for gaining insight into long-term climate change over the western Canadian Arctic region. Mass wastage of ice caps across the Canadian high Arctic is currently a significant contributor to global sea-level rise.

Burn, Chris

Carleton University Ottawa, ON christopher_burn@carleton.ca

File No: 12 404 325 Licence No: 14942

Region: IN Location: Garry Island, Mackenzie delta area; Illisarvik,

western Richards Island; Inuvik, near the Dempster highway; Paulatuk, near the community; Red Lake; Bar C;

Seal Lake; Dennis Lagoon

Permafrost and climate change, western arctic Canada

In 2011, the researchers visited three sites for their research activities. They made two separate visits of 10 days at the Illisarvik drained lake site on Richards Island. There, they began to measure the small earthquakes that happen when ice wedges crack. The researchers installed little data loggers that are meant to record ground shocks. They have not been used in the western Arctic before, so their installation was an experiment. The researchers went to Paulatuk, in July, with Dr. Ross Mackay. Their research there was concentrated on ground temperatures and abrasion of rocks, as before. They continued their studies of permafrost conditions near Inuvik. In 2012, they will write up data on tilting of trees above permafrost. They have published two papers this year, one on changes in vegetation at the pingo near Paulatuk, and one on the place names of Herschel Island. The paper on vegetation at Pinguksayuk is the first paper to report changes in the distribution of willows in the western Arctic. It compared photos taken in 1910 with present conditions. A large amount of their work in 2011 was focussed on preparing a book about Herschel Island for publication. This will be available in April 2012. It will be 252 pages long and have over 400 photos, both historical and taken in the last 10 years.

Clark, lan

University of Ottawa Ottawa, ON idclark@uottawa.ca

File No: 12 404 534 **Licence No:** 14919

Region: GW **Location:** Near Fort McPherson

Chronology of thaw flow and geochemistry of associated massive ground ice. Fort McPherson, Northwest Territory, Canada.

In May and June 2011 fieldwork was undertaken in the Stony Creek watershed (NT), to study the impacts of retrogressive thaw slumps on the terrestrial and aquatic ecosystems. The work was performed by students and professors from the University of Ottawa, with valuable field assistance provided by Gwich'in monitors from Fort McPherson. Below the researchers provide the tasks performed during their field season and anticipated outcomes.: (1) Data loggers that record water quality measurements, such as water level, conductivity and turbidity, were installed in streams above and below two thaw slumps, to monitor the quality of clear tundra streams and streams impacted by slump runoff. At these two sites, stream velocity measurements were taken to calculate ionic and sediment fluxes used to determine the contribution of slump runoff to the overall stream budget; (2) A survey of stream water quality in the Stony Creek watershed was performed, during both field campaigns. Here, stream water samples were collected at the mouth of Stony Creek, at several locations above, within and below the thaw slumps and from clear tundra streams. The samples were analyzed for major dissolved ions and suspended sediments. The data will allow the researchers to assess the impacts of slumps on the water quality downstream; (3) Permafrost drilling was undertaken in a thaw slump, to obtain samples of ancient peat and soil, in August, 2011. The peat will be analyzed for iodine, iodine-129 and carbon-14 using facilities at the University of Ottawa. The permafrost was also analyzed for major ions and stable isotopes to characterize the sediment in the thaw slumps and permafrost melting. The outcome of these analyses will allow the researchers to investigate changes in the fallout of iodine-129 in the recent past, as well as hundreds of years in the past.

Corriveau. Louise

Geological Survey of Canada Québec City, QB lcorrive@nrcan.qc.ca

File No: 12 404 716 Licence No: 14844

Region: SA, NS **Location:** Grouard Lake; Lou Lake; Gamètì; Cole Lake

GEM Great Bear magmatic zone/iron oxide copper-gold deposit project

The Great Bear magmatic zone is a geological belt that extends from Great Bear Lake to Great Slave Lake. The belt hosts two iron oxide-copper-gold (IOCG) deposits and has a very high mineral potential for other undiscovered IOCG and affiliated deposits. It remains, however, significantly under explored. The current project, within the Geomapping for Energy and Minerals program, and its partners have now laid a solid geoscientific foundation for exploring IOCG in the region and for land use planning. In summer 2011, NRCan conducted fieldwork in partnership with the Community Government of Gamètì, academia, and Fortune Minerals. The project models, techniques and mapping protocols for IOCG systems were tested southwest of Lou Lake and east of Gamètì. While based in Gamètì, Tłycho community members were employed and helped test the mapping protocols on an area centered on the historic Fab Lake showings. Field observations confirm the effectiveness of the protocols and highlight the ability of IOCG systems to create affiliated mineralization, such as albite-hosted uranium. Interim results were presented nationally and internationally, including at the Yellowknife Geoscience Forum, with formal publications to follow. Results are attracting attention of exploration companies and scientists worldwide and pave the way for a new cycle of mineral exploration in the region and new collaborations.

Coulton, Daniel

Golder Associates Ltd. Yellowknife, NT daniel coulton@golder.com

File No: 12 404 763 **Licence No:** 14898

Region: NS Location: Fortune Mineral's NICO property, along the

route of a proposed all-weather access road from Highway

3

Environmental baseline surveys of the Fortune Minerals Ltd. NICO project

No research was conducted under this licence in 2011.

Dallimore, Scott

Geological Survey of Canada Sidney, BC sdallimo@nrcan.gc.ca

File No: 12 404 359 **Licence No:** 14836

Region: IN, GW Location: Mackenzie Delta; Richards Island

Mackenzie Delta shallow gas and permafrost studies

This multi-year project attempts to quantify geohazards and environmental considerations related to release of methane gas from aquatic and terrestrial areas in the outer Mackenzie Delta, with special emphasis on the controls of permafrost and gas hydrates. The researchers conducted two field programs in 2011. In late-March, they deployed two aquatic moorings, to record the acoustics (sound characteristics) of gas discharge at two of their gas seep study sites. The researchers were then able to convert the acoustic data in gas flux measurements, to investigate temporal variability in gas discharge. They also deployed two dissolved oxygen sensors, to investigate changes in dissolved oxygen concentration over time. In October, the researchers deployed water column samplers for analysis of water properties at three sites. Their research results have been submitted to a journal for publication and have been presented at several national and

international meetings. This work will ultimately assist in understanding if the warming permafrost and gas hydrate deposits found in outer the Mackenzie Delta pose a hazard to hydrocarbon development or surface activities and also assist to calibrate fluxes of greenhouse gases.

Duffe, Jason Environment Canada Ottawa, ON

jason.duffe@ec.gc.ca

File No: 12 404 743 Licence No: 14923

Region: IN **Location:** The coastline of the Beaufort Sea, from the

Alaska/Yukon border to the Northwest Territories/Nunavut

border

Assessing the potential for environmental sensitivity index mapping in the Arctic using synthetic aperture radar

In late July-early August, the researchers collected geo-tagged oblique video and audio commentary over almost 3000km of Northwest Territories' coastline, from Alaska to Nunavut. RADARSAT-2 and TerraSAR-X satellite imagery were acquired at different periods of time and at different incidence angles over the study sites. All study sites are also covered with optical data (SPOT and LANDSAT). Radar and optical data analyses will be conducted independently, to determine the most suitable techniques and optimal datasets to differentiate the shoreline types (intertidal and supratidal zones). Textural analysis and polarimetric data will be generated and used with ancillary datasets, such as bathymetric data, surface deposits maps and wind data to classify shoreline type. Comparisons between the traditional approach (oblique video) and remote sensing techniques will be conducted to verify if the satellite products are as reliable as the traditional approach. A series of ground plots was also completed in Ivaavik National Park and in the Anderson River Migratory Bird Sanctuary. An intensive ground program was also completed on Herschel Island, where the researchers also acquired 2 hyperspectral images from the European satellite CRIS. The shoreline video will be interpreted into a Beaufort Coastal Sensitivity Atlas and research results on satellite analysis will be completed through the winter and reported next year.

Duthie, Andrew

Rescan Environmental Services Ltd. Yellowknife, NT aduthie@rescan.com

File No: 12 404 752 Licence No: 14884

Region: NS Location: Matthews Lake; Courageous Lake; Dumbell

Lake; Jolly Lake; Sandy Lake

Courageous Lake project

The 2011 environmental baseline program collected data, to characterize the physical and biological setting of the proposed Courageous Lake project area. The following activities were carried out:

<u>Meteorology:</u> The meteorological station and wind tower were maintained. Ten snow course surveys were completed at representative locations across the project area.

<u>Air Quality:</u> Dust fall monitoring was undertaken at five stations, every 30 days over the summer. Noise: Six stations were monitored for 24 hours in spring.

<u>Hydrology</u>: Eight hydrological stations were established. Water current velocity and related flow discharge measurements were determined throughout the open-water season. Current meters were installed in Courageous Lake, during the open water season, to obtain data.

<u>Bathymetry</u>: Surveys were completed at Courageous Lake, Matthews Lake, and two unnamed lakes of interest.

<u>Hydrogeology</u>: Two groundwater wells were drilled and Westbays installed. Packer testing was undertaken, to determine hydraulic parameters of aquifers. Water samples were taken from aquifers below the permafrost. Data loggers were installed on existing thermistor strings. Measurements were collected from thermistor strings.

<u>Aquatics</u>: Water quality was sampled at 14 lakes and 9 streams three times over the summer. A subset of these lakes was sampled in March. Sediment quality and primary and secondary producer communities were sampled in mid-summer.

<u>Fish and fish habitat surveys</u>: 65 lakes and 43 streams were sampled. 15 lakes and 13 streams were identified as fish bearing. 2 lakes and 25 streams were found to be dry or not flowing during freshet. Sensitive Habitat Inventory and Mapping (SHIM) were undertaken for Matthews Creek. Ecosystems: Terrain mapping and soil samples were collected, in conjunction with vegetation surveys, in the project area. Field surveys were conducted to identify wetlands in the proposed project area. Baseline reports will on the above aspects will be available in March 2012, when data analysis and report writing has been completed.

Fortier, Martin University Laval

Quebec, QB

martin.fortier@arcticnet.ulaval.ca

File No: 12 404 652 **Licence No:** 14917

Region: IN Location: Beaufort Sea/Mackenzie Shelf/Amundsen Gulf

region

ArcticNet: an integrated regional impact study of the coastal western Canadian Arctic.

Since 2004, ArcticNet has been using the Canadian research icebreaker CCGS Amundsen to carry out sampling operations in the Beaufort Sea/Mackenzie Shelf/Amundsen Gulf region, as part of its ongoing marine-based research program. The central aim of this research program is to study, on a long-term basis, how climate induced changes are impacting marine ecology. contaminant transport, biogeochemical fluxes, and exchange processes across the ocean-sea ice-atmosphere interface in the Canadian Arctic Ocean. In 2011, sampling operations in the Beaufort Sea/Mackenzie Shelf/Amundsen Gulf region were carried out from the CCGS Amundsen, from 11 August to 04 October. During these 55 days, researchers sampled at over 50 oceanographic stations. Sampling operations included deployments of a CTD-Rosette, box corer, Agassiz trawl, and plankton nets. A total of 4 sub-surface oceanographic moorings were deployed. In addition, a multitude of oceanic and atmospheric parameters were measured continuously, using the Amundsen's impressive array of continuous samplers (SM-ADCP, EK-60 scientific echosounder, water surface pCO2 and CTD on track system, foredeck and top bridge meteorological towers, ceilometers, radiometer and all-sky camera). The ship's EM302 multibeam sonar and Knudsen sub-bottom profiler collected over 15,000 km of high-resolution bathymetry and sub-bottom data. From the vessel's wheelhouse, hired Inuvialuit Marine Wildlife Observers sighted and identified marine mammals and seabirds. Data collected from this multi-year program will contribute to a better understanding of the impacts of climate variability and change on the

physical, biological and geochemical processes in the Beaufort Sea/Mackenzie Shelf/Amundsen Gulf region.

Grogan, Paul Queen's University Kingston, ON groganp@queensu.ca

File No: 12 404 687 **Licence No:** 14943

Region: NS Location: Daring Lake Terrestrial Ecosystem Research

Station

Controls on carbon and nutrient cycling in arctic tundra

This research looks at: What are the principal controls on the functioning of common tundra ecosystem types, and how are they likely to be affected directly and indirectly by climate change? In 2011, most of the summer was spent at Daring Lake collecting vegetation and soil samples from a greenhouse warming experiment. Samples are currently being processed, in order to determine the potential for warming to alter nitrogen and phosphorus availability, as well as vegetation change in response to rising air and soil temperatures. Other progress in 2011 includes: a manuscript is currently in review with *Global Change Biology*, demonstrating that birch shrub apical growth, at the Daring Lake research site, is limited as much by the availability of phosphorus as it is by nitrogen. This is a very surprising result. For many researchers across the Arctic, the focus has been on nitrogen. The biogeochemical cycling of phosphorus is fundamentally different. Based on this result, the researchers' understanding of how low arctic terrestrial ecosystems function, as well as how plant growth and vegetation change will be affected by climate warming, will be determined by the changes in both the availability of phosphorus and nitrogen.

Haas, Claudia

Environment and Natural Resources Yellowknife, NT claudia_haas@gov.nt.ca

File No: 12 404 774 **Licence No:** 14946

Region: DC Location: Ekali Lake (Ezáa Łue Túe); Sanguez Lake

(Tł'onie Túé); Gargan Lake (Tłįtętįį); Deep Lake (Dechį

Ná?a); McGill Lake (Tthets'éhk'e')

Ecological assessment fieldwork for Lue Túé Sulái (the Five Fish Lakes) candidate cultural conservation area

The survey was conducted in August 2011 and was supported by the Jean Marie River community and a larger DFN-AAROM collaborative management initiative. Three of the five lakes within the candidate area were sampled: Ekali, Sanguez, and Gargan Lakes. An inflow, centre and outflow site were established for each lake. Some measurements were taken at the sites and water samples from the surface were shipped to and analysed by Taiga Labs, Yellowknife. The water quality within all three lakes is generally good. Water temperature was uniformly warm, pH was slightly basic, major ion concentrations were moderate, and nutrient, chlorophyll and metal concentrations were low. However, dissolved oxygen was often reduced, mostly at depths below 5 meters in Ekali and Sanguez, but 50% saturation values were recorded for all depths within Lake Gargan. Shoreline observations were made for each station. Vegetation was similar and ranged from Black Spruce and Tamarack through Birch, Aspen & Willow to Alders, Dogwoods and Rosehips. Emergent vegetation generally occurred in the shallows and near shore sediment

ranged from organic, woody debris to sandy and small rocks. Elders provided traditional knowledge, to identify spawning areas and good fishing sites for each lake and the creeks between them. General hydrological observations were also documented.

Halverson, Galen McGill University Montreal, QB galen.halverson@mcgill.ca

File No: 12 404 769 Licence No: 14921

Region: SA **Location:** The Northern Mackenzie Mountains, specifically

in the Sheep Lake/Mountain River area

Late Proterozoic stratigraphy of northwestern Canada and its record of Earth system evolution

In July 2011, a research group consisting of geologists from Queen's University and Harvard University visited outcrops of the latest Precambrian sedimentary rocks in the Sekwi Brook region of the Northwest Territories. The purpose of this research was to better understand the age and depositional setting of sedimentary rocks deposited during intervals of extreme glaciation and in which some of the earliest animal fossils in the world are found. Fieldwork consisted of mapping of these rocks and collecting rock samples for geochemical analysis. Early results include: (1) the most precise age estimate yet produced for one of these glacial events; (2)new data bearing on the water depth and oxygen content of the sedimentary basin in which these early animals resided. Importantly, the geological mapping has shown that some of the fossil occurrences, which based on prior mapping appeared to significantly pre-date other similar fossils globally, are in fact younger than thought and hence do not constitute an outlier that forces rethinking of earliest animal evolution. Nevertheless, the new work provides a valuable new geological framework for interpreting the environmental setting in which these early animals lived. This project ties in with an ongoing research program on similarly-aged rocks in Yukon.

Herber, Andreas

Alfred Wegener Institute Bremerhaven Germany andreas.herber@awi.de

File No: 12 404 710 **Licence No:** 14895

Region: IN Location: Sachs Harbour; A N/NW route approximately

200 Km from Sachs Harbour and then returning to Sachs

Harbour along the same path

Polar airborne measurements and arctic regional climate model simulation project

The campaign polar airborne measurements and arctic regional climate model simulation project (PAMARCMIP) 2011 was performed from March 25, 2011 until May 06, 2011. The aircraft was operated by the Canadian Aviation partner Kenn Borek Air Ltd. Scientists from different research institutes, including Germany, Canada, and USA were involved in the project. The researchers performed airborne measurements (127 flight hours) in the inner Arctic over six weeks and successfully completed a traverse from the North American Arctic (Barrow, Alaska) to the European Arctic (Longyearbyen, Svalbard). Northward flights were performed over the Arctic Ocean from Barrow, Inuvik, Eureka, Alert, Station Nord as well as Longyearbyen. The researchers arrived in Inuvik on April 3, 2011 and left Inuvik on April 6, 2011. During this period they performed two research flights, measuring ice thickness, trace gases, aerosols and meteorological

parameters over a key region of the Arctic. Unfortunately the planned landing in Sachs Harbour, with the aim to extend the endurance for the research flights, was cancelled, due to bad weather conditions. Measurements were also made during the ferry flights from Barrow to Inuvik on April 3, 2011 and from Inuvik to Resolute Bay on April 6, in the altitude range of 10 000 feet.

Hicks, Faye

University of Alberta Edmonton, AB faye.hicks@ualberta.ca

File No: 12 404 493 **Licence No:** 14892

Region: SS **Location:** Along the Hay River

Hay River ice jam study

The 2010 field research program brought members of the University of Alberta (U of A) team, and colleagues from the Department of Indian Affairs and Northern Development (DIAND) to the Town of Hay River, to observe, measure, and document river breakup ice conditions (April 19 to May 10). During breakup, U of A/DIAND field crews worked with the Town Flood Watch Committee, to measure ice jams and to document the river's breakup progression. Breakup was relatively uneventful that year, with low water/ice levels and no flood threat. Nevertheless, there were numerous ice jam formation and release events along the river, providing excellent scientific data to aid in the advancement of the researchers' flood forecasting models. Operational testing of these U of A ice jam flood forecasting models continued during breakup 2011. The timing of the onset of breakup, the expected peak snowmelt runoff stream flow and the time of arrival of the ice runs from were all predicted with reasonable accuracy. Further research is needed to identify when these incoming ice runs will actually stall upstream of the community, and breakup 2011 provided valuable data towards understanding this scenario. The U of A and DIAND researchers will be providing the Town of Hay River with an update on the flood forecasting model development project prior to breakup in 2012.

Hilton, Robert

Durham University Durham, County Durham United Kingdom r.g.hilton@durham.ac.uk

File No: 12 404 717 **Licence No:** 14925

Region: GW, DC, SS **Location:** The Mackenzie River and its main tributaries

Geological carbon in the Mackenzie River Basin: sources and sinks of atmospheric carbon dioxide

In May and June 2011, the research team spent their third consecutive year sampling river sediments in the Mackenzie River Basin. Essential logistical support was provided by the Aurora Research Institute, Environment Canada, and the local communities. As in June 2009, river water and suspended sediment samples were collected from 'depth profiles' within river channels, using the researchers' custom-built, clean, depth sampler. At the same time, researchers used an 'Acoustic Doppler Current Profiler', to measure the speed of the water in detail. Analysis is ongoing. In 2011, the researchers hoped to collect samples from the river as close to the freshet (ice break up) as possible, because this is when the rivers carry most of their sediment. The campaign was a success. The May 2011 samples are the most sediment-rich collected in the three years of work on the Mackenzie River. The increase in sediment concentration with water depth is also useful to the researchers. It means each sample reflects different sizes and masses

of particles, which can be investigated. Researchers are currently analyzing the samples for their chemistry, to better understand the source of carbon contained in the water and sediments.

Holmes, Robert

Woods Hole Research Center Falmouth, MA United States rmholmes@whrc.org

File No: 12 404 713 **Licence No:** 14881

Region: GW **Location:** The Mackenzie River, near Inuvik; from shore,

and near the ferry crossing, near Tsiigehtchic

Arctic great rivers observatory

This project studies the 6 largest rivers that flow into the Arctic Ocean: in North America, the Mackenzie and Yukon; and in Russia, the Ob', Yenisey, Lena, and Kolyma. The researchers are measuring the concentration of naturally occurring chemicals, such as carbon, nitrogen, and phosphorus, in these rivers to obtain baseline information about the flow of these chemicals to the ocean, and to help them understand how climate change is impacting Arctic rivers. This is a 3 year project, and the researchers are now nearing the end of the third year. Most of their samples have been collected, but laboratory analyses are still underway. All data from this project is posted on a public website (http://arcticgreatrivers.org) and is available for free download by the public. The researchers took 2 sampling trips to the Mackenzie River in 2011. In late May and June, they took daily 1 liter samples of river water from the shore near Inuvik, as well as 3 samples by boat near the Tsiigehtchic ferry crossing. In September, they took one sample by boat near Tsiigehtchic. All boat samples were less than 15 liters of water. During their 2011 sampling year, they were fortunate to have Will Storr (of Fort McPherson / Tsiigehtchic) provide local field support.

Kanigan, Julian

Indian and Northern Affairs Canada Yellowknife, NT julian.kanigan@inac.gc.ca

File No: 12 404 661 Licence No: 14937

Region: DC **Location:** Historic seismic lines near the Mackenzie

Highway, between 61-62°N and 120°30'-121°30'W; several sampling sites will be located in the Scotty Creek Research

Basin

Investigating the effects of winter overland travel in sub-arctic boreal forest

The objective of this project is to investigate terrain conditions associated with transportation infrastructure, specifically seismic lines, in discontinuous permafrost. In July 2011, researchers met with staff from the Liidlii Kue and Jean Marie River First Nations, to discuss the project and opportunities for local involvement. Measurements were obtained at several road-accessible seismic lines. In areas underlain by permafrost, active layers were significantly thicker along seismic lines than in undisturbed areas. Subsidence in the order of centimeters was observed along the seismic lines, leading to wetter conditions and different vegetation. In August 2011, sites were established in 3 common terrain units adjacent to the Liard Highway: peat land, fen, and till. The sites were instrumented with shallow (1 m) ground temperature cables, soil moisture and air temperature sensors. Results from these sites will establish the natural variability of active layer freeze back between different terrain types. Deep (10 m) ground temperature cables were

installed in the Scotty Creek basin at a channel fen, and at the center and edge of a degrading peat land. Shallow and deep ground temperature cables were also instrumented along a nearby winter road and seismic line. Results may indicate how line widths, line orientation, and groundice content relate to permafrost resilience along disturbance corridors.

Kokelj, Steve

Indian and Northern Affairs Canada Yellowknife, NT steve.kokelj@inac.gc.ca

File No: 12 404 545 **Licence No:** 14840

Region: IN, GW Location: Mackenzie Delta

Environmental studies across treeline

This study is a northern-based collaborative program initiated by the Renewable Resources Directorate, AANDC, in 2004. It is designed to enhance the understanding of environmental conditions in the region, through scientific studies and monitoring. In 2011, the researchers continued to monitor permafrost and active layer temperatures in the Delta region, by visiting previously established sites by helicopter and boat. This was the third year of active-layer freeze back monitoring, and results indicate that different terrain types freeze back in the same order each year, but there is significant variation in the timing of freeze back between years. This information will be useful for decision makers for permitting winter overland travel. In 2010, the researchers removed tall shrubs from an abandoned drilling mud sump in the Mackenzie Delta, in an attempt to promote freezing conditions. Lower ground temperatures were measured in 2011 in the sump cap at depths up to 3 m. Continued monitoring may indicate that vegetation removal is a valuable long term sump maintenance technique. Ground temperatures were measured near proposed Mackenzie Gas Pipeline stream crossings. The thermal regime of stream valleys is significantly different than adjacent uplands, and an understanding of these differences is important for pipeline construction in the region.

Kokelj, Steve

Indian and Northern Affairs Canada Yellowknife, NT kokeljsv@inac.gc.ca

File No: 12 404 545 Licence No: 14842

Region: GW **Location:** Stoney Creek catchment, which runs parallel to the Dempster Highway and empties into the Peel River at

Fort McPherson

Evaluating the environmental impacts of permafrost mega-disturbances along the Dempster Highway, NWT

This project studies the cumulative impacts of permafrost slumping on the land and water and addresses several landscape change, water and fisheries questions that have been determined through the regional Renewable Resource Council (RRC) gatherings, Gwich'in Renewable Resources Board meetings and the Gwich'in Water Strategy Workshop. This project studies the impacts of big slumps on streams and fish in the Peel Plateau. Mapping shows that there are hundreds of big slumps like those seen off of the Dempster Highway. The slumps are impacting the Rat, Willow, Vittrekwa, Trail, Road and Caribou Rivers, as well as Stony Creek. By looking at old air photographs the researchers determined that the slumps are much bigger than in the 1970s. Studies tell them that the big slumps cause major changes to landscape and the streams and these changes impact what can live in the streams. The streams become choked with mud.

Slumps are having impacts on water in the Peel River. The support of the Tetl'it RRC and the community of Fort McPherson have been a key to the success of this project. In 2011, the Cumulative Impact Monitoring Program (CIMP) funded about 100 days of employment to community members from Fort McPherson. Community members helped researchers decide where to sample water, to travel safely and respectfully on the land and they have made many observations of changes to the environment. Community monitors have collected information on plants, berries, permafrost, water and the health of the streams. The Tetl'it RRC played an important coordinating role and administered funds to community researchers. The study is providing information to support fish and wildlife management, and planning and maintenance of community and transportation infrastructure. The data users include the Department of Fisheries and Oceans, the Community of Fort McPherson, the Department of Transportation and the Gwich'in Land Use Planning Board.

Kors-Olthof, Rita

Nehtruh-EBA Consulting Ltd.(on behalf of the Hamlet of Aklavik) Yellowknife, NT rkors-olthof@eba.ca

File No: 13 404 770 **Licence No:** 14928

Region: IN,GW **Location:** Along about 5 km of traditional trail between about 2 km west of Aklavik and 7 km west of Aklavik, west

of the junction with the Peel Channel

Proposed Aklavik west road and bridge - Hydrotechnical and geotechnical field investigations

In 2011, Nehtruh-EBA Consulting Ltd. did a research program at Bridge Creek, about 3.5 km west on the traditional trail from Aklavik, NT to the Richardson Mountains. The people in Aklavik have made some different bridges over the creek, but the bridges usually wash away in spring. So, in June 2011, Nehtruh-EBA sent a hydrotechnical engineer and a surveyor to look at the bridge site, and take some measurements of the stream bed and flood levels. They also took measurements at the Water Survey of Canada gauge on the Peel Channel, so that they would have some flood history with which to compare. Two local environmental monitors helped with the work. Now the hydrotechnical engineer can design a bridge high enough not to wash away when there is a flood. The information from the surveyor helps him to do his calculations. In August 2011, Nehtruh-EBA asked an air photo company to fly over the first part of the traditional trail and take some photos looking straight down. The photos overlap, so if you look at them through special glasses, it is like a 3D picture. This 3D picture helps the engineers see problem areas along the trail, and helps them think of ways to keep the trail good for travelling. It also helps them find sand or gravel to improve the trail if needed.

Lafleur, Peter

Trent University Peterborough, ON plafleur@trentu.ca

File No: 12 404 621 Licence No: 14851 Region: NS Location: Daring Lake

Exchange of carbon gas fluxes over low arctic tundra

The researchers continued their research into carbon fluxes from arctic tundra, near Daring Lake, NT, in 2011. Instruments were set up in early May and continued operating until late August. The researchers made measurements at 4 different tundra types: fen, mixed heath, low shrub heath

and tall shrubs. At each site, they measured carbon dioxide fluxes between the tundra and atmosphere. The overall objective is to see if the tundra is taking more carbon dioxide out of the atmosphere by plant photosynthesis than it is releasing by respiration. If more goes in than goes out, the tundra is a sink for carbon and if more goes out, it is a source. As carbon dioxide levels in the atmosphere build up from burning gas and oil, the researchers hope the tundra is a sink, which indeed their 2011 results seem to show for all the sites. They measured important differences in how much carbon dioxide is taken out among the sites. For example, the tall shrubs are a larger sink than the heath and low shrubs, but about the same as the fen. This research helps them to understand how arctic tundra will influence the amount of carbon dioxide in the atmosphere and thus how it might influence the climate today and into the future.

Landry, Francois

Rescan Environmental Services Ltd. Vancouver, BC flandry@rescan.com

File No: 12 404 767 **Licence No**: 14913

Region: SS **Location:** Near the former Pine Point Mine

Pine Point project (N-204)

Seven dust fall monitoring stations were installed and over 90 samples collected, along with noise monitoring. Water bodies at all deposit areas, as well as Great Slave Lake, were surveyed to characterize water quality and measure baseline metals in sediment. Fish populations were surveyed at over 50 sites in the deposit areas and habitat assessed. A 170 meter deep borehole was drilled and tested at the R-190 deposit and hydraulic testing was done at the upper Pine Point Formation. Six monitoring wells were drilled and installed at the N-204 deposit, and hydraulic testing and sample collection was completed. Four site visits were completed for hydrologic flow characterization, and hydrometric monitoring stations were installed along the Buffalo River and Twin Creek, to record surface water elevations. Fifty-onesamples from deposit areas were tested for metal leaching potential. Soils and vegetation field data were collected at 80 inspection sites within the six deposit areas, to classify terrain and ecosystems, and to characterize soils for soil salvage. Forty-five plant samples and 22 soil samples were collected from a subset of the inspection sites, to determine levels of baseline metals. The study area was surveyed for wetlands. which were then mapped. An archaeological assessment was conducted and four prehistoric sites were recorded. Potential land users and key socio-economic informants will be identified and interviewed in the fall and winter, to determine existing levels of land use near the project area and to characterize the social and economic environment. Traditional knowledge studies are planned to be conducted in 2012, in collaboration with aboriginal groups near the Pine Point project area (Akaitcho Territory Government, Dehcho First Nations, Deninu Kue First Nation, Hay River Dene Band/Katlodeeche First Nation, Northwest Territory Métis Nation, and West Point First Nation).

Langhorne, Amy

Golder Associates Ltd.
Saskatoon, SK
amy_langhorne@golder.com

File No: 12 404 733 Licence No: 14894

Region: NS, SS **Location:** The Kennady Lake watershed

De Beers - Gahcho Kué environmental monitoring program

Baseline environmental studies were completed during the spring, summer and fall in the area surrounding the Gahcho Kué project in 2011. The surveys were conducted with the assistance of representatives of the Yellowknife Dene First Nation and Łutsel K'e Dene First Nation. The studies encompassed weather, air quality, fisheries, water quality, and hydrology. The hydrological regime was monitored through measurement of water levels and river flow between each water body. Meteorological data was recorded starting in August from the weather station at site. Noise monitoring was not required in 2011. Fish surveys were completed in small lakes and streams throughout the local study area and at Kennady Lake. The surveys included gill netting, electrofishing and minnow trapping. Fish habitat was also mapped in the surrounding streams and small lakes. Water samples were collected to update water quality information for Kennady Lake, and surrounding and downstream lakes. Stream outlets and lakes were surveyed, along with the watershed adjacent to Kennady Lake. Benthic invertebrate sampling was also completed in the lakes.

Lawson, Nick

Det'on Cho Stantec Yellowknife, NT nick.lawson@stantec.com

File No: 12 404 773 Licence No: 14944

Region: NS, SS Location: Within Avalon's Thor Lake Property, 100 km

southeast of Yellowknife

2011 baseline studies for Avalon Rare Metals Inc. proposed Thor Lake rare earth element project - surface water hydrology and climate

The objective of the surface water hydrology field program was to characterize the surface water hydrology and meteorological conditions at the Thor Lake site. Surface water hydrology fieldwork, during 2011, included water level monitoring in the following lakes at the project site: Thor, Long and Cressy lakes. Water levels were recorded using a Hobo pressure transducer and water level gauge secured to the lake or stream bed. Stream flow monitoring was completed at the outlets of Thor, Long, Fred and Murky lakes. Stream flow measurements were taken following standardized methods for stream flow in May and October. Meteorological conditions were monitored at the Thor Lake site, using an AXYS Watchman 500 weather station. Data were periodically downloaded and compiled; station maintenance was also completed. Lake water levels, stream flows, and meteorological data was compiled, analyzed, and compared to regional data. Deton'Cho Stantec produced an updated technical data report in December 2011.

Lennie-Misgeld, Peter NWT Hydro Corporation

Yellowknife, NT

plennie-misgeld@ntpc.com

File No: 12 404 708 **Licence No:** 14835

Region: NS, SS **Location:** Barnston River; Beaulieu River; Hoarfrost River;

Waldron River

NT Hydro hydrology monitoring program

In May 2010, NT Hydro's contractor, Water Survey of Canada, installed water gauging stations on the Hoarfrost, Barnston, Beaulieu and Waldron Rivers. Gauging stations are remotely operated and collect hydrology data on a full time continuous basis. The goal of the program is to collect hydrology information, to better understand the hydrology and hydro potential of these rivers in this area of the Northwest Territories. The four stations continue to collect data to develop a

complete hydrology record. Hydrology information collected includes: water level, water and air temperature, water volume and velocity. Data will continue to be collected for the next 2-3 years, to develop a hydrology record for the rivers. Once enough data has been compiled, NT, Hydro will be able to evaluate the hydro potential of these rivers.

Lintern, Gwyn

Geological Survey of Canada Sidney, BC glintern@nrcan.gc.ca

File No: 12 404 612 **Licence No:** 14935

Region: IN Location: Mackenzie Delta

Coastal geoscience research in the Beaufort Sea and Mackenzie Delta

No research was conducted under this licence in 2011.

MacNaughton, Robert

Geological Survey of Canada Calgary, AB robert.macnaughton@nrcan-rncan.gc.ca

File No: 12 404 529 **Licence No:** 14858

Region: SA Location: Mackenzie Mountains near Norman Wells and

Tulít'a

Geological fieldwork in Mackenzie Plain and adjacent mountains.

A team of eleven scientists from the Geological Survey of Canada (Calgary), University of Calgary, University of Ottawa, University of Saskatchewan, Texas A&M University, and James Madison University did geological fieldwork based out of Norman Wells, for five weeks in July and August, 2011. Working with them was a Wildlife Monitor from Norman Wells. Two scientists also worked out of Tulít'a for one week in July, accompanied by a local Monitor. Helicopter, accommodation, and food services were provided by local businesses. Fieldwork involved helicopter visits or overland hiking to 458 rock outcrops on ridges and streams from the eastern Mackenzie Mountains to the Franklin Mountains. Locations and rock descriptions were recorded, and rock thicknesses and orientations were measured. Approximately 300 rock samples were collected, varying from fist size to slightly larger than a loaf of bread. Samples were shipped to labs at the Geological Survey of Canada in Calgary, or at the above mentioned universities, where they are undergoing paleontological, geochronological and organic chemistry analyses. Data are being used to produce new geological reports and maps of bedrock geology for the Norman Wells and Tulít'a region (NTS map areas 96C, 96D, 96E, and 96F).

Marsh, Philip

Environment Canada Saskatoon, SK philip.marsh@ec.gc.ca

File No: 12 404 378 Licence No: 14837

Region: IN, GW Location: Trail Valley Creek; Havikpak Creek; Denis

Lagoon; Big Lake, 65 km W of Tuktoyaktuk, 130 km N of

Inuvik; 75 km E of Tuktoyaktuk

Hydrological studies, Mackenzie Delta region

With a changing climate and increasing development, there is an urgent need for appropriate hydrological information (snow cover, soil moisture, soil temperature, and stream discharge) in the western Canadian Arctic. For example, the design of roads and pipelines requires estimates of maximum stream discharge, while rules controlling land access in the fall require estimates of snow cover and whether the soil is frozen. However, with a changing climate, the recent past may not be a reliable guide to the hydrological conditions in the near future. As a result, in order to limit the environmental impact of development, the research team needs better methods to predict future conditions. The research program is aimed at developing such improved methods, and over the last year has: (1) collected hydrologic data at two study sites, in order to extend the 20+ year data set; (2) extended the data collection by installing new state of the art equipment. This year, the research team installed a continuous GPS unit that will collect important snow and soil information; (3) continued to develop better methods to predict future changes in snow cover, soil moisture, ground thaw, and stream flow. Recent results consider the factors controlling the thaw of the upper layer of the ground over the summer period. This is an important step towards better predictions of the impact of a changing climate and developments on the hydrology of the region. Other results have considered the role of lakes on the hydrology and ecology of the Mackenzie Delta.

McCallum, Dee

De Beers Canada Inc. Yellowknife, NT dee.mccallum@ca.debeersgroup.com

Region: NS, SS **Location:** Snap Lake and the regional study area (33 km

radius from camp)

De Beers Snap Lake Mine - 2011-2014 environmental monitoring program

Monitoring of water quality, sediment quality, zooplankton, phytoplankton, benthic communities, fish, fish habitat, and fish health was successfully carried out in the 2011 field season. The hydrology program monitored lake levels, stream-flow, and outflows; measured site runoff; and collected hydro-meteorological data. Additional water quality monitoring was conducted during construction and installation of the new diffuser. The 2011 aquatics program results have yet to be analyzed and reported. This information will be available in the 2011 Aquatic Effects Monitoring Program (AEMP) Annual Report for Snap Lake which will be submitted to the Mackenzie Valley Land and Water Board (MVLWB) by March 31st, 2012 and available on the MVLWB Public Registry online at (http://www.mvlwb.ca/mv/registry.aspx). Many parameters were measured as part of the 2011 geochemistry monitoring program. An annual site inspection to monitor site runoff/seepage, review the placement of materials, and identify any signs of acid generation was also carried out in September. The summary field report for this inspection is available on the Public Registry. The air quality monitoring program involved the collection and processing of meteorological data from the onsite weather station, as well as the measurement of particulate matter and dust-fall from sampling stations on site. No vegetation monitoring was carried out in 2011. As per the Vegetation Monitoring Plan, the next vegetation monitoring at Snap Lake will take place in 2013. The results of all these monitoring programs will be submitted in the 2011 Annual Report, which will be available on the MVLWB's Public Registry.

Miles, Warner

Geological Survey of Canada, NRCan Ottawa, ON wmiles@nrcan.gc.ca

File No: 12 404 727 **Licence No**: 14876

Region: NS **Location:** The project area is on the Hearne Channel of

the East Arm of Great Slave Lake, 90 km southeast of

Yellowknife

Blatchford Lake, NT airborne gravity gradiometry survey

The objective of this research was to acquire high-resolution gravity gradiometer and aeromagnetic data, in the Blatchford Lake, NT area, centered on the Thor Lake rare earth element (REE) deposit. The gravitational field measured by this survey reflects lateral variations in the density of underlying rocks. The aeromagnetic survey measured magnetic properties of bedrock. Both data types are tools used in geological mapping. Understanding these gravity and magnetic data will help geologists map the area, assist mineral exploration activities, and provide information necessary for communities, aboriginal associations, and government to make land use decisions. The survey collected approximately 3,066 line km of data flown along parallel lines spaced 250 m apart. The flying height was at a nominal terrain clearance of 100 m. The horizontal gradient of gravity and the intensity of the total magnetic field were measured from the aircraft. The survey was flown between March 19, 2011 and March 23, 2011. Final data have been accepted for the survey. The data and maps were published on October 14, 2011. The data are available for free download from the Geoscience Data Repository for Aeromagnetic and Electromagnetic Data (http://gdr.nrcan.gc.ca/aeromag) and digital versions of the maps are similarly available from MIRAGE (http://gdr.nrcan.gc.ca/mirage). The survey results were presented at a poster session of the Yellowknife Geoscience Forum on November 15-17, 2011. The poster was entitled Geological significance of a new high resolution gravity gradiometric and magnetic survey over the Blatchford Lake Complex. The data acquired over the Blatchford Lake area are of high quality and will serve their intended purpose.

Miles, Warner

Geological Survey of Canada, NRCan Ottawa, ON wmiles@nrcan.gc.ca

File No: 12 404 718 Licence No: 14983

Region: SS **Location:** The survey will be flown in an area 230 km from

the Łutsel K'eDene First Nation, 305 km from the Deninu K'ue First Nation, and 215 km from the Smith's Landing

and Salt River First Nation locations

South Rae, NWT aeromagnetic survey

No research was conducted under this licence in 2011.

Milton, Jack

University of British Columbia Vancouver, BC jmilton@eos.ubc.ca

File No: 12 404 734 **Licence No:** 14845

Region: SA, DC Location: Mackenzie Mountains - the Backbone Ranges

and the Redstone Plateau, Coates Lake; Fortress

Mountain; Ten Stone Range

Geology of the Redstone Copperbelt

Four weeks were spent in the field around Coates Lake in the Central Mackenzie Mountains. Detailed geological mapping was carried out, to support ongoing research and thesis work at the University of British Columbia (UBC), Vancouver, BC. This was the final field season for the project and the fieldwork has provided a wealth of information, regarding copper mineralization in the Mackenzie Mountains. Further laboratory studies will continue at UBC and the results will be written up as a doctoral thesis. Initial results of the field studies have furthered scientific understanding of how copper is transported and concentrated in certain areas of the Earth. These results can be used to increase the efficiency of mineral exploration projects in the Northwest Territories and also in other parts of the world where similar geological environments can be found, for example: central Africa, Poland, eastern Russia and Afghanistan.

Mloszewski, Aleksandra

University of Alberta Edmonton, AB mloszews@ualberta.ca

File No: 12 404 759 **Licence No:** 14979

Region: GW, NS **Location:** Point Lake; Russell Lake; Damoti Lake; Bell

Lake; Rapitan; Yellowknife

Investigating the influence of Archean seawater composition on the evolution and diversity of microbial metallo-enzyme evolution, through the chemistry of Archean banded iron formation

The objective of this research project is to examine the chemistry of the oceans directly preceding the oxidation of the atmosphere at ca. 2.4 billion years ago (The Great Oxidation Event), by studying recently-discovered Neoarchean (2.8 to 2.6 GA) banded iron formations (BIF) in the Northwest Territories. In August (2010), the researchers explored these units for the first time and took transects of the sampling areas and a number of samples for petrographic and chemical analyses. In 2011, analyses determined the stratigraphic, initial petrographic and geochemical framework needed in order to conduct detailed geochemical analytical work that will be completed in 2012. Sampling sites included the ca. 2.62 billion year old BIF at Point Lake, which are interbedded with greywacke-mudstone turbidites, the ca. 2.85 billion year old BIF in the Central Slave Cover Group ~30 km north of Yellowknife, and detailed sampling of the BIF belonging to the Central Slave Cover Group at the Northwest Territories Geoscience Office (NTGO) core lab. Thin sections for petrographic and chemical analyses have been made of these samples and chemical analyses are underway.

Moore. Kristin

Diavik Diamond Mine Inc. Yellowknife, NT kristin.moore@riotinto.com

File No: 12 404 766 Licence No: 14907 Region: NS Location: Lac de Gras

Diavik aquatic effects monitoring program 2011

Network program stations indicated an "early warning" or "low level" effect on water chemistry within Lac de Gras resulting from the mine. Analysis of benthic invertebrates indicated a range of effect designations. Effects on Procladius sp. density and percent Chironomidae were classified as "early warning" or "low level"; the effect on total benthic invertebrate density was classified as a "moderate", and the effect on Heterotrissocladius sp. density was classified as a "high" level effect. Overall, benthic invertebrate monitoring results indicate mild nutrient enrichment. Results

of a study to examine changes in the amount, number and types of tiny animals (zooplankton) and algae (phytoplankton) that live in the water of Lac de Gras indicate a pattern consistent with nutrient enrichment from the mine, similar to eutrophication indicators, measured as part of the aquatic effects monitoring program. Phytoplankton and total phosphorus measurements in the near-field area resulted in a "moderate" level effect designation while higher zooplankton biomass near the effluent resulted in a "high" level effect designation. Results of the lake trout study suggest that there has been an increase in mercury in the muscle tissue of lake trout, in both Lac de Gras and Lac du Sauvage, since 2005. The increase from baseline resulted in a "low" level effect designation. However, since the increase was observed in both lakes, it cannot be directly linked to the mine. The weight-of-evidence analysis confirmed the nutrient enrichment effect and concluded that there is strong evidence for a mild increase in lake productivity as a result of nutrient increases in Lac de Gras. There is some evidence suggesting potential low-level toxicological impairment of the benthic invertebrate community, although these findings have high uncertainty, because the link to contaminant exposure is not strong and the responses indicating possible impairment are not consistent with the multiple other responses indicating enrichment.

Mumford, Thomas

Carleton University Ottawa, ON thomas.mumford@gmail.com

File No: 12 404 736 **Licence No:** 14948

Region: SS Location: Blachford Lake Intrusive Suite of rocks,

centered on Thor Lake, about 100 km southeast of

Yellowknife

Petrogenesis of the Blachford Lake intrusive suite

Work completed during 2010-2011 involved processing samples collected during the 2010 field season. The samples were divided into two portions, one part was crushed and powdered for analysis to determine composition, and the other was used to determine mineralogy. A geochronological study was also initiated to determine the age of the Thor Lake syenite, a phase in the Blachford Lake intrusive suite, which has some historic controversy. The previous date of this unit was almost 100 million years (Ma) younger than the rest of the units of the Blachford Lake Intrusive Suite, far beyond what was expected and analytical error. Based on their study, the researchers have shown the previous date to be incorrect. Fieldwork, during the summer of 2011, consisted of sampling carbonate bearing dykes along the shore of the Hearne Channel, Great Slave Lake. These dykes were sampled to determine if they had a relationship to the sedimentary rocks on the south-side of the channel, or to the Blachford Lake Complex; geochemical results which are currently pending will be used to distinguish. Rock sampling was also done on diamond drill holes provided by Avalon Rare Metals Inc. These drill holes intersected unusual rock types that were not encountered surface sampling, during the 2009 and 2010 field seasons. These new rock types are extremely rare and may have serious implications to how the Blatchford Lake intrusive suite formed.

O'Neill, Norm

Université de Sherbrooke Sherbrooke, QB

File No: 12 404 712 **Licence No**: 14936

Region: NS Location: Yellowknife, Aurora College

Sunphotometer measurements at Yellowknife

Measurements of aerosol optical depth (AOD), which is an indicator of the vertical concentration of aerosols, as well as their size, were acquired in Yellowknife, at the AEROCAN sunphotometry network, from January 19 to September 13, 2011. The researchers have not had time to analyze these data, in general, but they did look at some events in detail. The sunphotometer captured the extraordinarily intense smoke (sub-micron aerosol) event at Yellowknife on May 16, 2011, which could be attributed to the fires of northern Alberta. See the Rapidfire image at http://rapidfire.sci.gsfc.nasa.gov/cgi-

bin/imagery/single.cgi?image=Canada.A2011136.1840.2km.jpg

Paradis, Suzanne

Geological Survey of Canada Sidney, BC suparadi@nrcan.gc.ca

File No: 12 404 772 Licence No: 14941

Region: SA, DC Location: Howards Pass deposit; Prairie Creek deposit;

Gayna River deposits; Various sites within the central

Mackenzie Mountains

Hydrothermal event recognition and vectoring to SEDEX ore system in shale basins, Yukon and NWT

Rocks are made of various components and occasionally they have metals in them. The researchers want to find a way to 1) identify some proximal (close) and distal (at a distance) mineralogical and chemical elements, that will indicate that rocks in a sedimentary basin are likely or not likely to be rich in metals; and 2) understand how metals were emplaced in rocks, and how they disperse in the soils and water once the rocks are weathered. This has implications for the land, environment and mineral identification. To achieve the objectives mentioned above, preliminary fieldwork was done during 3 weeks in July and August 2011, and will continue during the summers of 2012 to 2014. The fieldwork consisted of familiarizing themselves with the regional and basin-scale geology; and collecting representative small samples from drill holes stored at one exploration site (i.e., Howards Pass deposit). The samples will be subsequently analyzed in laboratories, using various techniques, to help define the mineral and chemical composition of the rocks. Analysis will start in the fall of 2011, and will continue over the next few years. Microanalytical methods for in-situ measurement of trace elements of mineral phases (apatite, pyrite, feldspar etc.) within host-, footwall, and hanging wall sedimentary rocks will be developed.

Pickart, Robert

Woods Hole Oceanographic Institution Woods Hole, MA United States rpickart@whoi.edu

File No: 12 404 742 **Licence No:** 14839

Region: IN **Location:** The shelf edge, from the US/Canada border to

the vicinity of Banks Island

Assessment of the western Arctic boundary current

Cruise HLY1103 of the US Coast Guard Cutter Healy took place from 2-27 October, 2011. The title of the field program is "Assessing the western Arctic boundary current and its role in the arctic ecosystem and climate change", funded by the US National Science Foundation as part of the

Arctic Observing Network (AON). The project is a collaboration between US and Canadian scientists. The researchers are using a combination of year-round subsurface moorings in the boundary current (deployed upstream in US waters), and seasonal (summertime) shipboard observations, including measurements downstream in Canadian waters. During cruise HLY1103, they successfully deployed all of the moorings, and carried out a hydrographic survey of the boundary current from Barrow Canyon, along the continental slope into Canadian waters to the mouth of M'Clure Strait. The survey consisted of 10 cross-slope transects, using a conductivity/temperature/depth (CTD) package equipped with a transmissometer, fluorometer, and oxygen sensor. Niskin bottles were used for water sample measurements of salinity, dissolved oxygen, nutrients, dissolved inorganic carbon, total alkalinity, oxygen isotopes, and chlorophyll. Velocity measurements were made using the hull-mounted acoustic Doppler current profiler (ADCP).

Pisaric, Michael

Carleton University
Ottawa, ON
michael pisaric@carleton.ca

File No: 12 404 640 **Licence No**: 14908

Region: IN, GW

Location: Blueberry site, north of Inuvik from Noel Lake towards Swimming Point: the Dood Zone in the outer

towards Swimming Point; the Dead Zone in the outer Mackenzie Delta; the Kendall Island Bird Sanctuary; Richards Island; lakes north of Fort McPherson, on the

western edge of the Mackenzie Delta

Examining the impacts of climate change on aquatic and terrestrial ecosystems of the Mackenzie region, NWT

In 2011, the researchers collected sediment from the bottom of lakes near Noel Lake, Husky Lake, north of Fort McPherson, and ~60 km west of Swimming Point. Their objectives are to document the impacts of changing climate on these lakes, especially the impact of thawing permafrost and storm surges. This summer they visited ~10 lakes, to study how algae and insects preserved in the sediment are being affected by these disturbances. Early results of the storm surge project indicate that larger storm surges occur during periods of warmer temperatures, suggesting climate warming may result in more and larger storm surges near the coast. Around Fort McPherson the researchers sampled a lake near Husky Lake with a small stream flowing into it from the mountains. Up in the mountains the permafrost has melted and the ground is collapsing and is being carried by the stream into the lake. In April, the researchers collected sediment from this lake and found that ~20 cm of sediment had been deposited in this lake during the last 10 years, which is a remarkable amount of sediment. In other lakes around Inuvik, it would usually take more than 100 years to accumulate this much sediment.

Quinton, William

Wilfrid Laurier University Waterloo, ON wquinton@wlu.ca

File No: 12 404 570 Licence No: 14874

Region: DC **Location:** The Scotty Creek drainage area, approximately

62 km SSW of Fort Simpson

Understanding and prediction of permafrost thaw impacts on northern water resources

Research at Scotty Creek is focused on 1) understanding the rates and patterns of permafrost thaw, and the physical and biological processes that control it; 2) developing science-based tools to predict the rate and pattern of permafrost thaw over the next 50 years; 3) understanding and predicting the impact of permafrost thaw on ecosystems and water resources; and 4) developing appropriate mitigation strategies. Scotty Creek is typical of the southern margin of permafrost, where permafrost is relatively warm, thin and discontinuous. As a result, permafrost thaw often leads to permafrost disappearance. Over the last half century, permafrost has reduced from about 72% of Scotty Creek to about 40%, and the rate of permafrost disappearance is accelerating. Current research is focused on 1) developing new conceptual and mathematical models that simulate water flow and storage processes at the southern margin of permafrost, 2) developing a new permafrost thaw model that includes the effects of climate warming and human-induced disturbances, such as seismic lines, winter roads and pipelines; and 3) coupling the hydrological model with the permafrost model, to predict the spatial distribution of permafrost and river flow regimes under possible scenarios of climate change and human-induced disturbances. This project hosted a public workshop in Yellowknife (Oct., 2011), that focused on interactive training of new science-based predictive tools, needed to properly manage northern water resources in the wetland-dominated regions with thawing, discontinuous permafrost. This project is part of the new Laurier-GNWT Partnership, and as such collaborates closely with the GNWT and its federal partners, NGOs, First Nations communities and local stake-holders, for the purpose of strengthening the NWT Water Strategy.

Reford, Stephen

Darnley Bay Resources Ltd. Toronto, ON sreford@darnleybay.com

File No: 12 404 745 **Licence No:** 14857

Region: IN **Location:** Darnley Bay Resources land holdings near

Paulatuk

Darnley Bay Resources Ltd. 2010 - 2012 field program

This summary covers the second year of a three-year program of exploration for metals and diamonds in the Paulatuk area. The following fieldwork took place in November 2010 and March-April 2011: (1) Geological Prospecting and Sampling was not carried out, due to snow cover; (2) Claim Staking and Bathymetric Survey was not carried out during the second year; (3) Ground Geophysical Surveys were not carried out during the second year; (4) Kimberlite/Metals Drilling was not carried out during the second year. Results of the drilling in 2010 were received in early 2011 and announced. Three new kimberlite pipes were intersected on the Parry Peninsula, one contained diamonds and numerous indicator minerals. The other two did not have enough kimberlite material for analysis. In addition, xenoliths were extracted from the kimberlite core, for a research project currently underway at the University of Alberta and the Northwest Territories Geoscience Office, focused on mantle studies. The third drillhole for metals was completed in 2010, south of Paulatuk. The hole on a gravity target was terminated after 158 m of overburden (glacial till and unconsolidated Cretaceous sediments) and 84 m of Devonian vuggy dolomite, due to technical difficulties. The same target was drilled again in 2011, but encountered the same difficulties and was terminated at a depth of 242 m, without reaching its target.

Rose, Rachael

University of Calgary Calgary, AB roserm@ucalgary.ca

File No: 12 404 775 Licence No: 14951

Region: SA **Location:** Powell Creek; Shortcut Creek; Little Chicago

Western Imperial formation

Plans were changed, due to circumstances in the field. Rather than multiple days and sites, the researchers visited a single location on one day, July 27 2011. The location was Lac Charrue, approximately 50 km northeast of Little Chicago. Samples were collected for detrital zircon analysis. Detrital zircon analysis of other sandstones in the northern Mackenzie Valley had shown that the sandstone rocks contained sand grains that were initially from either northwestern Alaska or Siberia. Those sand grains were transported from those locations to the northern Mackenzie Valley area, approximately 360 million years ago, when those landmasses were located where the Arctic Ocean is presently located. The researchers aim to improve their understanding of how NW Alaska and Siberia drifted away from northern Canada, during the formation of the Arctic Ocean. Sandstones collected under this research licence were crushed, to separate individual sand grains for analysis. The researchers did identify Alaskan and Siberian sand grains and they will publish those results in a scientific journal.

Russell, Hazen

Geological Survey of Canada Ottawa, ON hrussell@nrcan.gc.ca

File No: 12 404 765 Licence No: 14906

Region: NS, SS Location: MacKay Lake; Beaverhill Lake

Heavy mineral indicator tracing in glacial-fluvial systems

No research was conducted under this licence in 2011.

Skeries, Kristina Queen's University Kingston, ON k.skeries@queensu.ca

File No: 12 404 762 **Licence No:** 14896

Region: SA, DC **Location:** Coates Lake deposit; Bear Twit deposit; Prairie Creek deposit; Howards Pass deposit; Mactung deposit

Geochemical and mineralogical controls on metal dispersal downstream of mineralization in the Mackenzie Mountains. Canada

In July and August of 2011, sampling was done in, and around, the Prairie Creek Mine site. Water and sediment (very fine grained, as well as larger grained) was sampled from the streams. Water and sediment (only fine grained) was also sampled from within the ore stock pile and waste rock pile on site. Over the past year, these samples have been analyzed, using a variety of different methods. Chemical analysis was done for total concentrations of major and trace elements in the water and sediments. The sediments have also been analyzed under a Scanning Electron Microscope and by synchrotron-based techniques, both of which allow the investigator to look at individual grains, to get chemical data for specific spots on those grains, and to figure out which minerals the grains contain. Results from all analyses are still being processed and interpreted. Preliminary results show that there is chemical weathering occurring and that dissolved trace metals are found in very small quantities in the streams. They are often adsorbed onto iron-

containing grains and clays. In the ore stock pile and waste rock pile, metals are found in higher concentrations. Modeling software will be used to try to predict the movement of these metals.

Smith, Sharon

Geological Survey of Canada Ottawa, ON sharon.smith@nrcan.gc.ca

File No: 12 404 657 **Licence No:** 14918

Region: IN, GW, SA, DC **Location:** Jean Marie River; Fort Simpson; Wrigley;

Tulít'a; Norman Wells; Fort Good Hope; Tsiigehtchic; Tuktoyaktuk and locations surrounding these communities

Permafrost monitoring and collection of baseline terrain information in the Mackenzie Valley corridor, NWT

Permafrost monitoring sites throughout the Mackenzie corridor (Inuvialuit, Gwich'in, Sahtú, Dehcho regions) were visited in August and September 2011, to acquire ground temperature and active layer data. Two students and an Inuvik resident were engaged through ARI, to assist with data collection in the Inuvialuit and Gwich'in regions. Data records for 40 monitoring sites established in 2007-2008 were extended, to better characterize the permafrost conditions. These records are helping the researchers understand the natural variability in permafrost thermal and active layer conditions and ensure availability of adequate baseline permafrost information to support land management decisions in the region. Their results show that permafrost in the discontinuous permafrost zone, which covers a large portion of the corridor, is generally warmer than -2° C. Permafrost temperatures generally continue to increase and ongoing maintenance of monitoring sites and data collection is planned to better characterize the impact of climate change on the permafrost environment. A detailed report, including graphical and tabular summaries of data, is currently being prepared and will be sent to relevant organizations in the region.

Snyder, David

Natural Resources Canada Ottawa, ON dsnyder@nrcan.gc.ca

File No: 12 404 548 Licence No: 14855

Region: SA, NS, SS **Location:** Castor lake; Gamètì airport; Hepburn Lake;

Sulky Lake; Lac des Bois; Simpson Lake; Colville Lake;

Kugluktuk

Teleseismic studies in the Wopmay

During 2011, nine seismic stations were removed in the East Arm regions of Great Slave Lake; one remains. One new station was sited on Johnson Point, Banks Island. Seven stations were maintained in the northern Great Bear Lake region, as well as on southern Victoria Island. This project is attempting to define the northwest margin of the Achaean Slave block at great depth, in support of diamond exploration. All active stations successfully recorded more than 100 distant earthquakes, in 2011. Analysis of the final data from the East Arm suggests that mantle rocks, associated with the Slave block, to the northwest, form a wedge of mantle rock that continues as far south as Gardenia Lake at 170 km depth.

Sofko, George

University of Saskatchwan Saskatoon, SK

george.sofko@usask.ca

File No: 12 404 636 Licence No: 14848 Region: IN, GW Location: Inuvik

PolarDARN: The northern hemisphere polar portion of the international SuperDARN [Super Dual Auroral Radar Network] program

The most important achievement in the past year is the successful construction and installation of a new digital beam forming (DDS) system. The system was installed at Inuvik, on a site visit, from 29 November - 3 December 2010. The Inuvik radar is a world-class facility for ionospheric monitoring. When a DDS system is deployed at Rankin Inlet, as well as at the other two Canadianoperated auroral-zone SuperDARN radars near Prince George and Saskatoon, by the end of 2011, the researchers will be able to perform more complicated/useful scanning programs than the other 23 radars in the network. The PolarDARN radars are positioned in the polar cap region, which is highly dynamical and is directly controlled by space weather conditions in near-Earth space. The polar caps are the regions that contain those magnetic field lines that are connected directly between the Earth and the interplanetary medium. It is by studying the direct interaction between the interplanetary medium and the Earth's upper atmosphere that the researchers can begin to understand the influences that phenomena like solar storms have to communications, satellites, astronauts, large power grids at ground level, and the influence of space weather upon weather at the Earth's surface. The PolarDARN radars will be located and orientated so that they will provide the opportunity for research collaboration with the "incoherent scatter radar" (ISR) radar project happening in Nunavut. They will also enhance the ability to do HF radio wave propagation experiments in the polar cap region. For the times when HF radio wave transmission is the only viable communication option for aircraft in the polar regions, understanding the propagation of these waves will be very important. It has clearly been a very busy and productive year for the Inuvik radar team, and the future promises to hold even greater research and collaboration opportunities.

Spence, Christopher Environment Canada Saskatoon, SK chris.spence@ec.gc.ca

File No: 12 404 535 **Licence No:** 14920

Region: NS Location: The Baker Creek basin

Investigations of the water cycle and hydrological processes of the subarctic Canadian Shield

Field activities in 2011 in the Baker Creek research catchment began with spring snow surveys and the activation of climate towers and water level stations in April. There were no people living and working in the research catchment in 2010. Along with the continued remote measurements of meteorological conditions, evaporation, soil moisture and stream flow, a hydrochemistry sampling program, began in 2010, continued through 2011. This program involved sampling stream flow in tributaries and at lake outlets along Baker Creek bi-weekly. Groundwater was also sampled. Samples were analyzed for ions, pH, metals, nutrients and carbon and nitrogen. This work is in support of determining how stream chemistry and frozen ground relate to wetter autumn conditions during freeze up. These research questions are in response to observations, which have shown that stream flow in small subarctic Canadian Shield catchments changed from a predominantly nival (snowmelt) to a combined nival/pluvial (snowmelt and rainfall) regime in the late 1990's. The autumn of 2011 was wet and the 2011 annual peak discharge was in October,

rather than during spring snowmelt. Hydrochemistry and stream flow data during this freeze-up event were collected and are now being analyzed.

Steele, Michael

University of Washington Seattle, WA United States mas@apl.washington.edu

File No: 12 404 757 **Licence No:** 14915

Region: IN Location: Southern Beaufort Sea

UpTempO: measuring the upper ocean temperature of the Arctic Ocean

One UpTempO buoy was deployed from the Canadian Coast Guard ship Amundsen, in August 2011. The ocean thermistors on the buoy failed immediately. A second buoy was deployed later that same month and this one worked successfully. As expected, it drifted with the prevailing currents westward through the southern Beaufort Sea, measuring the fall cooling of the upper ocean. The buoy failed in late October 2011, probably as a result of sea ice impact.

Turetsky, Merritt University of Guelph Guelph, ON

mrt@uoguelph.ca

File No: 12 404 776 **Licence No:** 14955

Region: DC **Location:** Streams and rivers along the Mackenzie

Highway from the Alberta border until it meets the Liard

Highway near Fort Simpson

Composition of natural dissolved organic carbon in streams along latitudinal transect

In August 2011, the researchers sampled 65 rivers in the Northwest Territories and northern Alberta, during a 3 day road trip, taking 2x50 ml whole water samples at each river. The purpose was to assess the composition of dissolved organic carbon for rivers within and outside the permafrost region. Sampling was successful and analysis back in Guelph, Ontario, was also successful. The researchers found that the aromaticity of dissolved organic carbon decreased with latitude of the sampling river, however, they are still looking into possible explanations for this pattern. Aromaticity of dissolved organic carbon regulates how well it supports microbial respiration, but also how well UV-light is attenuated. Possible explanation include the presence or absence of permafrost, extent of peat lands along the transect, differences in water temperature, or differences in how much groundwater that enters the rivers.

Urbanic, Jane Challen

Environment Canada Burlington, ON jane.challen-urbanic@ec.gc.ca

File No: 12 404 741 **Licence No**: 14856

Region: IN, GW, DC, NS Location: Inuvik; Paulatuk; Ulukhaktok; Tsiigehtchic; Fort

Providence and Rae

Arctic wastewater research

From May to October of 2011, Environment Canada conducted sampling at wastewater systems in Canada's Arctic region for the third year. Extensive sampling was conducted at the lagoons in Paulatuk, Inuvik, Tsiigehtchic, and Rae. Environment Canada also visited Ulukhaktok, to remove thermistors that were placed in the wastewater lagoon the year prior. The sewage lagoons in Paulatuk, Inuvik and Tsiigehtchic discharge continuously, during the open-water season. Sampling was done in these communities in June. Paulatuk and Inuvik were also sampled in September. A controlled discharge of Rae's sewage lagoon began in August and samples were collected throughout the discharge. At each community, samples were taken of raw sewage (influent), lagoon effluent and sludge. Samples were also taken of the overland flow (wetland) prior to the receiving environment. Solids in the Paulatuk and Inuvik lagoon effluent were lower in June than in September. Higher solids are expected in the fall, because algae are discharged with the effluent. Solids concentrations were high in Rae (>50mg/L), due to sediment flowing through the discharge pipe. In most cases, solids concentrations were reduced by >90% through the lagoon. Removal of organics through the lagoon systems was similar. The results show that wetlands also contribute to the overall treatment of the wastewater. In most cases, wetland treatment reduced solids and organic concentrations by an additional 20% from the lagoon effluent. Community reports containing detailed data will be sent to the ARI and the communities by November 30, 2011. In 2012, Environment Canada plans to return to Paulatuk and Rae. New sites will likely include Edzo, Fort Providence and Tuktoyaktuk, funding permitting.

Vonk, Jorien

ETH-Zurich
ERDW - Geologisches Institut
Zurich
Switzerland
jorien.vonk@erdw.ethz.ch

File No: 12 404 764 Licence No: 14900

Region: IN, GW Location: Mackenzie River Delta

Deltaic lake sediments as recorders of past carbon export from arctic river drainage basins. The Mackenzie River is the largest source of river sediments and carbon to the Arctic Ocean. Its delta, the Mackenzie Delta, is the second largest river delta in the world, and is covered with thousands of small, shallow lakes. During the period of ice break-up in May-June, when water levels are high, these lakes receive and store a lot of river sediments. To determine the spatial spread of sediment delivery to the delta, samples were collected from six lakes and delta channels all across the Mackenzie Delta. In addition, sediments were collected from the Mackenzie main channel. Sampling was performed from helicopter floats, on June 3rd 2011. The sampling results are still being processed, but preliminary results show that lakes in the western part of the delta show a different sediment pattern than in the eastern and northern part of the delta. This can probably be explained by the inflow of the Peel River, draining into the Mackenzie Delta in the southeast. If river discharge and/or the timing of the ice break-up changes due to climate change, this will most likely also change the sediment distribution into the delta lakes.

Wang, Zhaohui (Aleck)

Woods Hole Oceanographic Institution Woods Hole, MA zawang@whoi.edu

File No: 12 404 740 **Licence No:** 14838

Region: IN Location: Mackenzie River, near Inuvik

Towards long-term monitoring of the CO₂ system in arctic rivers

This year, the researchers have continued to collect monthly river samples from Mackenzie River, near Inuvik, NT, Canada. All sampling was very successful. The researchers have collected a total of 18 water samples. All samples will be measured for total dissolved inorganic carbon (DIC) concentration and alkalinity. About half of the samples have been processed, and the results are promising. The researchers will continue processing the samples, and maintain monthly sampling into 2012. The available data show significant seasonal changes in DIC concentrations and alkalinity in the east channel of Mackenzie River at the Inuvik dock. From DIC concentrations and alkalinity, they can calculate pH and partial pressure of carbon dioxide (pCO2) in water. The results also show large seasonal changes of pH and pCO2. They also found that there is a significant positive correlation between monthly DIC or alkalinity concentration and mean basin-wide temperature: higher temperature, higher DIC or alkalinity concentration in Mackenzie River water. The researchers are in the process of data analysis and developing a manuscript that describes carbonate chemistry in the Mackenzie River basin.

Wolfe, Stephen

Natural Resources Canada Ottawa, ON swolfe@nrcan.gc.ca

File No: 12 404 549 Licence No: 14854

Region: NS **Location:** West of Yellowknife; along the Ingraham Trail, east of Yellowknife; the Baker Creek Watershed, north of

Yellowknife; along the Tibbit to Contwoyto winter road, based out of the Lockhart and Lac de Gras facilities

North Slave permafrost study: Characterizing and predicting discontinuous permafrost for climate change adaptation

Fieldwork was conducted between June and September, 2011, in the Great Slave region, along Highways 3 and 4, and the Tibbitt to Contwoyto winter road. Permafrost cores, ranging from 1.2 to 7.3 meter depth, were obtained from six sites in peatland, spruce and birch forest settings. Thaw depths, soil types, visible ice moisture contents, and bulk densities were determined, with grain size, water geochemistry and geotechnical tests to be determined. These data will be used as part of a graduate thesis study and for understanding geotechnical conditions associated with permafrost soils in the area. Temperature data continue to be collected and monitored at a number of sites including: active layer temperatures from birch, spruce forest and peatland sites; ground temperatures from burn sites and peatland, birch and spruce forest sites; and air temperature sites; shallow-water; and lake-bottom sites. These data are used to understand potential climatic gradients and the effects of water on local permafrost conditions. PCSP-supported helicopter surveys were undertaken, to validate remote sensing interpretations of surficial geology and vegetation cover mapping in NTS map sheets 85I, J, O and P, 86A and 75M.

Worthy, Douglas

Environment Canada Toronto, ON doug.worthy@ec.gc.ca

File No: 12 404 760 Licence No: 14871

Region: NS Location: NWT Power Corporation's Frank Channel

Substation, located near Behchokò

High-precision atmospheric carbon dioxide and methane measurements at Behchokò, NWT

Environment Canada conducts atmospheric measurements of carbon dioxide, methane and other GHGs, from coastal, interior and arctic regions in Canada. The researchers' goal is to provide high quality data, to permit and improve their ability to estimate sources, such as carbon dioxide, from the burning of oil and gas, as well as natural sources, such as methane emissions, from wetlands. Of particular interest, and thus the reason behind the measurement program at Behchokò, is in regards to the general concern that the Arctic may undergo drastic changes, if the arctic warming trend continues. This is especially true for methane, since arctic methane sources may be widespread. The climate feedbacks from such changes may potentially be very large. Because Behchokò is located in a discontinuous permafrost zone, and within the tree line transition zone, this site provides an excellent platform for observing potential emissions of carbon dioxide and methane from the high terrestrial and subarctic region. The data records are too short to make any conclusive statements at this time, but the researchers are confident that these records will be invaluable, if significant emissions of methane and carbon dioxide from the subarctic do start to occur.

Wrona, Frederick

University of Victoria Victoria, BC wrona@mail.geog.uvic.ca

File No: 12 404 711 Licence No: 14947 Region: IN Location: Noell Lake

Noell Lake ice study - hydro-ecological responses of arctic tundra lakes to climate change and landscape perturbation

The objective of this research is to improve knowledge on lake ice and its effect on the food webs/productivity of small arctic lake systems, in order to better predict changes that could occur under changing climate. In late-September 2010, prior to freeze-up, an automated ice buoy and subsurface mooring system was deployed in Noell Lake, for continuous monitoring of weather conditions, lake ice cover (i.e., formation, growth over winter, breakup in spring), light penetration into the lake through ice in winter, and water quality. The installation was successful and data was collected by the system, throughout the 2010-11 winter. In late June/early July 2011, after the ice was gone, the buoy and mooring system was removed from the lake for servicing. Due to some logistical challenges, the monitoring system was not in place to capture the first part of the summer season. In September, the buoy system was redeployed and it is expected that the system will provide continuous data (winter, spring, summer and fall), in 2011-12, from Noell Lake. These data are allowing the researchers to examine lake ice and its effects on the food web/productivity through the winter, as well as the character of food webs/productivity during the ice-free season.

Wrona, Frederick

University of Victoria Victoria, BC wrona@mail.geog.uvic.ca

File No: 12 404 711 Licence No: 14832

Region: IN Location: Mackenzie Delta Lakes

Amendment - Hydro-ecological responses of arctic tundra lakes to climate change and landscape perturbation

Through the researchers' on-going investigations on small arctic ponds/lakes, it became evident that some of the small pond/lake food-webs may include very small fish, such as stickleback and pond smelt, as top-down controls on the food-web. In 2009, this research component was added to their overall research program, to determine if any of the small ponds/lakes they study contain fish. It was found that 7 of the 11 lakes visited did indeed host fish. In 2010, they visited some of these lakes again, to collect additional samples of some species of fish and increase their overall sample size. They also visited 15 "new" ponds/lakes, not sampled in 2009, that drain into Noell Lake, to enhance their investigations. Although the researchers were licensed to do so, it was decided that they did not need additional information on fish, and no fish collections were taken in 2011. All their information on fish, including which fish types were found in which lakes and which lakes did not host fish, are being made available as a DFO Data Report.

Wrona, Frederick University of Victoria Victoria, BC wrona@mail.geog.uvic.ca

File No: 12 404 711 **Licence No**: 14833

Region: IN Location: Mackenzie Delta Lakes

Hydro-ecological responses of arctic tundra Lakes to climate change and landscape perturbation

An overall goal of this work is to better understand the effects of changing climate, using permafrost degradation as an analogue for changes under a warming climate, on the supply of nutrients to tundra lakes, and in turn, its effect on lake geochemistry, as well as the biological communities within the lakes. Preliminary results indicate that the high inter-annual and seasonal variability in temperature controls key hydrological processes, such as ice-on and ice-off dates and the timing of the spring snow melt. Additionally, inter-annual variability in snowfall and rainfall has significant controlling effects on the magnitude of the spring snow melt and summer surface flow generation, respectively. This apparent variability in both climate and hydrology affects the geochemistry of both surface flow and lake water. Most noticeable, in spring, when snow melt water has relatively low ionic concentrations, there is a dilution effect on surface flows into the lakes, and in turn, the lake water. However, runoff from shoreline permafrost slumps exhibit relatively high concentrations of ions than from other catchment sources, and contributes to relatively higher concentrations in the lake water. Data analyses, interpretation of results, and assessment of potential effects on aquatic biology (food-webs) is ongoing.

SOCIAL SCIENCES 2011

Abele, Frances

Carleton University
Ottawa, ON
frances abele@carleton.ca

File No: 12 410 857 **Licence No:** 14879

Region: NS Location: Yellowknife; Behchokò

Housing and being homeless in Yellowknife

This research project began in 2009, and thus far has resulted in three publications. They are as follows: (1) A 2010 article, co-authored by Frances Abele, Nick Falvo and Arlene Hache: http://homelesshub.ca/%28S%2820gliz35wmmqflb1dxypfz45%29%29/Library/Homeless-in-the-Homeland--A-Growing-Problem-For-Indigenous-People-in-Canadas-North-49863.aspx; (2) A 2011 policy report on homelessness, written by Nick Falvo, in partnership with the Centre for Northern Families: www.homelesshub.ca/Yellowknife; (3) A 2011 peer-reviewed chapter on government-assisted housing in the Northwest Territories (NT), which appeared in How Ottawa Spends (McGill-Queen's University Press): www.homelesshub.ca/NWT. Dr. Abele and Mr. Falvo still intend to collaborate on an historical article on government-assisted housing in the NT. It would focus on the period from roughly 1945 to 2000.

Baumann, Britt

University of Waterloo Kitchener, ON bbaumann@uwaterloo.ca

File No: 12 410 895 **Licence No:** 14930

Region: IN, NS **Location:** Inuvik; Yellowknife; Tuktoyaktuk

The impact of the priest decline on the Canadian Roman Catholic Church

This doctoral dissertation examines the impact of the decline in the number of priests within the Canadian Roman Catholic Church. Part of this project examines the priest shortages that exist in remote areas in northern Canada. Interviews with Roman Catholic activists reveal that many Canadian parishes are experiencing a shortage of priests. There are reports that some Roman Catholic communities in the north may only see a priest once or twice per year and that Roman Catholic lay persons are attending to the spiritual needs of parishioners. As well, some northern priests must cover multiple parishes. These priests spend a great deal of time travelling between parishes to administer sacraments requested by northern Roman Catholics. The next phase of this project will take place at the Diocese of Mackenzie-Fort Smith. The researchers will gather qualitative data on the lived experiences of northern Roman Catholics as they develop what they refer to as "a new way of being church" that no longer depends on priests from the south. Data

will be collected through interviews with the Diocese administrators, as well as people that are involved in Roman Catholic lay ministry. This research is a valuable addition to the sociological literature as it will improve people's understanding of the constituency of the Canadian Roman Catholic Church. It is imperative to include the voices of northern Roman Catholics in any study of the Canadian Roman Catholic Church. Additionally, northern Roman Catholics have had an effect upon the Roman Catholic Church due to both their challenges and their innovations. This study will provide the researchers with specific knowledge about the unique features of northern Roman Catholic practice, which is increasingly focused upon aboriginal leadership.

Cash, Penny

University of British Columbia, Okanagan Kelowna, BC penny.cash@ubc.ca

File No: 12 410 886 Licence No: 14888 Region: NS Location: Yellowknife

Quality workplace environment

Quality workplace environments are known to be a positive determinant in the recruitment and retention of nurse educators. This project is designed to survey nurse educators in all public postsecondary schools of nursing in British Columbia (BC) and the Northwest Territories (NT) to establish what nurse educators consider to be important elements in their work environment and to determine if these are experienced. This project builds on a pilot project (H08-0021), which involves the development of an instrument to measure quality workplace environments, as well as the pilot testing of that instrument. This instrument was used in all of the surveys. All nurse educators in public post-secondary schools of nursing in BC and NT were invited to participate in an online survey. Using a mixed methods approach, descriptive statistics will be used to examine the survey responses. Qualitative responses will be examined for themes and used to triangulate the statistical findings. This work is ongoing. Next Steps: the researchers have proposed an amendment to collect data through focus groups with nurse educators in BC and NT. In the survey, subjects had time and space restrictions on their qualitative comments. Focus groups will help to further inform the themes emerging from the qualitative data. The researchers are planning 4 to 8 focus groups, which will involve all nurse educators currently employed in public postsecondary education institutions in BC and NT. This will include nurse educators teaching in programs ranging from Health Care Assistant Certificates to Doctoral Programs.

Delormier, Treena

Centre for Indigenous Peoples' Nutrition and Environment Ste. Anne de Bellevue, QB treena@treena.ca

Traditional and market food: focus on fish consumption in Tulít'a, NWT

In June 2010, the community of Tulít'a raised concerns about a public health advisory to eat less lake trout from Kelly Lake due to high levels of mercury measured in them. The community requested a study to learn if the fish they were eating could harm them, while also keeping in mind the nutritional and cultural benefits of fish. Sixty-seven community members took part in the study, twelve were children. Participants gave a hair sample for mercury testing and answered questions about their fish eating habits. Seven community knowledge holders and a public health officer gave interviews on the cultural meaning of fish as a traditional food, as well as the effects

of the advisory on how people view eating fish. Hair mercury tests showed that people were consuming very low amounts of mercury. Only two adult men required follow-up. Dietary results show that people eat more fish in summer than winter. Whitefish and lake trout are eaten most often, but grayling, loche, inconnu and cisco are eaten too. Fish provides important nutrients. Interviews noted that traditionally fish was a dependable food source, ensured the survival of the Dene & Métis peoples, and is also deeply culturally important. Fish is still preferred as a food today and traditional food harvesting needs to continue. Negative reaction to the public health advisory on mercury was largely due to the way information reached the community. This study has changed the way public health advisories are done in the Northwest Territories.

Denning, Bryany

Institute of Circumpolar Health Research Yellowknife, NT bryanydenning@gmail.com

File No: 12 410 896 Licence No: 14933 Region: NS Location: Yellowknife

Street outreach community consultation

The purpose of this study was to describe the health and addictions assets in the downtown core of Yellowknife, NT, in order to better address the high hepatitis C (HCV) and sexually transmitted infection (STI) rates observed there, especially infections transmitted through illicit drug use. Over the summer of 2011, data was collected through focus groups with nineteen street people, a Photovoice workshop with eight street people and seven one-on-one interviews with service providers in the downtown core. A number of themes emerged through the consultation process. including: needle exchange; non-beverage alcohol use; mixed messages to youth; sex trade; interagency coordination; and need for outreach services. The community consultation results indicate that the high HCV and STI rates are likely not related to illicit drug use, such as transmission by crack cocaine pipes. However, links may be found through further investigation into the following areas: (1) the development of harm reduction mechanisms that address the needs of the street population; (2) non-beverage alcohol use and how it impacts the HCV and/or STI rates in Yellowknife; (3) youth-based social marketing tools that use messaging consistent with local laws and/or policy; (4) how the sex trade in Yellowknife can be effectively addressed; (5) how communication, coordination and collaboration can be increased between service providers and service agencies in Yellowknife; and (6) defining the appropriate outreach measures to reach Yellowknife's marginalized street population.

Duran. Nelida

UCLA School Of Public Health Dallas, TX United States nduran@ucla.edu

File No: 12 410 881 Licence No: 14973 Region: NS Location: Ndilo; Dettah

The effects of a changed climate and environment on the nutrition and health of Dene First Nations

No research was conducted under this licence in 2011.

Duthie, Andrew

Rescan Environmental Services Ltd.

Yellowknife, NT aduthie@rescan.com

File No: 12 410 891 **Licence No:** 14912

Region: DC, NS, SS **Location:** Courageous Lake

Courageous Lake project - Social and economic sciences research

The objective of this study is to characterize the social and economic environment of the proposed Courageous Lake Project and land use in the area. The 2011 environmental baseline program collected data from various government, academic and other public sources to characterize the social and economic environment of the proposed Courageous Lake Project. Contact with various First Nations and the Métis was initiated to discuss research methodology, but due to various delays no survey work was completed. Meetings are anticipated to occur in November. A desk-based portion of the study compiled information from available sources, including government databases regarding existing land use activities to help identify land users and tenure holders, including commercial and recreational land users, as well as aboriginal groups. The land management context for the study area was investigated, including the identification of relevant land/resource management strategies and objectives.

Fresque, Jennifer

Wilfrid Laurier University Waterloo, ON fres3130@mylaurier.ca

File No: 12 410 885 Licence No: 14869 Location: Fort Resolution

Linking place identity, environmental change and adaptation in the context of changing water conditions in Fort Resolution, NT

Preliminary findings show that water in/around Fort Resolution is perceived by community residents to be changing dramatically. Water levels in Great Slave Lake, Slave River and Delta and other important waterways have declined by several inches in the last few years. Residents noted that water looks, tastes and smells different in many traditional areas. Many expressed mistrust of water quality, both on the land and in the community, and take alternate measures for drinking water consumption. People are connected to places in the South Slave Region because of heritage, what the land provides, a sense of identity, social connections, and for well-being. Many places are now inaccessible due to declining water levels, making it harder for trappers to engage in traditional subsistence activities. Many residents described feelings of loss because places important to them are changing. There is sadness and frustration that people can no longer use the land and water in the same way as before. People expressed frustration that many of the impacts on regional waters are coming from outside the community, and often beyond territorial borders. It is imperative that steps are taken to ensure community concerns about water are heard and addressed appropriately at multiple levels.

Gagnon, Graham

Dalhousie University Halifax, NS ggagnon@dal.ca

File No: 12 410 890 **Licence No:** 14882

Region: IN, GW Location: Ulukhaktok; Aklavik

Success factors for small sustainable drinking water systems

Water is used in the home for cooking, cleaning, bathing and drinking. Everyone needs to have access to safe water in order to be healthy. In most communities in Canada, water is treated and then delivered to homes by a pipe or by truck. Small communities (those with fewer than 5,000 people) often face a number of challenges when providing safe water. These include, but are not limited to, the cost of water treatment and delivery. The purpose of this study is to understand what these challenges are, and to understand how some small communities have overcome these challenges in order to provide safe water. To do this, interviews were conducted with decision-makers, water operators, and community members, in seven communities from across Canada, including one community in the Northwest Territories (Ulukhaktok). The research found that the community had improved their water treatment so they could meet new regulations set by the territory. Although people were happy with their water supply, most preferred to gather water from the land. This information will be used to better understand issues that small communities face with respect to water quality and health.

Hampton, Mary

University of Regina Regina, SK mary.hampton@uregina.ca

File No: 12 410 906 **Licence No:** 14988

Region: IN, GW, NS, SS, SA, DC Location: All communities

Rural and Northern Community Response to Intimate Partner Violence

For the year 2011, the Northwest Territories team of the Rural and Northern Community Response to Intimate Partner Violence (IPV) accomplished several goals. They successfully hired two undergraduate students from Aurora College as research assistants. One student is from the Nursing Program and the other student is currently enrolled in the Social Work Program. The team is very satisfied with their work productivity, learning achievements, and contributions to the project. Furthermore, these students will be retained for the 2012/2013 academic year. One of the team's major accomplishments this year was the completion of an environmental scan of supports and resources located throughout the NT (e.g., shelters, victim services, treatment programs, justice system supports/courts, health program, etc.). This data was submitted to the project geographer who will integrate the data into the geographic information system (GIS) mapping. The team was also able to establish a relationship with a G Division RCMP member, whose work focuses on domestic violence. He and the national office have helped the researchers collect incidence data across the NT. This will also be integrated into the GIS mapping. For 2012, the NT team will have a face-to-face project meeting with the entire planning team, representing all 4 jurisdictions (Alberta, Saskatchewan, Manitoba, and Northwest Territories). This will determine the steps for the project's second year including the completion of GIS mapping, a review of the results, and begin interviews with front-line service providers across the NT who have been identified by the team's aboriginal project sub-committee as having expertise in IPV (e.g., directors of shelters, justice workers, elders). The researchers are in the process of seeking NT representation on the aboriginal project sub-committee.

Hodgkins, Andrew

University of Alberta Edmonton, AB hodgkins@ualberta.ca

File No: 12 410 649 Licence No: 14883 Region: IN, GW Location: Inuvik

Vocational education and training partnerships in northern Canada

In November, the researcher visited Inuvik for 10 days and interviewed students at Aurora Campus who were taking Aboriginal Skills & Employment Partnership (ASEP) funded trades programs. Stakeholders related to education, training, and employment were also interviewed. A total of 17 interviews occurred at this stage. Follow-up interviews with the students will be carried out in the spring 2011.

Levitan, Tyler
Carleton University
Ottawa, ON
tyler levitan@hotmail.com

File No: 12 410 900 **Licence No**: 14968

Region: NS Location: Yellowknife; NDilo; Dettah; Behchokò

The Canadian state's relation to Impact and Benefit Agreements in NWT

This research looked at the Canadian state's relationship to Impact and Benefit Agreements (IBAs) signed over to the diamond mines in the Northwest Territories (NT). Since these agreements are bilateral and take place outside of the regulatory regime within the NT, it is important to understand better what interests the state has in these agreements. Interviews were conducted with federal and territorial government officials, leaders within the Akaitcho Treaty 8 First Nations and the Tłįchǫ First Nations, consultants, and lawyers. Many of the interviews were conducted in and around Yellowknife. The research concluded that the Canadian state has been downloading its responsibilities onto third parties through these agreements. Therefore, the state is being relieved of it responsibilities in the consultation and accommodation of potential infringements to aboriginal rights and title. Through abstaining from regulating these agreements, the state has largely given way to the forces of the market to dictate the negotiation, signing, and implementation of these agreements. It is through these agreements that indigenous communities in Canada can be further incorporated into the capitalist system, through increased participation in the labor market and the emergence and growth of aboriginal-owned and operated businesses. This suits the interests of the state by making these communities more self-reliant.

Little, Lois

Lutra Associates Ltd. Yellowknife, NT xlutra@ssimicro.com

File No: 12 410 306 Licence No: 14926 Region: IN, GW, NS, SS, SA, DC Location: All communities

Evaluation of early childhood development training

The evaluation was intended to identify ways to improve the quality of training. Input was gathered from almost 120 early childhood students, practitioners, educators, and leaders in the field. It is clear that tinkering with the existing training program offered by Aurora College is not enough. The GNWT Department of Education, Culture and Employment and Aurora College need to work together to deal with the issues that impact on attracting, retaining, and training northerners for early childhood work. These issues include: low wages and benefits, low value placed on this work, job insecurity, and inequities among early childhood workplaces. The evaluation calls for changes to: existing early childhood training, in terms of expanding delivery modalities; establishing partnerships to deliver accredited training; implementing prior learning assessment and recognition; and linking early childhood, aboriginal language, and teacher education training.

The evaluation also recommends a professional early childhood association, occupational standards, credentialing and registering processes, standardized pay structure, and more focus on student supports, promotion, and performance monitoring.

Loovers, Jan Peter

Royal Anthropological Institute and Goldsmith University of London London, England p.loovers@abdn.ac.uk

File No: 12 410 667 **Licence No:** 14893

Region: GW Location: Fort McPherson; Inuvik; Base camp on Peel

River

Tracing trails with Gwich'in: Poetics, well-being, memory, and land in circumpolar Canada This Royal Anthropological Institute Urgent Anthropology Fellowship research has illustrated that there is an intimate relation between land, poetics, well-being, and memory. This research has shown that 'language loss' between and within different generations is complex and nuances need to be made. Language revitalization projects need to include such complex dynamics. The programs for language revitalization can be threefold: on the land, in the community with Tukudh Bible classes, and in the community at school and in homes. Taking ownership of the language, like Gwich'in language workers and elders emphasize, will play an integral role in this. With Gwich'in elders passing away, work on the Tsii Deii language remains pivotal. Recording elders singing Gwich'in hymns and reading the Tukudh Bible needs to be continued. Gwich'in people continue to live life out on the land – hunting, fishing, trapping, picking berries and travelling. Large-scale resource extraction developments in the Peel River Watershed have been a matter of great concern to the Gwich'in. These developments could jeopardize life out on the land, and subsequently, the language revitalization, well-being, and memory of the Teetl'it Gwich'in people.

Luig, Thea University of Alberta

Edmonton, AB tluig@ualberta.ca

File No: 12 410 892 **Licence No:** 14911

Region: GW Location: Fort McPherson

Collaborative research on community well-being

This research on the role of volunteering, sewing, and human-land relationships in community well-being is conducted in collaboration with the community of Fort McPherson and the Gwich'in Social and Cultural Institute. The methodology of this qualitative study involves apprenticing with elders and community members in the three focus areas, conducting life-story and semi-structured interviews, as well as documenting these activities on video. The theme emerging from the first five months of research concerns the balance of the capacity to care for others, either individually or collectively, as expressed in volunteering time and resources, and the capacity to care for oneself as expressed in hunting, snaring, getting wood, sewing, cooking, being physically fit, and pursuing education. The importance of this balance for well-being is recognized more easily while staying on the land. A project collaboration with the youth council using photography for the creation of digital stories and posters to explore the topic of well-being is in the planning stages and will begin in January 2012. The second stage of the project will focus on the recording

of life stories, the completion of the youth photo project, and the production of written and visual material for knowledge sharing in the community.

Mair, Heather University of Waterloo Waterloo, ON

hmair@uwaterloo.ca

File No: 12 410 902 Licence No: 14991 Region: SS Location: Hay River

Tourism development plan for Hay River

This project involved working with members of the community of Hay River to update their tourism development plan. The researchers' interviews and observations indicated that the community possesses many opportunities to become a go-to destination for travelers who are already in the north. A strategic plan and two presentations about the plan were made to the community about the researchers' findings, including recommendations for future development.

Noble, Bram

University of Saskatchewan Saskatoon, SK b.noble@usask.ca

File No: 12 404 726 **Licence No:** 14865

Region: IN Location: Inuvik; Tuktoyaktuk

Strategic environmental assessment roles and stakes in arctic oil and gas exploration and development

Canada's Beaufort Sea is rich in oil and gas resources. Currently, 'environmental impact assessment' is the tool of choice for assessing and managing the impacts of development in the offshore. The problem is that this approach looks only at each project, one at a time, and does not address the total or cumulative impacts of offshore development on the ecosystem or on communities. Neither does it plan for development, explore different development options, or identify the potential impacts of each, in order to determine how best to move forward in the planning and development of the offshore. In 2004 the Inuvialuit Game Council wrote to the federal Minister of the Environment, requesting a more regional and 'strategic environment assessment' of future energy development in the region. Although the Beaufort Region Environmental Assessment (BREA) was recently launched, its focus remains, like previous efforts, on data collection, as opposed to regional planning for the future. Norway, the United Kingdom and Atlantic Canada all have formal systems for strategic environmental assessment offshore, but no such system exists in Canada's Arctic. This research examined government, industry, environmental, and Inuvialuit perspectives on the need for, benefits of, and risks associated with strategic environmental assessment in the Beaufort Sea. Results indicate a number of opportunities, including: improved regulatory efficiency for proponents; better regional science and planning practices; improved northern influence over future development; an opportunity to assess cumulative effects; more meaningful project-based assessment; and greater certainty for industry. At the same time, there are a number of perceived risks, including: foregoing anticipated development opportunities; the loss of flexibility in decision making; adding another layer of bureaucracy; the uncertainty of a new approach.

Oehler, Alexander

University of Northern British Columbia

Inuvik, NT oehler@unbc.ca

File No: 12 410 897 Licence No: 14956 Region: IN Location: Inuvik

Inuvialuit language and identity: perspectives on the symbolic meaning of Inuvialuktun in the Canadian western Arctic

The purpose of this study was to examine how Inuvialuit beneficiaries in Inuvik felt about the importance of Inuvialuktun, especially in relation to their own cultural identity. This information was intended to help Inuvialuit language planners and instructors in promoting Inuvialuktun. The study consulted 45 Inuvialuit beneficiaries between the ages of 6 and 59. Data was gathered through interviews, group meetings, and questionnaires conducted with individuals who were part of the college, learning centre, secondary school, and the community at large. The study identified some separation in the beliefs held by individuals about their heritage language. While some saw Inuvialuktun as a pillar of their cultural identity, others saw it merely as a benefit, while favoring land-ties as a primary sign of their cultural identity. Moreover, learning desire fluctuated according to age, peaking in young adults who were building identities to pass on to younger generations. The data also confirmed that Inuvialuktun had to compete with English on unequal terms, and within a generally accepted hierarchy upheld by economic demands. Although Inuvialuktun was held in high esteem, most people were satisfied knowing only a few words, as this connected them to their heritage. Several recommendations for language revitalization were made, based on participants' insights.

O'Keefe, Jeffrey

Aurora College Fort Smith, NT jokeefe@auroracollege.nt.ca

File No: 12 410 891 Licence No: 14905

Region: IN, GW, SA, DC, NS, SS Location: South Slave; Dehcho; Tłycho; Sahtú; Beaufort-

Delta; Yellowknife

Teacher performance appraisals: A tool for teacher growth and improvement

The objective of this qualitative research is to explore how teacher performance appraisals contribute to teacher growth and improvement over time in the Northwest Territories (NT). Teachers' perceptions about the value, benefit, and purpose of teacher performance appraisals are being analyzed. This study attempts to generate data about the amount of time, energy, and effort that teachers put into annual performance appraisals and how teachers act upon the recommendations and feedback provided by the principal. Interviews were conducted with 24 teachers in the K-12 education system in the NT. The interviews consisted of several open-ended questions designed to have teachers share their perceptions about their own experiences with performance appraisals and the motivational value of these appraisals. The subjects were selected to include teachers with varying amounts of experience as teachers in the NT and to ensure that they have experience with teacher performance appraisal practices in the NT. Teachers interviewed came from the South Slave, Dehcho, Tłլcho, Sahtú, Beaufort-Delta and Yellowknife regions. The interviews have all been completed, and data from these interviews are being compiled and analyzed at the time of this summary. It is anticipated that data analysis will continue into early 2013 with research being completed by the end of 2013.

Parker, Aliana

University of Victoria

Victoria, BC alianaparker@gmail.com

File No: 12 410 904 **Licence No:** 14977

Region: GW, NS **Location:** North Slave; Gwich'in Territory

Learning the language of the land: The representation of land in web-based indigenous language education

This research focused on connections between indigenous languages and land, incorporating a synthesis of current literature, interviews with indigenous language and culture experts, and a survey of indigenous language education websites. Essential ties between land and language were demonstrated to be integral parts of indigenous cultures. These ties were explored in the framework of an intimate relationship with the land that involves living on the land, learning from the land, belonging to the land, and respecting the land. The current situation of language, territory and culture loss that indigenous communities face has influenced that relationship in many ways, affecting all aspects of indigenous life and culture. These ties between language and land were then considered in an educational context through a survey of 14 indigenous language education websites from within Canada and the United States. The survey revealed that all aspects of the relationship to the land, as described above, appear in different ways on the websites, and that there are many opportunities for more fully representing land in online education. The research showed that indigenous languages have a complex and deep connection to the land that is essential to indigenous culture and that plays a significant role in online language education.

Pearce, Tristan

ArcticNorth Consulting Guelph, ON tristanpearce@gmail.com

File No: 12 410 650 Licence No: 14843

Region: IN Location: Aklavik; Ulukhaktok; Paulatuk

Adaptation planning for climate change in Ulukhaktok, Paulatuk and Aklavik, NWT

This project worked with community members in Paulatuk, Ulukhaktok and Aklavik to continue adaptation planning for climate change. Workshops were held to identify how climate change is affecting the communities and adaptation options. Opportunities to include adaptation in community plans/decision making processes were identified and adaptation actions were prioritized. A landscape scientist worked with community public works and housing to address climate risks affecting community infrastructure. Community adaptation plans were produced for each community, as well as landscape hazard maps for Ulukhaktok and Paulatuk. In Ulukhaktok, a workshop on cruise-ship tourism was held to document community concerns and opportunities related to increasing cruise-ship tourism. An oral history project was developed in partnership with the Inuvialuit Cultural Resource Centre and Helen Kalvak School to help preserve, organize and make oral history accessible within the community (www.nauvikhaq.com). In Paulatuk, where food insecurity was highlighted as a priority concern, a community kitchen series was piloted in collaboration with the community counselor and local volunteers. The series presented an opportunity for food-insecure families to learn and prepare new, nutritious recipes using affordable ingredients (ingredients were free for participants), as well as a space to discuss food security challenges and pose nutrition questions.

Rawluk, Andrea

University of Alberta Edmonton, AB ajrawluk@ualberta.ca

File No: 12 410 859 Licence No: 14878 Region: IN, GW Location: Aklavik

Intergenerational resilience in Aklavik, NWT

In February 2011, Andrea Rawluk returned the preliminary results from her Masters research to Aklavik. Interview transcripts were returned to everyone that participated in the project. As well, Rawluk presented the preliminary results to the Aklavik Renewable Resource Council (RRC) on March 10, 2011. The results were also shared with anyone who was interested at a combined community workshop and community feast held at the Moose Kerr School. The preliminary results suggest that Gwich'in and Inuvialuit elders define resilience similarly to other indigenous cultures, whilst offering additional perspectives. Fewer youth reported having traditional language, knowledge and spirituality than elders, but expressed a desire to learn them and described spiritual experiences. All generations had similar perspectives about what changes were negative and positive for the community and the land and how they would like to see the future of the community. At the workshop, Rawluk received feedback from some community members and decided to ask three community elders to participate in longer interviews during her stay. In these interviews, the elders described that love, spirituality, and patience were at the foundation of resilience for the people of their community. Since the sharing of the results with Aklavik, the thesis has been completed.

Reinfort, Breanne University of Manitoba Winnipeg, MB b.reinfort@gmail.com

File No: 12 410 852 Licence No: 14860 Region: IN Location: Sachs Harbour

Arctic contaminants: Exploring effective and appropriate communication between Inuvialuit communities and researchers

In November 2010, collaborative surveys were created but responses were low. Compared to interviews, which offer richer descriptions and context, it was determined that the surveys were better suited for use as pilot studies and were discontinued. Interviews were completed in June 2011, representing 27% of the Sachs Harbour population. Focus groups were not to begin until the target range (25-30%) was met. Verification of transcripts with participants is in progress. In November 2011, focus groups were going to start, but funding issues prevented this. To finish the project and deliver on set objectives, arrangements are being made for January/February 2012. Themes identified in interviews include: relationships with researchers (positive/negative/neutral impressions of interactions, frequency/duration of time in community); conceptions/perceptions of contaminants (yes/no concerns/risks, local knowledge of what contaminants are/look like, etc.); methods of communication (positive/negative impressions, recommendations). Some people found the information presented by researchers to be helpful, informative and easy to understand; however, some found it to be confusing and complicated. Most locals acknowledge the importance of learning about contaminants, especially for future generations. Meetings and presentations by researchers are viewed positively because researchers are visible in the

community, but spending time one-on-one or in small groups using storytelling and pictures are potential ways to better communicate the importance of contaminants on a local level.

Sabin, Jerald

University of Toronto Toronto, ON jerald.sabin@utoronto.ca

File No: 12 410 855 **Licence No:** 14972 **Region:** NS **Location:** Yellowknife

Alternatives North: A history

This research produced a history of Alternatives North, a social justice coalition based in Yellowknife, Northwest Territories. The research produced a book chapter on the subject. The chapter argues that the success of Alternatives North is the result of its ability to capitalize on political opportunities in the Northwest Territories and its unique organizational structure that stresses collaboration and flexibility. Relying on a consensus-based model of decision-making, the coalition has focused its efforts around three policy areas: social and economic justice; environmental sustainability; and peace. Knitting together its activities is a well-established normative framework that enables the diverse membership of Alternatives North to choose projects and allocate its time and resources. The chapter outlines the origins of Alternatives North in the Canadian nationalist and anti-free trade movements of the late 1980s and early 1990s, and the evolution of the organization's structure and operation in response to political and economic upheavals in the Northwest Territories over the subsequent two decades. It concludes with a longer discussion of Alternatives North's significance for northerners, progressives, and students of political science.

Saxon, Leslie

University of Victoria Victoria, BC saxon@uvic.ca

File No: 12 410 210 Licence No: 14922 Region: NS Location: Behchokò

Tłycho on-line and print dictionary

This research has resulted in updating of the Tłįcho Yatiì Multimedia Dictionary (online at http://tlicho.ling.uvic.ca) and the development of the iPhone/iPad/iPod application called "Yati", released in May 2012 (download it at http://itunes.apple.com/ca/app/yati/id525154015?mt=8). Researchers of the Tłįcho Community Services Agency and the Department of Linguistics, University of Victoria, worked together in increasing the number of words in these dictionary formats and recording sounds for these "talking dictionaries". The researchers produced a set of instructions on how to use the dictionary. A dictionary is an on-going project and the goal is to continue dictionary work into the future.

Schurr, Theodore

University of Pennsylvania Philadelphia, PA United States tgschurr@sas.upenn.edu

File No: 12 410 845 Licence No: 14861

Region: SA, NS Location: North Slave; Sahtú

The Genographic Project: Anthropological genetic analyses of indigenous human populations of North America - North Slave and Sahtú Dene field research

No field work was conducted under this licence in 2011. Instead, the researchers focused on analysis using the samples collected in 2010. The results show that the vast majority of individuals from the Tłycho communities have maternal genetic lineages belonging to indigenous origins. While sharing a few maternal genetic origins with the Gwich'in, the Tłychołychon, the Tave maternal geFrom a paternal standpoint, some 65% of Tłįcho male individuals belonged to indigenous genetic origins, with the rest representing non-native lineages. Like the Gwich'in, the Tłycho had genetic markers that also appear in Athapaskan (Dene) speaking populations in Alaska. The researchers also found one paternal lineage that was unique to the Tłycho. In addition, the population history of Athapaskan (Dene) speakers appears to be rather complex, with the Tłycho being distinctive from other Athapaskan groups, including the Gwich'in. This highresolution analysis further makes clear that paternal genetic material diversity among the first Native Americans is greater than previously recognized. In addition to these questions, the research data provides new details about the geographic distribution history of Athapaskan (Dene) peoples, including their origins and pattern of dispersal across the circumarctic region. The data also allows the researchers to test theories about Dene prehistory based on linguistic evidence, including supposed links to populations in Siberia. Furthermore, they will be able to expand their understanding of the recent history of aboriginal communities from the Mackenzie River valley and Great Slave Lake region. The researchers are in the process of writing and publishing papers describing their findings, which will be co-authored with members of the aboriginal communities and other Northwest Territories' researchers involved in this research.

Schurr, Theodore

University of Pennsylvania Philadelphia, PA United States tgschurr@sas.upenn.edu

File No: 12 410 845 **Licence No:** 14862

Region: IN, GW **Location:** Inuvialuit Settlement Region; Gwich'in

Settlement Region

The Genographic Project: Anthropological genetic analyses of indigenous human populations of North America

No field work was conducted under this licence in 2011. Instead, the researchers focused on analysis using the samples collected in 2009-2010. The results show that the vast majority of individuals (>94%) from the Gwich'in and Inuvialuit communities have maternal genetic lineages belonging to indigenous origins. While sharing a few maternal genetic origins, the Inuvialuit were largely distinctive from the Gwich'in. Most Inuvialuit belonged to one maternal ancestral origin, and within this group there were two sub-populations: Mackenzie River delta and those with genealogical ties to Alaska, and eastern part of the region and those with ties to Nunavut. This genetic difference and geographic gradient suggests that two distinct prehistoric migrations may have contributed to the gene pool of the contemporary Inuvialuit. From a paternal standpoint, some 65% of male individuals belonged to indigenous genetic origins; the rest represents non-native lineages. Only the Gwich'in showed genetic origins that appear in Athapaskan (Dene) speaking populations of Alaska and Canada. The researchers also found one paternal lineage that is unique to the Inuvialuit, and which is present in all the Inuvialuit, Inupiat and Yupik populations, and may also be in the Aleut populations. This suggests that it represents a founding

male lineage for all of these circumarctic populations. The paternal genetic data further suggested that Canadian Inuit and Dene-speaking populations are genetically distinct from one another, and that the formation of these groups was the result of two population expansions that occurred after the initial movement of people into the Americas. In addition to these questions, the research data provides new details about the geographic distribution history of Athapaskan (Dene) peoples. The data also allows the researchers to test theories about Dene prehistory based on linguistic evidence, including supposed links to populations in Siberia. Furthermore, they will be able to expand their understanding of the recent history of aboriginal communities from the Mackenzie River valley. The researchers are in the process of writing and publishing papers describing their findings, which will be co-authored with members of the aboriginal communities and other Northwest Territories' researchers involved in this research.

Schurr. Theodore

University of Pennsylvania Philadelphia, PA United States tgschurr@sas.upenn.edu

File No: 12 410 845 Licence No: 14863

Region: DC, SS **Location:** South Slave; Dehcho

The genographic project: Anthropological genetic analyses of indigenous human populations of North America - South Slave and Dehcho

No research was conducted under this licence in 2011.

Simmons, Deborah

University of Manitoba Yellowknife, NT simmons@cc.umanitoba.ca

File No: 12 410 678 Licence No: 14909 Region: SA Location: Dél_inę

Mapping, language and stories in Déline

The mapping, language and stories program is an outgrowth of research initiated in 2006. Funding was obtained to conduct research about land-based language, stories and spatial knowledge, focusing on the five themes of climate change, abandoned mines, health, the social economy, and governance, under the working title Learning About Changes. During the summer of 2011, doctoral research was conducted by Ingeborg Fink (Endangered Languages Documentation) and Sarah Gordon (Health, Healing, and the Stories of the Sahtúot'ıne). This phase was also a period of analysis and synthesis of previous research, leading to collaborative development of a framework for future research under the working title Stories and Songs as Policy. Co-authored presentations were made at the results conference of the Social Economy Network of Northern Canada (SERNNoCa); Strategies for Moving Forward, the 2nd International Conference on Language Documentation and Conservation (ICLDC); the Pan-Arctic Results Workshop of the Climate Change and Health Adaptation Program for Northern First Nations and Inuit Communities; and People in Places: Engaging Together in Integrated Resource Management. Contributions were made to two forthcoming books, a special issue of Rangifer journal, Sustaining Caribou and their Landscapes - Knowledge to Action, as well as a Health Canada Success Stories booklet and website.

Southcott, Chris

Lakehead University Thunder Bay, csouthco@lakeheadu.ca

File No: 12 410 800 **Licence No:** 14931

Region: IN, GW, SA, DC, NS, SS Location: All communities

Mapping the social economy in northern Canada - Northwest Territories project

No research was conducted under this licence in 2011.

Stirling, Mark

Royal Roads University Calgary, AB mark stirling514@hotmail.com

Taiga Adventure Camp feasibility study

The goal of the project was to examine the possibility of offering a new service in the Northwest Territories (NT). To accomplish this, the market, stakeholders, company capability, and financial considerations were studied. A survey was given to men that live in the NT. The survey was done in paper format and consisted of a series of closed and open-ended questions. The survey was used to determine if the people responding would like the suggested service, if they have had any previous experience with a similar service and if they would be interested in using that service again. The results of the survey helped determine the demand for the proposed service. A report was submitted to the client as part of an overall project to help the client figure out if it is worth offering the new service.

Stuhl, Andrew

University of Wisconsin–Madison Inuvik, NT andrew.stuhl@gmail.com

File No: 12 410 875 **Licence No:** 14868

Region: IN, GW Location: Inuvik; Yellowknife; Aklavik; Tuktoyaktuk

Environment, commerce, and science in western Arctic history

Research was conducted while living in Inuvik between January 1, 2011 and June 7, 2011. Information was collected in the following ways: speaking with Inuvik residents, visiting certain places in the Beaufort-Delta, and spending considerable time in the town's libraries, especially the Inuvialuit Cultural Resource Centre and the Inuvik Centennial Library. Archives and other libraries that hold records pertaining to northern history still need to be visited. Consequently, final research results were not available at the time of this summary. Some relevant results concerning the history of reindeer herding in the area were presented to Inuvik high school students. Additional information regarding the history of exploration in the area was shared during a five day field program in Ivvavik National Park. The central questions guiding the research include the following: (1) Since the late 1860s, how has scientific research shaped the land of the western Arctic, a region which includes the Arctic portions of the Northwest Territories, Yukon Territory, and Alaska? (2) What have been arctic science's relationships with federal governments, private

industries, and native communities? (3) How has scientific knowledge about arctic environments changed over time?

Todd, Zoe University of Aberdeen Edmonton, AB zoe.todd@gmail.com

File No: 12 410 815 **Licence No:** 14978

Region: IN Location: Paulatuk; Yellowknife; Inuvik

Lands, lakes and livelihoods: women's subsistence fishing in Paulatuk, NT

This research will investigate the ways people talk about, and practice, fishing in Paulatuk, NT. The methods will include participant observation, apprenticeship, oral history and archival research. This work will inform anthropological understandings of women's harvesting activities in northern Canada, and will generate detailed, locally informed data on people's relationships with the environment in the past and the present in the community. The researcher believes that research on women's subsistence practices can provide insight into contemporary relationships between people and the environment. In 2011, archival work was conducted at the Hudson's Bay Archives in Winnipeg. The notes that were taken from the Hudson's Bay outpost journals from Letty Harbour, between 1932 and 1934, as well as some notes from the Fort Anderson journal (1861), are currently being transcribed. The researcher is working closely with Anne Thrasher to share these notes. She plans to return to the Hudson's Bay archives in 2012 after fieldwork in Paulatuk, and will work closely with local residents to co-ordinate ways to obtain photos of the journals so that they may be shared with the community. This is in addition to the joint archival work that she will do with a community member at the Prince of Wales Northern Heritage Centre in Yellowknife in 2012. The researcher plans to travel to Paulatuk in early January in order to begin consulting with local community members about how to proceed with the remainder of the project (ie: interviews, fishing, workshops).

Vannini, Phillip

Royal Roads University Victoria, BC phillip.vannini@royalroads.ca

File No: 12 410 904 Licence No: 14987

Region: IN Location: Aklavik; Inuvik; Tuktoyaktuk

Dwelling off-grids

Geography has largely evolved as a land-based discipline. If people are serious about reversing geography's land-centric bias, they should not erect boundaries amongst water worlds or between land and water. Fieldwork was conducted during the month of February 2012 to study off-grid living in Canada's Northwest Territories. Research included participant observation and 15 interviews conducted in the Mackenzie Delta region of Canada's Northwest Territories, Inuvik in particular. Ice roads are an intricate assemblage of trails carved by human movement and by the movements of water across the seasons. The ice road meshwork is a mutating, temporary, ephemeral landscape, embedded within its environment. Indeed its formation is an event - turning water into a land-like icescape. As river and ocean waters meet cold air, ice roads form and change. As sun rays shine on ice roads, their surfaces change. As water accumulated in the shape of snow meets winds, snowdrifts begin to form, confusing distinctions between road and snow banks. It is by recognizing the transformational flows of water and by working with the uniqueness of these changing materials that inhabitants of the Mackenzie Delta region (and its

visitors) can use roads and thus access each other's communities and the rest of Canada. It is by taking part in the region's constant self-transformation that they create and follow their roads, routes, and trails, the territorialization of the ice road meshwork. Then liquefaction is an event leading to its deterritorialization and to the emergence of a delta that is no longer drivable but navigable. Whether congealed or liquefied, frozen solid, muddy, or in the midst of breaking up or thawing out, the water-land-air meshwork reveals a domain of entanglement open to different relations with humans for different access assemblages.

Wood, Mellissa

Government of the Northwest Territories Yellowknife, NT mellissa wood@gov.nt.ca

File No: 12 410 898 Licence No: 14959 Region: NS Location: Yellowknife

Collaboration: NWT sport and recreation sector

The sport and recreation sector in the Northwest Territories consists of many relationships and contractual agreements. Within the sector, many organizations are responsible for providing programming to improve the lives of Northwest Territories (NT) residents through physical activity, sport and recreation. This research project examined effective collaboration between one government division and one program delivery non-government organization within the NT sport and recreation sector. Through a series of one on one interviews and a facilitated focus group, it was determined that effective collaboration is based on an evolution of relationship. Resulting recommendations provided to the government division propose that value be placed on development opportunities for staff while investing in a structure which will support staff in their collaborative activities. Specifically, recommendations included: building staff interpersonal skills; learning by engaging staff in internal collaborative projects and then shifting to a sector-wide focus; and declaring and aligning personal, divisional and organizational values to create and foster a shared vision.

TRADITIONAL KNOWLEDGE 2011

Borowitz, Michelle

University of Alberta Edmonton, AB borowitz@ualberta.ca

File No: 12 410 873 Licence No: 14864 Region: SS Location: Fort Resolution

Human dimension of river resource development and transboundary water security in the Peace-Slave River Basin

In August 2011, the researcher returned to Fort Resolution to continue her fieldwork. During this time, she followed up with the research participants she interviewed in 2010 and was able to participate in both Cultural Week and Deninoo Days. She was not able to meet with each person that she had previously interviewed, so she will be returning to Fort Resolution in 2012 in order to follow-up with the research participants she missed in August 2011. Her research is progressing well and she is looking forward to returning to Fort Resolution, to not only complete her project but to re-connect with the community members who welcomed her warmly and gave her their encouragement.

Chavarie, Louise

University of Alberta Edmonton, AB chavarie@ualberta.ca

File No: 12 410 894 **Licence No:** 14927

Region: SA Location: Déline; Great Bear Lake

The biology and ecology of sympatric polymorphic lake trout, *Salvelinus namaycush*, in Great Bear Lake, Northwest Territories

Stock assessment monitoring research has been conducted on the lake trout in Great Bear Lake since 2000. In order to investigate the occurrence of four forms of shallow-water lake trout, however, many questions remained regarding lake trout ecology. With the collaboration of the Déline Renewable Resources Council, a focus group was held with six participants from the community, followed by individual interviews. Both the focus group and individual interviews involved a mixed method of semi-directed and structured exchanges on lake trout distribution, movement, habitat, diet and morphology. A slide presentation about the scientific research on this subject was offered to participants at the beginning of the traditional knowledge study.

Documentation tools, such as Google Earth, photos, audio recordings, projectors and transcription, were used to facilitate the study. A post-study validation process asked the people from the community to verify all material produced to date (i.e. the poster, presentation and report).

Davison, Tracy

Environment and Natural Resources Inuvik, NT tracy_davison@gov.nt.ca

File No: 12 410 899 Licence No: 14967 Region: IN Location: Ulukhaktok

Peary caribou and Dolphin Union caribou traditional knowledge in the ISR

Interviews were conducted in Ulukhaktok in September 2011 and January 2012. A total of 11 traditional knowledge holders were identified by the Olokhaktomiut Hunters and Trappers Committee and interviewed. A total of 8 interviews were done in September; a community assistant was hired to assist with the interviews, and a translator was hired for interviewees who wanted to be interviewed in their traditional language. A total of 3 interviews were done in January; a community assistant/translator was hired to assist with these interviews. All interviews were audio recorded and later transcribed. There was also a mapping component to the interviews. Maps were scanned and digitized after the interviews. There are currently no results available, as the information needs to be complied and then verified by the interviewees.

Lantz, Trevor

University of Victoria Victoria, BC tlantz@uvic.ca

File No: 12 410 906 **Licence No:** 14992

Region: IN,GW Location: Husky Lakes; Hendrickson Island; Peel

Plateau/Dempster Highway; Aklavik; Inuvik; Tuktoyaktuk;

Tsiigehtchic; Fort McPherson

Using Inuvialuit and Gwich'in observations to monitor environmental change in the Mackenzie Delta region

No research was conducted under this licence in 2011.

Maraj, Ramona

Environment Yukon Whitehorse, YT ramona.maraj@gov.yk.ca

File No: 12 410 865 Licence No: 14867

Region: IN Location: Aklavik; Inuvik; Ulukhaktok; Sachs Harbour;

Tuktoyaktuk; Paulatuk

Polar bear traditional knowledge for the Beaufort Sea

No research was conducted under this licence in 2011.

Nesbitt, Lorien

Lorien Environmental Consulting Vancouver, BC lorien.nesbitt@gmail.com

File No: 12 410 821 Licence No: 14897 Region: SA Location: Dél₁ne

Planning for climate change impacts on the aquatic ecosystems of Great Bear Lake and its watershed

The Déline Renewable Resources Council, with the support of the Déline Land Corporation and Indian and Northern Affairs Canada, completed a project in 2011 to investigate the impacts of climate change and commercial development on Great Bear Lake using a combination of scientific and traditional knowledge. They conducted a review of the relevant scientific literature, completed a climate change model for Great Bear Lake, and discussed traditional ecological knowledge of Great Bear Lake and climate change impacts during a series of workshops with Déline elders and hunters. The information collected was integrated in their evaluation of the vulnerability of Great Bear Lake to climate change impacts and commercial development, and informed the design of a community-based water monitoring program. The vulnerability assessment and other project materials are available on the project website at www.greatbearlake.org.

Parlee, Brenda

University of Alberta Edmonton, AB brenda.parlee@ualberta.ca

File No: 12 410 522 Licence No: 14984

Region: SA **Location:** Fort Good Hope

Community perspectives on the health of caribou, moose, and deer populations around Fort Good Hope

No research was conducted under this licence in 2011.

Sandlos, John

Memorial University of Newfoundland St. John's, NL isandlos@mun.ca

File No: 12 410 847 **Licence No**: 14866

Region: SA, NS, SS **Location:** Former Pine Point mine/townsite; Fort Resolution; Hay River; Giant and Con mine sites;

Yellowknife; Dettah; Dél_lne; Port Radium mine/townsite

Abandoned mines in northern Canada: historical consequences and mitigation of current impacts

In May 2011, the research team conducted extensive archival research on the history of Giant Mine, as well as the aboriginal employment policy in the mineral industry. Some members of the research team also worked in partnership with the Goyatiko Language Society to begin oral history interviews in Dettah and Ndilo about the historical impacts of Giant Mine. To date, Goyatiko researchers have conducted, transcribed, and translated eleven oral history interviews. A workshop will be held in Dettah in November 2011 to discuss the results of this research. In the

Pine Point area, workshops were conducted in Fort Resolution and Katlodeeche First Nation to communicate research results. During these workshops, options for community use of the research results were discussed with attendees. Options included using the results as an educational resource and as part of a community history, amongst others. A paper on the history of Pine Point has been accepted for publication in the journal "Environment and History", and will appear in early 2012. In Déline, a graduate student spent the summer months conducting fieldwork with the goal of developing community contacts and partnerships for her research on how local people interpret artistic and media representations of uranium mining in the Sahtú region. The abandoned mines project has developed a new website, with a blog documenting activities. You can find the comments. their blog. and post http://www.abandonedminesnc.com/.

Simmons, Deborah

University of Manitoba Yellowknife, NT simmons@cc.umanitoba.ca

File No: 12 410 678 Licence No: 14904 Region: SA Location: Dél_inę

Caribou and communities in the Sahtú region

This program was initiated in response to announcements of declining barren-ground caribou herds. Activities took place in each of the five Sahtú communities in 2007-2010, along with two regional harvester workshops and an international conference (the North American Caribou Workshop, or NACW). During 2011, core activities related to this program included the following: the development of a proposal for a Sahtú regional validation workshop; a knowledge exchange including youth, harvesters and leaders through the Indigenous Talking Circle at the international Arctic Ungulate Conference; and preparation, editing and publication of the NACW proceedings in special issue #20 of the journal "Rangifer". This journal submission was titled "Sustaining Caribou and their Landscapes — Knowledge to Action", and included submissions from aboriginal authors speaking to key issues in caribou research and management.

Snortland, Jody

Wek'èezhìi Renewable Resources Board Yellowknife, NT jsnortland@wrrb.ca

File No: 12 410 636 Licence No: 14965
Region: NS Location: Slemon Lake

Ihda k'ètì aquatic ecosystem monitoring project

The main goals of this project were to share and document Tłįchǫ knowledge and western scientific knowledge on the aquatic environment in Russell Lake. The project engaged local community members in sampling and recording a standard set of observations, using both Tłįchǫ and western scientific knowledge. It involved community members in a meaningful manner in all aspects of conducting contaminants related research, including the actual pursuit of monitoring and research objectives. A monitoring camp was held on Russell Lake, a location that supports a strong aboriginal subsistence fishery. Water, sediment and fish were sampled by elders, youth and fisheries scientists. Elders provided assessments of fish health, and described the indicators they use to identify fish health. Scientists sampled fish tissues and demonstrated to elders and youth the methods for collecting fish tissues for analysis. A results workshop was held in Behchokö to present the results of the fish tissue analysis, water and sediment quality sampling.

Community members were informed and educated on the status of contaminants in the fish they may be eating and that these foods remain healthy choices, perhaps within certain limits. Annual implementation of the program and consistent use of the monitoring protocols developed this year will be the key to achieving the main goals of long-term monitoring: detecting change over time and space.

Svoboda, Michael

Arctic Borderlands Ecological Knowledge Coop Whitehorse, YT michael.svoboda@ec.gc.ca

File No: 12 410 811 **Licence No:** 14989

Region: IN, GW **Location:** Fort McPherson; Tsiigehtchic; Aklavik; Inuvik;

Tuktoyaktuk

Arctic Borderlands Ecological Knowledge Coop: Community based ecological monitoring program

The Coop uses both local and scientific knowledge to monitor and assess environmental changes in the range of the Porcupine caribou herd, and nearby coastal and marine areas. Interviews with local experts are conducted every year by community researchers. People share what they see and hear about fish, berries, caribou, unusual animal sightings, weather conditions, and other things while they are out on the land. This year was the first year that the researchers used the reviewed and updated community questionnaire. Community interviews were completed for Tsiigehtchic, Fort McPherson, Aklavik, Inuvik, Old Crow, and Arctic Village. Also, community reports were completed and mailed to program partners and community participants at the end of 2011. The Coop website is an important communication tool, and contains proceedings from past meetings as well as the results from the community monitoring program. Lastly, two videos were published on their website main page. The videos, along with all other documents, can be viewed at www.taiga.net/coop.

ARCHAEOLOGY 2011

Andrews, Tom

Prince of Wales Northern Heritage Centre (GNWT)

Permit No: 2011-004 **Class:** 2

Region: SA Location: Tulít'a District, Sahtú Settlement Area

NWT ice patch monitoring project Tłįchǫ

Bad weather—rain, fog, and low clouds—plagued the researchers' fieldwork this year, significantly restricting their ability to reach the ice patches by helicopter. Though they were camped in the mountains from August 15th to 18th, the weather permitted only a few hours of flying each day. Usually, this was later in the afternoon. This year, their Tulít'a partner was unable to join them due to other commitments, but his seat in the helicopter was filled by a PhD student from the University of Alberta. The student will focus his PhD thesis on the broader cultural and ecological context of ice patch use in the Northwest Territories. As with the 2010 field season, the researchers were shocked at the amount of melting at several sites. It seems that as the ice patches melt, they reach a critical tipping point where enough dung is exposed to dramatically change the albedo of the patch, leading to rapid melting. They have seen this in recent years, most dramatically at the KhTe-2 site. Despite poor weather, they discovered a new archaeological ice patch site, KhTf-3, at an elevation of just over 2000 metres. They recovered approximately ¾ of the proximal end of a wooden arrow from this site. The nock end was recovered, the distal end with the projectile point was not.

Bussey, Jean

Tibbitt to Contwoyto Winter Road Joint Venture

Region: NS Location: North Slave, Tłįcho Settlement Area

Tibbitt to Contwoyto winter road project

In 2011, an archaeological investigation was conducted for the Joint Venture that operates the Tibbitt to Contwoyto winter road. This road runs from the south end of Tibbitt Lake near Yellowknife to almost the north end of Contwoyto Lake in Nunavut. This ice road was used every winter for over 25 years, but since the winter of 2008 to 2009 it has not been constructed north of Lac de Gras due a lack of mining activity. In previous years, a number of archaeological sites located near the winter road or its associated developments were marked by stakes to ensure avoidance during winter activities. Monitoring of the protected archaeological sites was a major component of the 2011 investigation. In addition, ground reconnaissance was done at two potential gravel sources located near Lockhart Lake camp. In total, there are seven sites that are protected from accidental impact by the installation of markers. Whenever possible, these markers are 30 m from archaeological sites, but in most instances this is not possible because

the development occurred prior to archaeological investigations. Damaged stakes were replaced when necessary and the top of all markers were sprayed with fluorescent paint to make them more visible in winter. In the process of visiting the protected archaeological sites, other portages were examined from the air to confirm their status and ensure no new disturbances have occurred in areas with archaeological potential. During the ground reconnaissance, an archaeological site with multiple localities was discovered at each of the proposed gravel source near Lockhart Lake. At the preferred source, there were four localities with primarily sparse archaeological material. A few flakes of non-quartz material were collected and the small sample of quartz flakes present was left in situ. There is little potential for significant archaeological material at this site. At the other proposed gravel source, three localities with intact archaeological material were located and all specimens were left in situ. The second gravel source has greater potential to yield intact and significant archaeological material. If this potential source is selected, more extensive testing, and possibly excavation, will be required in addition to more ground reconnaissance.

Cary, Henry Parks Canada

Region: SA Location: Sahtú Settlement Area, Délıne District

Parks Canada investigations at Cloud Bay, Great Bear Lake

On 11 August 2011, two people travelled from Inuvik to Saoyú-?ehdacho National Historic Site to assess the location for a new cabin in Cloud Bay, a small inlet on the Saoyú Peninsula in Keith Arm, Great Bear Lake. The 9.8 × 6.1 m traditionally-built cabin will be used as a teaching and healing camp for elders and youth, and accessed by boat or float plane from Déline. Parks Canada had conducted an assessment for this project in 2010 on an adjacent beach, but this location was rejected in favour of building the cabin further north. Archaeological sites have been found near the study area, most notably an early occupation found on Dog Point in 1951, so further assessment was necessary to ensure that remains would not be disturbed during the cabin's construction. After landing on the beach in Cloud Bay and a brief search of the surrounding area, an elder from Déline selected a cobble bench approximately 160 m inland as the building site. A surface survey was conducted and the depth of the cobble was assessed. After digging below 30 cm, no archaeological deposits or artefacts were found, and the excavation was abandoned. The proposed foundation area, a section from the beach to the study area, and the high water mark nearest the proposed cabin location were mapped using GPS. On the return trip to Déline a series of oblique aerial views of the Déline Fisheries and Fort Franklin National Historic Site Designated Place were taken The Cloud Bay cabin will be constructed in Spring 2012, and will not require further archaeological assessment.

Cary, Henry Parks Canada

Region: GW Location: Gwich'in Settlement Area

Hannah's Field archaeological assessment

In 2011, the Teetl'it Gwich'in Council wished to move two large log cabins from the Tl'oondih Healing Camp to 'Hannah's Field,' a partially vacant lot due east of the Fort McPherson Anglican Church and cemetery. There was some concern that this new construction would impact archaeological remains at the site. Although unexplored, the property was thought to have a number of features given its proximity to Fort McPherson National Historic Site, and because several finds had been made while excavating a children's playground within a hundred metres

of Hannah's Field. Two Parks Canada employees travelled to Fort McPherson on 6 October 2011 to visit the proposed construction area and determine whether archaeological excavation was necessary before the cabins were moved to the site. Despite the light snow cover, the researchers could readily see the foundations of structures built on the proposed site in the past 100 years. They determined that the option with least impact to the archaeological remains was an on-grade pillar structure. This would involve laying a series of gravel beds on grade, which would provide a base for the horizontal wood pillars supporting the cabin structure at its corner and mid-points. They also recommended that the gravel beds be separated from the ground surface using geotextile. Since this option does not require excavation, the researchers did not recommend that the construction be monitored by an archaeologist. They also assessed the Historic Sites and Monuments Board plaque at Fort McPherson National Historic Site. They found the two posts holding the plague to be rotted, and one had snapped. Additionally, the plague's location behind the Anglican Church is not easily found by visitors approaching from the road. Since the plaque will be replaced, the community has requested that it be moved to a new location on the roadside of the church. The researchers agree with this proposal, but suggest that any new plaque location be tested by archaeological excavation in advance.

Clarke, Grant

Aurora Geosciences Ltd. and TNR Gold Corp.

Region: NS **Location:** Akaitcho Territory, North Slave Region

Moose property archaeological investigation

A Heritage Resources Impact Assessment was completed at Moose property, located on the north shore of Great Slave Lake, in advance of mineral exploration. The purpose of the investigation was to identify, record, and evaluate previously unrecorded archaeological sites and to provide recommendations to mitigate potential impacts if development sites were identified within the study area. These investigations focused on the proposed drilling locations and were carried out over two days. The first day of fieldwork was completed in August 2011. The second day of fieldwork took place in September and provided an opportunity for a site visit with elders and youth representing the Łutsel K'e First Nation, the Yellowknives Dené First Nation, the Deninu Kue First Nation and the Fort Resolution Métis Council. The visit allowed the representatives an opportunity to visit the site and provide advice regarding the nature and significance of the area. and to comment on the potential of the area for archaeological resources. No previously recorded sites were identified within the study area during the researchers' pre-field studies. Field investigations focused on areas exhibiting moderate to high potential for archaeological materials that may be impacted by the proposed development. Field methods consisted of pedestrian transects with the intent to identify and assess any archaeological sites that may be present. This included the excavation of limited shovel tests to investigate for buried archaeological materials. Much of the area investigated, however, was bedrock with little to no sediment. No precontact archaeological sites were identified and it was generally believed that the area has moderate archaeological potential at best. The abandoned workings and some machinery are still evident from the abandoned De Staffany mine and previous camp area. An area that is currently used as a camp by local people and exploration crews was also noted. A total of 11 shovel tests were excavated in these two areas, but no archaeological materials were recorded.

Clarke, Grant

The Department of Transportation, GNWT

Permit No: 2011-014 Class: 2

Region: IN Location: Inuvialuit Settlement Area

Tuktoyaktuk to Inuvik Highway archaeological investigation

The proposed Highway runs north from Inuvik to Tuktovaktuk on Inuvialuit-owned landsas well as Crown Land. The proposed Highway measures 138 km in length in its current alignment, starting at the end of Navy Road in Inuvik (km 0) and ending in Tuktoyaktuk (km 138). The objectives of the Heritage Resources Impact Assessment (HRIA) were to identify, record and assess heritage resources that might be impacted by the proposed Highway project, and to devise appropriate mitigation strategies should any be found in conflict with the proposed Highway. The field investigations of the HRIA were completed in September 2011 over a six day period. The field crew consisted of two IMG Golder employees and four Inuvik community members who assisted during the field program and provided advice on the cultural significance of the landscape. Prefield research noted that there were no previously recorded sites within the proposed right-of-way routes. Five archaeological sites were previously recorded in areas that are potential borrow sites for gravel. The assessment was conducted along the planned Highway right-of-way and at several proposed borrow source locations. Aerial helicopter surveys were used to conduct preliminary reconnaissance to confirm areas with moderate to high potential for the presence of cultural materials. Areas deemed to have potential were ground truthed using pedestrian surveys and shovel tests. In total approximately 189 shovel tests were excavated in the study area, and one previously recorded site was revisited to identify any possible conflict with the proposed program. No artefacts were recovered from the test sites.

Clarke, Grant

Aurora Geosciences Ltd. and Boxxer Gold Corp.

Permit No: 2011-017 **Class:** 2

Region: NS Location: Akaitcho Territory, North Slave Region

Ven Lake property archaeological investigation

In October of 2011, Golder Associates Ltd. conducted a Heritage Resources Impact Assessment at a series of potential diamond drill locations near Ven Lake, Northwest Territories. The work was conducted on behalf of Aurora Geosciences Ltd. and Boxxer Gold Corporation. No archaeological sites have been previously recorded in the local study area, although sites are known to be present in the region. The investigations were conducted to identify, record, and evaluate the proposed location for the presence of previously unrecorded archaeological sites, and to provide recommendations to mitigate potential impacts if sites were identified. The field crew consisted of an employee of Golder Associates Ltd. (Yellowknife) and a member of the Yellowknives Dené First Nation. Procedures used for this project were standard for projects of this nature in the region, and included pre-field studies, helicopter overflights, on-ground reconnaissance, reporting and formulation of recommendations. Project planning also included provisions for a representative of the local community to accompany the field crew during the field inspection and to provide advice regarding the nature and significance of the sites in the area. Field investigations began at an existing, and currently unoccupied, camp located on the west shore of Ven Lake. An existing series of cut lines connected the areas of investigation. Evidence of previous exploration, including bulk trenches and cleared outcrops, were noted during the assessment. The fieldwork focused on areas exhibiting moderate to high potential for archaeological materials that may be impacted by the proposed development. These areas were shovel tested in an effort to identify any buried cultural materials or palaeosols. Because much of the area was exposed bedrock outcrops, limited shovel testing was conducted, but all completed tests were negative. No archaeological sites were recorded as a result of the investigation.

Harris, **Ryan** Parks Canada Region: IN Location: Inuvialuit Settlement Area

H.M.S. Investigator underwater archaeological survey

No summary was provided for this 2011 permit.

Lobb, Wayne Murray

Mackenzie Valley Highway Project

Region: SA **Location:** Sahtú Settlement Area, K'ahsho Got'ine District

Mackenzie Valley Highway project - Gibson Gap to Thunder River

An Archaeological Impact Assessment (AIA) was conducted near Fort Good Hope. Northwest Territories for the proposed K'asho Got'ine Highway (KGH). The proposed highway would convert the winter road from Norman Wells to Fort Good Hope into a year round road, which would then be extended northward toward Little Chicago. The goal of this project was to relocate and reexamine 13 known archaeological sites that are on or near the proposed KGH route. In addition, six other archaeological sites were to be examined at borrow sources if time was available. These 19 sites consisted of paleontological sites, historic structures, historic trails, and pre-historic archaeological sites. The fieldwork was conducted from July 12th to 16th. The field program was carried out by hiking, boat, and helicopter at various points during the survey. Twelve sites on the main KGH route were revisited. Four of the sites were determined to be outside of the KGH rightof-way. A new paleontological site was recorded during a survey of the Jackfish Summer Trail (MbTb-12). Unfortunately, due to helicopter issues, the rest of the archaeological sites, including one on the proposed KGH right-of-way and six in borrow source locations, could not be revisited. Approximately 80 artifacts were recovered, consisting of fossils and stone tool debitage (debitage is the by-product from stone tool manufacturing). The new artifacts are from four of the archaeological sites surveyed in 2011. In addition, one of the sites on the Jackfish Summer Trail (MbTb-11) featured the remnants of at least two hearths. It is hoped that this past year's fieldwork will lay the groundwork for any future development of the KGH and borrow sources.

MacKay, Glen

Prince of Wales Northern Heritage Centre (GNWT)

Region: DC Location: Dehcho Region

Kakisa archaeology project

The Prince of Wales Northern Heritage Centre (PWNHC) conducted a community archaeology project in partnership with the Ka'a'gee Tu First Nation in 2011. The goal was to help the community document cultural values in the Ka'a'gee Tu Candidate Area, which is being considered for protection as a National Wildlife Area through the NWT Protected Areas Strategy. Over three days in July, archaeologists from the PWNHC worked with an elder from the Ka'a'gee Tu First Nation to document important cultural places around Kakisa Lake. A previous archaeological survey was conducted around Kakisa Lake in the late 1970s. The researchers relocated several of the archaeological sites recorded during that survey in order to evaluate their conditions and determine more precise locations for these sites using GPS. The sites included two log cabin villages that were occupied in the 40s and 50s, and a lithic scatter located on the Muskeg River (which flows into the east side of Kakisa Lake). A small test excavation at the latter site indicates that it contains at least two precontact components. The researchers also recorded

two new sites: a precontact lithic scatter, and an historic fish camp. They hope to conduct more extensive surveys of Kakisa and Tathlina Lakes in future seasons.

MacKay, Glen

Prince of Wales Northern Heritage Centre (GNWT)

Region: DC Location: Dehcho Region

Dehcho highways archaeology project

The Prince of Wales Northern Heritage Centre (PWNHC) conducted a community archaeology project in partnership with the Jean Marie River First Nation in 2011. The goal of the project was to document cultural values in the Łue Túé Sųlái Candidate Cultural Conservation Area, which is being considered for protection through the NWT Protected Areas Strategy. During one week in July and another week in September, archaeologists from the PWNHC worked with elders from the Jean Marie River First Nation to survey portions of two small lakes in the Łue Túé Sųlái area: Ekali and Sanguez Lakes. The researchers recorded eight new archaeological sites, including five precontact lithic scatters, a cabin, a cache, and a trail. According to the oral traditions of the people of Jean Marie River, these small inland fish lakes were important winter harvesting areas, where fish caught through the ice and small game provided important staples for the winter months. The researchers expect the density of archaeological sites in the Łue Túé Sųlái area to be high, and are planning more extensive surveys and test excavations for 2012.

Prager, Gabriella

EBA Engineering Consultants Ltd.

Region: NS Location: Akaitcho Region, North Slave Region

Nechalacho rare earth element project

The Nechalacho Rare Earth Metals Project is located on the north side of the east arm of Great Slave Lake, approximately 95 km southeast of Yellowknife, with mine development focused around Thor Lake, about 4 km due north of Great Slave Lake. The archaeological team for this project consisted of two archaeologists and a local person from each of the three closest communities (Dettah, ŁutselK'e and Fort Resolution). Archaeological investigations were conducted in August 2011, and consisted of pedestrian surveys of the proposed mine site and associated developments. During this field project, six archaeological sites were found, all along the north sides of two small lakes associated with proposed tailings containment locations. These sites are small, comprising one or two stone circles that were probably tent rings, hearths and a rock structure identified by the researchers' local crew members as a marten trap. The small sizes of the circles suggest that they probably represent one night stops by a lone traveller. These sites provide information about past people's travel routes and regional use patterns. Their presence suggests that interior parts of the study area were used more than expected. Further detailed recording and investigations will be completed at all newly recorded sites. Three sites found during the original 1988 archaeological survey of the Thor Lake project were revisited. One is the existing exploration camp, one was a small quartz scatter and the third contained three quartz tool fragments found on different beach ridges on Great Slave Lake. Although no additional artifacts were found this year at the latter site, it is the only one that may require further work. Since it is close to the proposed dock and associated laydown areas, the site boundaries need to be defined so that impacts can be avoided if possible.

Seip, Lisa

Seabridge Gold Inc.

Region: NS Location: Tłįcho Settlement Area and Akaitcho Region

Courageous Lake

This work was conducted by two archaeologists and two community members, and was a continuation of a baseline study that was conducted in 2010. Work was focused around the southern end of Courageous Lake and the areas surrounding Matthews Lake. Community meetings were conducted with the Lustel K'e Dene, the Tłycho, the North Slave Métis Alliance, and the Yellowknife Dene between March 15th and 20th. Topics discussed included archaeological baseline studies conducted in 2010 and archaeological studies proposed for the current year. Community site visits were held between August 22nd and August 25th and included members of the North Slave Métis Alliance, the Northwest Territory Métis Nation, the Tłıcho, and the Lustel K'e Dene. The meetings and site visits allowed community members to learn about and provide feedback regarding the baseline programs to date. The fieldwork resulted in the recording of 55 new archaeological sites, including 32 lithic sites, 13 rock cairns, one rock feature site, eight multicomponent sites (containing a combination of lithics, rock cairns, other rock alignments, and/or historic features), and one site containing a modified piece of caribou antler. Additionally, one previously recorded site (LbNw- 3) containing two grave sites was revisited and found to be in good condition. Additional archaeological studies are planned for 2012. Avoidance is the preferred management recommendation for all sites, and if avoidance is not possible then systematic data recovery is recommended. As the project is currently in the design phase no impacts are anticipated in 2011.

Walker, Daniel

Tamerlane Ventures Inc.

Region: SS Location: Akaitcho Region, South Slave

Pine Point project

In 2011, archaeological investigations of the proposed Pine Point project were conducted. This project is located between Hay River and Fort Resolution, south of Great Slave Lake. Fieldwork was conducted from August 22 to September 3 under snow free conditions. The field investigations were conducted by three archaeologists and three local community members from the Deninu Kue First Nation, the Fort Resolution Métis Council, and the Hay River Métis Council. The main objective of this investigation was to identify, evaluate, and record any archaeological sites located within seven proposed deposit areas. During pedestrian surveys of the project area subsurface testing was conducted on terrain features determined to have the potential to contain cultural materials, as well as a sampling of areas determine to have low potential to contain cultural material in order to confirm that assessment. As a result of this investigation, four new prehistoric archaeological sites containing lithic material created during stone tool production were recorded.

Wickham, Michelle

Bison Historical Services Ltd.

Region: SA Location: Tulít'a District, Sahtú Settlement Area

Slater River exploration program

In September of 2011, Bison Historical Services Ltd. carried out a survey for heritage sites southeast of Norman Wells in the Sahtú Region. The goals of this investigation were to conduct a pre-impact examination of all areas that may be impacted by 2011/2012 development activities to ensure any unrecorded heritage resource locations will be avoided, and to relocate a known archaeological site (LgRs-2). Fieldwork was based out of Norman Wells and was carried out by helicopter and on foot. Investigations focused on high potential areas within the 3-D seismic area, as well as locations where construction activities will take place (e.g. winter access road, base camp, construction camp, two helipads, security shack, staging area, two well leases, and the shoreline of fourteen water sources). The 3-D seismic area was repeatedly overflown in systematic grids at low elevation and slow speed to identify any possible heritage concerns. A low elevation, slow spiraling over-flight was conducted at each location noted above. The majority of these locations were identified as having low heritage resource potential given some combination of low topographic relief, muskeg ground cover, black spruce vegetation, standing water and existing disturbances along the access road or existing seismic lines. The over-flights and photo documentation were deemed an appropriate level of assessment for these sites. Ten areas with high heritage resource potential (as identified during over-flights and pre-field map analysis) were further assessed through pedestrian and subsurface testing. All 180 shovel tests that were excavated yielded negative results. Only one known site (LgRs-2) occurs close to any of the currently proposed development activities. LqRs-2 is located approximately 1.4 km north of the access road and will not be impacted by the currently proposed construction activities. LqRs-2 was recorded in 2003 as a collapsed tripod with insulators and wire, part of the CN communications line installed around 1960. During this investigation, the tripod was relocated. The insulator and wires are still in good condition; however, the wooden poles are starting to decompose. No previously unidentified heritage sites were found within the Slater River Exploration Program, so it will not impact any known heritage sites.

WILDLIFE 2011

Abernathy, David

BHP Billiton Diamond Inc.

Yellowknife, NT

Permit No: 5693 **Species Studied:** Caribou, grizzly bear, wolves, falcons

and birds

Region: NS Location: BHP Billiton property, approximately 300km

northeast of Yellowknife

Wildlife effects monitoring program

The requirement for the wildlife effects monitoring at the Ekati Diamond Mine is described in the Environmental Agreement between Canada (DIAND), the GNWT (ENR) and BHP (January 1997).

Bidwell, Mark

Canadian Wildlife Service Saskatoon, SK

Permit No: 4821 **Species Studied:** Whooping crane **Region:** SS **Location:** South Slave Region

Whooping Crane ecology and rehabilitation

The goal of the whooping crane recovery strategy is to protect, restore and manage whooping cranes to be self-sustaining in the wild and to downlist the species from endangered to threatened.

Carriere, Suzanne

ENR Wildlife Yellowknife, NT

Permit No: 5764 **Species Studied:** Mice, voles, lemmings, shrews

Region: IN, GW, NS, SS, DC, SA Location: All regions in the NWT

NWT small mammel and hare transect survey

Establish the abilty to predict small mammal cycles throughout the Northwest Territories.

Cluff, Dean

ENR North Slave Yellowknife, NT

Permit No: 6883 **Species Studied:** Black bears **Region:** NS **Location:** North Slave Region

Black bear movement and ecology in the North Slave Region

No studies of black bears in the North Slave Region have been done, therefore very little information is known about these bears in the northernmost part of their range.

Cluff, Dean ENR North Slave Yellowknife, NT

Permit No: 5690 **Species Studied:** Tundra wolves

Region: NS Location: Central tundra region of the NT

Index abundance for tundra-denning wolves

Management plans prepared for barren-ground caribou herds recommend monitoring for trends in predator abundance.

Coulton, Dan

Golder Associates Limited

Yellowknife, NT

Permit No: 6877 **Species Studied:** Raptors and water birds

Region: NS **Location:** 15 km radius from Nico Project base camp

Baseline wildlife studies for Fortune Minerals NICO Project at Lou Lake study area

Wildlife surveys were completed, to augment existing information on wildlife species, the habitat surrounding the proposed mine site, and all-weather access road route.

Croft, Bruno

ENR North Slave Yellowknife, NT

Permit No: 6878 Species Studied: Caribou Region: NS Location: North Slave Region

Continue monitoring the health, condition and contaminants of the Bathurst and Bluenose East caribou in the North Slave Region

Health, body condition, disease, and parasites of barren-ground caribou provide important information on the status of the herds and on the potential for population growth.

Croft. Bruno

ENR North Slave Yellowknife, NT

Permit No: 6879 **Species Studied:** Caribou

Region: NS, SS Location: Behchokò; Whatì; Gamètì; Wekweètì; Dettah;

and Łutsel K'e

Continue monitoring the Bathurst and Bluenose East caribou herds

In order to insure that barren-ground caribou and caribou herds remain healthy, a number of monitioring actions must be undertaken on an annual basis, to provide decision makers with the information required to address management objectives.

Davidson, Tracy ENR - Inuvik region

Inuvik, NT

Permit No: 7410 **Species Studied:** Barren-ground caribou

Region: IN Location: Range of the Tuktoyaktuk Peninsula; Cape

Bathurst; Bluenose-West barren ground caribou herds

Late winter recruitment, and fall composition surveys of the Tuktoyaktuk Peninsula, Cape Bathurst, and Bluenose-West barren-ground caribou herds

Late winter recruitment survey, fall composition, and collar monitoring.

Davidson, Tracy

Gwich'in Renewable Resource Board

Inuvik, NT

Permit No: 7420 Species Studied: Moose

Region: GW Location: Various locations throughout the Richardson

Mountains, Mackenzie Delta, Eagle Plains

Moose abundance and composition survey

This survey will provide moose population density, composition and recruitment rates and will also give important insights into the influence of habitat characteristics and human-related disturbances on moose.

Decker, Robert

ENR South Slave Hay River, NT

Permit No: 5404 Species Studied: no specific wildlife species Region: IN Location: Sachs Harbour; Ulukhaktok

Ecological regions (ecosystem classification) of the Northwest Territories arctic islands

All provinces and territories have some form of hierarchical ecological land classification that integrates climate, physiography, and biotic factors in some fashion. These classifications can be a useful tool / framework for ecosystem based resource management and land use planning, as well as for understanding the effects of climate change, cumulative effects and natural disturbances on the landscape.

Derocher, Andrew

University of Alberta

Edmonton, AB

Permit No: 7425 **Species Studied:** Female polar bears

Region: IN, GW Location: Inuvik Region

Movement and habitat use by adult female polar bears

Adult female polar bears with cubs will be caught in the southern Beaufort Sea (From Herschel Island to Baillie Islands, NT) and instrumented with GPS satellite transmitters. The study will monitor the movements and habitat use of adult females.

Elkin, Brett ENR Wildlife

Yellowknife, NT

Permit No: 5761 Species Studied: All wildlife species

Region: IN, GW, NS, SS, DC, SA Location: NWT wide

Wildlife health, condition, and genetic monitoring

Although most wild animals are healthy, diseases and parasites can occassionally occur in any wildlife population. Some diseases and parasites are naturally occurring and appear to cause little problem in their host species, while others have the potential to impact wildlife, at both the individual animal and population level.

Enbridge Pipelines

Enbridge Pipelines (NW) Inc.

Norman Wells,

Permit No: 5760 **Species Studied:** Various wildlife species

Region: DC, SA **Location:** Selected areas of the Enbridge Pipeline, right -of

- way in the vicinity of various Dehcho and Sahtú

communities

Monitoring wildlife along Enbridge ROW

A community based wildlife monitoring program was established when pipeline operations began in the mid-1980s and ran for several years before being terminated.

English, Colleen

Rio Tinto, Diabik Diamond Mines Inc.

Yellowknife, NT

waterfowl and other aquatic birds, raptors

Region: NS **Location:** Diavik wildlife study area

2010 wildlife monitoring program for the Diavik diamond mine

To monitor wildlife within the vicinity of the mine.

Fast. Marie

Canadian Wildlife Service

Yellowknife, NT

Marsh bird surveys in Kwets'ootl'àà candidate protected area

Kwets'ootl'àà is a candidate National Wildlife Area being proposed through the NWT Protected Areas Strategy (Mackenzie Valley 5-Year Action Plan). As part of Step 5 of the NWT Protected Areas Strategy, an ecological assessment must be conducted, to identify the key ecological components in the candidate area, before a final decision can be made to proceed with legally designating the site.

Fronczak, David

United States Fish and Wildlife Service Division of Migratory Bird Management Bloomington, MN United States

Permit No: 4824 Species Studied: Ducks Region: NS Location: Mills Lake Marsh

Western Canada cooperative preseason waterfowl banding prgram - Mills Lake Station, NT Preseason banding of 2,000 mallards, 1,500 northern pintails, and 1,000 of all other waterfowl species.

Green, David

CWE

Burnaby, BC

Permit No: 5401 **Species Studied:** Yellow warblers

Region: IN, GW Location: Inuvik Region

Latitudinal variation in life history traits and carry-over effects of Yellow Warblers

Species with wide breeding distributions, such as Yellow Warblers, provide an excellent opportunity to study potential trade-offs among vital rates at different latitudes.

Groves, Debbie

US Fish and Wildlife Service Migratory Bird Management Juneau, Alaska United States

Permit No: 5403 **Species Studied:** Geese, swans, ducks, loons, gulls, terns

and owls

Region: IN Location: Banks Island; Tuktoyaktuk Peninsula; Western

Victoria Island

Aerial waterfowl surveys on Banks Island, Tuktoyaktuk Peninsula, and Western Victoria

Information on waterfowl abundance, distribution, and population trends in the Northwest Territories is needed to ensure that populations are conserved for the long-term use, as well as that appreciation by northern residents and all people residing within the species' migratory range is maintained.

Harpley, David

Canadian Zinc Vancouver, BC

Permit No: 5033 Species Studied: Woodland caribou

Region: DC **Location:** Prairie Creek Mine

Aeiral surveys of the Prairie Creek mine access road for caribou

Canadian Zinc's's access road links the Prairie Creek mine to the Liard Highway, near Nahanni Butte, and crosses terrain that is not belived to be home range for significant caribou populations.

Hegel, Troy

Environment Yukon (V5-A) Whitehorse, YT

Permit No: 5765 **Species Studied:** Nahanni and Coal River caribou herds

Region: DC, SA Location: Sahtú; Dehcho

Population monitoring of the Nahanni and Coal River caribou herds

Recent fall composition surveys, increasing hunter traffic on the Nahanni Range Road, increasing industrial development, and reports of low numbers of large bulls are all contributing factors heightening the level of concern regarding the South Nahanni and Coal River caribou herds.

Kardynal, Kevin

Canadian Wildlife Service

Yellowknife, NT

Permit No: 4822 **Species Studied:** Canada warbler, olive-sided flycatcher,

rusty blackbird

Region: DC **Location:** Protected area in the Dehcho region,

surrounding Kakisa

Nest searching in Ka'a'gee Tu candidate protected area

As part of Step 5 of the NWT Protected Areas Strategy, an ecological assessment must be conducted to identify the key ecological components in the candidate area before a final decision can be made to proceed with legally designating the site.

Kelly, Allicia

ENR South Slave Fort Smith, NT

Permit No: 4826 **Species Studied:** Moose population **Region:** SS **Location:** Slave River Lowlands

Moose population survey - Slave River Lowlands

Moose surveys were completed when moose are found in more open habitats and snow cover is complete. Survey to collect data on age and sex composition, as well as population abundance.

Kelly, Allicia

ENR South Slave Fort Smith, NT

Permit No: 4813 Species Studied: Moose Region: SS Location: South Slave Region

Moose - Baseline health survey

In the South Slave region, moose are one of the most important food animals to the communities. Understanding what types and levels of parasites and disease moose currently carry and the occurrence of new or emerging pathogens is an important first step in determining population health.

Kelly, Allicia ENR South Slave Fort Smith, NT

Permit No: 5757 **Species Studied:** Barren-ground caribou

Region: SS **Location:** ŁutselK'e; Fort Resolution; Fort Smith

Barren-ground caribou monitoring: population parameters, movements, distribution, health, and habitat use of the Beverly and Ahiak herds

Monitoring of the Beverly and Ahiak caribou herds is imporant to determine the status of these herds and to understand factors driving changes in herd status. This work is a part of strategy 5 of the barren-ground caribou management strategy for the northwest territories 2006-2010.

Kutz. Susan

University of Calgary

Calgary, AB

Permit No: 5158Species Studied: CaribouRegion: SALocation: Sahtú Region

Community-based monitoring of wildlife health

Climate and other landscape changes may impact the health of wild animals. This research establishes ongoing wildlife health monitoring programs, in collaboration with local hunters and trappers.

Larter, Nic

ENR Dehcho Fort Simpson, NT

Permit No: 5036 Species Studied: Moose

Region: NS Location: Dehcho Region, includes areas north of and

along the Mackenzie River

Geospatial moose survey in the Dehcho

Concerns have been voiced over depressed moose abundance, along high use areas in the Dehcho Region. There was unanimous agreement by all first nation representatives at the Regional Wildlife Workshop, in Fort Simpson, in September 2002, that there was a need for moose surveys, to establish some baseline information on moose densities in the region.

Larter, Nic ENR Dehcho Fort Simpson, NT

Permit No: 5032 Species Studied: Wood bison Region: DC Location: Fort Liard; Nahanni Butte

Nahanni wood bison population survey

The first and only survey to estimate population number of the Nahanni wood bison population was conducted in March 2004. At the time the population was estimated at about 400 adults and it was acknowledged that future population surveys would be required every 5-6 years.

Larter, Nic ENR Dehcho Fort Simpson, NT

Permit No: 5031 Species Studied: Boreal caribou

Region: DC Location: Dehcho Region

Dehcho boreal caribou population monitoring

A substantial portion of boreal caribou range falls within the boundaries of the Dehcho Region. Boreal caribou have recently been designated as threatened by COSEWIC.

Larter, Nic ENR Dehcho Fort Simpson, NT

Permit No: 5030 **Species Studied**: Moose **Region**: DC **Location**: Dehcho Region

Moose population monitoring

There were concerns voiced over moose abundance along high use areas in the Dehcho Region by all Dehcho communities during a wildlife workshop conducted by ENR in September 2002.

Larter, Nic ENR Dehcho Fort Simpson, NT

Permit No: 5028 Species Studied: Wood bison

Region: DC **Location:** Fort Liard; Nahanni Butte area, in the vicinity of

Liard and South Nahanni River Valleys

Monitoring of the Liard wood bison population

The Nahanni wood bison population is currently afforded a measure of protection against infection with *Brucella abortus* (causes brucellosis) and *Mycobacterium bovis* (causes tuberculosis), by maintaining a bison free zone, to prevent contact with infected bison from Wood Buffalo National Park.

Lausen, Cori

Birchdale Ecological

Kaslo, BC

Permit No: 4812 **Species Studied:** Bats **Region:** SS **Location:** Near Fort Smith

Swarming survey of bats near Wood Buffalo Nationsl Park

Bat biodiversity has not been surveyed in the South Slave Region. A hibernaculum for bats is known in Wood Buffalo National Park, suggesting bats may be overwintering in the southern part of the Northwest Territories.

Machtans, Craig

Canadian Wildlife Service, Environment Canada

Yellowknife, NT

Permit No: 5034 Species Studied: Forest songbirds

Region: DC Location: Fort Liard

Long-term population monitoring of songbirds at Fort Liard, NT

Data collected regarding song bird population is used for conservation management and policy decisions at Environment Canada.

Madsen, Erik

Tibbit to Contwoyto Winter Road Joint Venture C/O Diavik Diamond Mines Inc.

Yellowknife, NT

Aerial and ground-based caribou surveys of the winter road

In response to concern from community members, the Joint Venture is interested in monitoring caribou movements along and near the Tibbit to Contwoyto winter road, to obtain a better understanding of caribou presence in the area.

McCallum, Dee Snap Lake Mine Yellowknife, NT

Permit No: 6881 **Species Studied:** Caribou, grizzly bear, wolverines, wolves

and falcons

Region: SS Location: Snap Lake

Wildlife effects monitoring program

This program is designed to detect, monitor and measure environmental effects that may impact wildlife habitat, changes to wildlife behaviour and distribution, and wildlife mortalities associated with the mine activities.

Mulders, Robert ENR North Slave Yellowknife, NT

Permit No: 6880 Species Studied: Wolverine

Region: SS Location: NE of Wekweètì (Daring Lake, BHP and Diavik

study areas)

Wolverine DNA sampling

A DNA sampling protocol has been developed, that can estimate relative wolverine abundance across large landscapes. This technique enhances the ability to quantify relative wolverine abundance, provide trend information for wildlife monitoring programs, and is anticipated to permit the modelling and assessment for cumulative impacts of anthropogenic activity on a regional scale.

Panayi, Damian

Golder Associates Yellowknife, NT water birds, and raptor

Region: SS Location: Kennady lake area

Gahcho Kué environmental monitoring

The purpose of these studies is to gather additional baseline information on wildlife in relation to the proposed Gahcho Kué project.

Popko, Richard ENR Sahtú Region Norman Wells, NT

Permit No: 5159 Species Studied: Mallards and northern pintail ducks

Region: SA Location: Willow Lake

Western Canada cooperative duck banding program at Willow Lake

Dabbling ducks are migratory waterfowl that are hunted throughout their range. Banding large numbers of ducks across their summer range before the start of the hunting season and then documenting band returns from successful hunters allows the researchers to plot the harvest distribution.

Rausch, Jennie

Canadian Wildlife Service

Yellowknife, NT

Permit No: 5402 **Species Studied:** All shorebirds

Region: IN Location: Kendall Island Migratory Bird Sanctuary

Arctic shorebird monitoring program

The arctic shorebird monitoring program was initiated in response to widespread shorebird population declines noted on migration routes through southern Canada and the United States.

Reford, Stephen

Darnely Bay Resources Ltd.

Toronto, ON

Permit No: 7419 Species Studied: Birds Region: IN Location: Near Paulatuk

Darnely Bay Resources Ltd. 2010-2012 field program - pre-activity nest surveys

Identify if birds are nesting in or near the identified drill and camp sites; identify which bird species are nesting in or near identified drill and camp sites.

Robertson, Myra

Canadian Wildlife Service

Yellowknife, NT

Permit No: 5405 **Species Studied:** All goose species

Region: IN Location: Anderson River Delta Migratory Bird Sanctuary

Population management of geese and swans in the Inuvialuit Settlement Region, using aerial surveys and banding studies

Information on bird numbers, distribution, survival and productivity is needed to determine if current local and international harvest levels are sustainable and to ensure that populations are conserved for the long-term use of the Inuvialuit and other people residing or hunting within the migratory range of these species.

Sharam. Grea

Rescan Environmental Services

Yellowknife, NT **Permit No: 5697**

Species Studied: Caribou, wolverine, arctic fox, red fox, grizzly bear, voles, lemming, hares, ground squirrels

Region: NS Location: Couageous Lake

Courageous Lake project wildlife baseline program

Seabridge Gold is conducting mineral exploration near Courageous Lake, NT. Baseline wildlife studies were conducted in the area in 1982-1983 and 2004-2005, but this is the first year that Rescan will be conducting this work.

Wood, Cindy

Canadian Wildlife Service Yellowknife, NT

Permit No: 6887 Species Studied: Herring gulls

Location: North shores of North Arm of Great Slave Lake Region: NS

from Boundary Creek west to Frank Channel

Chemical management plan wildlife national monitoring program

Herring Gulls and other gull species have been used as the typical sentinel species for containinants in aquatic environments for over 30 years. They are used because they are known to accumulate organic contaminants, nest in colonies, making egg collection relatively simple, and lay three eggs, but seldom rear more than 2 chicks, thus the removal of a single egg from a nest does not typically reduce breeding success.

Wortham, Jim

US Fish and Wildlife Service c/o Canadian Wildlife Service Yellowknife, NT

Permit No: 5762 Species Studied: Swans and ducks

Location: From southern border of the NWT to Mackenzie Region: IN, GW, NS, SS, DC, SA

Delta region

Cooperative waterfowl population surveys in the Northwest Territories

Information on bird numbers, distribution, and population trends is needed to determine if current local and international harvest levels are sustainable and to ensure that populations are conserved for the long-term use and appreciation by northern residents.

FISHERIES PERMITS 2011

Ash, Jason

De Beers Canada Inc. jasonash@ca.debeersgroup.com

Licence Number: S-11/12-3002-YK

Species: All fish species **Location:** Lakes at Snap Lake Mine Site

De Beers Snap Lake Mine Monitoring Program

De Beers Canada Inc. is planning to undertake a number of aquatic programs in 2011 to satisfy the requirements of the Water Licence and Fisheries Authorization for the Snap Lake Mine. Five distinct programs are planned for 2011:

Snap Lake: 1) an open water zooplankton program (includes phytoplankton from Northeast Lake); 2) a fall benthos program (includes Northeast Lake); and 3) a small bodied fish monitoring program at the embankments and reference areas.

Northeast Lake: 1) an open water phytoplankton program (includes Snap Lake); and 2) a fall benthos program (includes Snap Lake).

Bisaillon, Jean-Francois

Parks Canada - Western Arctic Field Unit jean-francois.bisaillon@pc.gc.ca

Licence Number: S-11/12-3028-YK

Species: Arctic char, lake trout Location: Cashe Lake

Uyarsivik Monitoring Program for Tuktut Nogait National Park

The objectives of this study were: 1) to document and assess the density and abundance of Arctic char and Lake trout fish habitat; and 2) to evaluate the biological integrity of the fish population.

Bisaillon, Jean-Francois

Parks Canada - Western Arctic Field Unit jean-francois.bisaillon@pc.gc.ca

Licence Number: S-11/12-3029-YK

Species: Dolly varden (searun), arctic grayling, **Location:** Joe Creek fish hole (Ivvavik

whitefish (unspecificed), clam (sp)

National Park)

Integrated freshwater and Dolly Varden Monitoring Program for Ivvavik National Park

The objectives of this study were: 1) to document and assess the Dolly Varden fish habitat; and 2) to provide juvenile Dolly Varden relative density estimates.

Blais, Jules

University of Ottawa Jules.Blais@uottawa.ca

Licence Number: S-11/12-3006-YK

Species: All fish Species Location: Mackenzie Delta Area

Effect of permafrost thawing on contaminant bioavailability to fish in the Mackenzie Delta The objective of this study was: 1) to test if lakes affected by permafrost thaw slumps have altered biochemical profiles compared to unaltered lakes; 2) to test if thaw slumping will affect contaminant delivery to ecosystems of freshwater lakes; 3) to test if higher contaminant loads will be found in fish and macroinvertebrates of permafrost thaw-slump lakes compared to those without thaw slumps.

Bohnet, Seth

Rio Tinto

seth.bohnet@riotinto.com

Licence Number: S-11/12-3030-YK

Species: Lake trout Location: Lac de Gras Area, Lac du

Sauvage

Diavik Diamond Mines Inc. Annual Effects Monitoring Program - Non-lethal mercury in lake trout

Diavik Diamond Mines Inc. (DDMI) is planning to complete a non-lethal analysis of mercury concentrations in Lake Trout in association with the annual Aquatics Effects Monitoring Program (AEMP) for the Diavik Mine on Lac de Gras. These data will be used to assess mercury concentrations in large-bodied fish.

Bourke, Robin

Golder Associates Ltd pvecsei@golder.com

Licence Number: S-11/12-3010-YK

Species: All fish species Location: Hill Creek

Hill Creek Fish habitat

The objective of this study was to assess performance of three newly constructed riffles and the culvert. It was proposed that by modifying the stream downstream of the culvert, fish passage could be improved due to changes in velocities within the culvert crossing. The objectives include hydrometric monitoring, water temperature monitoring and fisheries observation.

Cote, Jason

Cambria Gordon Ltd.
JCote@cambriagordon.com

Licence Number: S-11/12-3045-YK

Species: All fish species Location: Lac la Martre

La Martre River Hydro Power Project - 2011 Fall fisheries program

The objective of this study was to determine the potential effects of a ramping event (sudden shutdown of the power generation facility) on fish and fish habitat within the lower reaches of the

La Martre River. This information would be added to the overall fish and fish habitat effects assessment and will identify potential mitigation measures, if required. Specific objectives of the field program include: 1) Identify fish habitat values and fish life-stage use of the braided side channels, vegetated benches and ephemeral areas of the lower La Martre River; 2) Identify relative abundance of fish use of the braided side channels, vegetated benches and ephemeral areas of the lower La Martre River in comparison to the mainstem channel; 3) Determine depth range of habitat use for each species and life-stage observed. Collect depth cross-sections within the braided channels in the Tailrace to 8 km Downstream River Section.

Fortier, Martin

ArcticNet Inc., GIROQ, University Laval martin.fortier@arcticnet.ulaval.ca

Licence Number: S-11/12-3047-YK

Species: All fish species Location: Amundsen Gulf, Mackenzie shelf

and southern Beaufort Sea

ArcticNet 2011 expedition: Integrated regional impact study on the Coastal Western Canadian Arctic

The goal of the ArcticNet marine-based research program is to study on a long-term basis how climate induced changes are impacting the marine ecosystem, contaminant transport, biogeochemical fluxes, and exchange processes across the ocean-sea ice-atmosphere interface in the Canadian Arctic Ocean. Ultimately, the knowledge generated from this multi-year program will be integrated into regional impact assessments to help decision makers develop effective adaptation strategies for the changing coastal Canadian Arctic.

Gantner, Nikolaus University of Victoria gantnern@uvic.ca

Licence Number: S-11/12-3043-YK

Species: All fish species Location: Eskimo (Husky) Lake

Evaluation of hydro-climatic drivers of contaminant transfer in aquatic food

The objectives of this study were: 1) to characterize the lake-ice conditions and associated lake productivity and aquatic food webs along a hydroclimatic/salinity gradient within the Husky Lakes Watershed (HLW); 2) to develop and validate a process-based hydro-ecological food-web subcomponent coupling changes in hydro-climatology and ice conditions to water column productivity and food webs; 3) to explain uptake of Hg including isotopes as tracer/marker through a comparison of: i) spatial comparison of Hg bioaccumulation in food webs; ii) Hg stable isotope ratios in biota along a salinity gradient in the HLW and with lakes outside the HLW. 4) to review existing and new documentation of Traditional Knowledge (TK) on historical and present ice and climate conditions and fish biology and subsistence fisheries; and 5) to combine both knowledge bases to help develop future strategic monitoring of locally relevant sites.

Guthrie, Glen

Sahtú Renewable Resource Board rrco@srrb.nt.ca

Licence Number: S-11/12-3009-YK **Species:** Benthic invertebrates

Location: Unnamed Creek (Between Edie and Hodgson lakes) Bosworth Creek, Oscar

Creek, Jungle Ridge Creek

Location: Lac de Gras (Diavik)

Bosworth Creek macroinvertebrate study

The objectives of this study were: 1) to collect baseline information on benthic invertebrate community in streams within the Sahtú; 2) to work with the community and Sahtú Renewable Resources Board (SRRB) to develop a community-based stream biomonitoring program that is based on standard assessment techniques using benthic invertebrates; 3) to create a guideline package for high school students that will provide them with the tools to implement and maintain the monitoring program; 4) to initiate a monitoring study of lower Bosworth Creek in Norman Wells, where a weir was removed and flow restored in 2005; and 5) to identify benthic invertebrates from a variety of stream habitats to add to the local species list.

Hanna, Bruce

Fisheries and Oceans Canada bruce.hanna@dfo-mpo.gc.ca

Licence Number: S-11/12-3027-YK **Species:** Lake trout, slimy sculpin

Lake trout spawning assessment for shoals and dike exterior at Diavik Diamond Mine

The objective of this study was to assess the dikes isolating the open pits from Lac de Gras at the Diavik diamond mine to determine if the dike exterior (and nearby shoal) are currently being used by lake trout for spawning. Incidental information on other fish species use of the area will be collected on an opportunistic basis.

Hanna. Bruce

Fisheries and Oceans Canada bruce.hanna@dfo-mpo.gc.ca

Licence Number: S-11/12-3048-YK

Species: Invertebrates Location: Yellowknife Area

Preliminary sediment assessment of Frame Lake

The objectives of this study were: 1) to evaluate and map the bottom sediments of Frame Lake in terms of basic composition, distribution and overall quantitiy. Frame Lake is a non fish bearing lake with in the city of Yellowknife NT; and 2) to recover and examine short and long core sequences with a view to reconstructing the environmental history of the lake and suronding area and thereby better understanding curren lake conditions.

Harwood, Lois

Fisheries and Oceans Canada lois.harwood@dfo-mpo.gc.ca

Licence Number: S-11/12-3022-YK

Species: Ringed seal, bearded seal Location: Ulukhaktok Area - Coastal Marine

Waters

Harvest assessment of reproduction, condition disease and contaminants of ringed and bearded seals

The objectives of this study were: 1) to sample and measure ringed seals taken in the annual harvest in the Ulukhaktok (n=100) area, using reproductive status and body condition as indicators of ecosystem productivity and fluctuations in the seal population; 2) to examine the aspects in objective 1 in the context of regional ice conditions; and, 3) to co-ordinate with, and provide samples for, "stock health" related studies, such as disease and contaminants; and 2) In community-based programs, to sample and measure any bearded seals that happen to be taken in the annual harvest in the Ulukhaktok (n=5) areas, to examine reproductive rates, growth, condition and prey preferences.

Hayward, David

Summit Environmental Consultants Inc. dh.summit-environmental.com

Licence Number: S-11/12-3024-YK

Species: All fish species Location: Hay River Area

Northwest Territories Highway 1 maintenance and widening

Baseline assessments of watercourses in two segments (Section 2 and Section 8) will be surveyed to identify issues and complications with the proposed road works for Highway 1; surveys will focus on fish presence/absence, barriers to fish access, road crossings and habitat values in order to determine mitigation measures during and following construction. The watercourse crossings to be assessed are: Section 2 (5 small stream crossings); and Section 8 (3 crossings, km 67-83 at least 3 crossings).

Hayward, David

Summit Environmental Consultants Inc. dh.summit-environmental.com

Licence Number: S-11/12-3044-YK

Species: All fish species Location: Project area: (113° 53' 36.42" to

113° 43'57.57")

Northwest Territories Highway 4 (Ingraham Trail) upgrades

The objective of this study was to conduct baseline assessments of watercourses within the project site will be surveyed within the proposed road works for Highway 4; surveys will focus on fish presence/absence, barriers to fish access, road crossings and habitat values in order to determine mitigation measures during and following construction.

Howland, Kimberly

Fisheries and Oceans Canada kimberly.howland@dfo-mpo.gc.ca

Licence Number: S-11/12-3015-YK

Species: Lake trout Location: Great Bear Lake

Monitoring of lake trout stocks in Great Bear Lake

The objectives of this study were: 1) to monitor size and age structure, fecundity (egg number per female), growth and mortality of lake trout populations from Dareli (Keith), Turili (McVicar), Kwit tla (McTavish), Tugacho (Dease) and Tirato (Smith) Arms of Sahtú (Great Bear Lake). These data will be used for stock assessment purposes and to follow changes in the biological characteristics

of lake trout stocks over time; 2) to determine the extent of movements (if any) by lake trout in Great Bear Lake through molecular genetics and a tagging program; and 3) to monitor species composition and if sufficient data are available, presence, size structure and other biological characteristics of by catch and invertebrate species.

Howland, Kimberly

Fisheries and Oceans Canada kimberly.howland@dfo-mpo.gc.ca

Licence Number: S-11/12-3020-YK

Species: Cisco Location: Great Bear Lake

Cisco diversity in Great Bear Lake, Northwest Territories

The objectives of this study were: 1) to examine the morphological, meristic and life history characteristics of archived ciscos collected from Great Bear Lake over the past 7 years to test the hypothesis that there are multiple forms/species including shortjaw; 2) to conduct targeted sampling and examination of characteristics for cisco from deeper regions of Great Bear Lake (>50 m) to increase sample size and increase the range of surveyed habitat; and 3) to compare Great Bear Lake cisco with shortjaw cisco identified in other lakes to verify species identification and provide information that will feed into broader questions regarding the taxonomy of shortjaw cisco.

Howland, Kimberly

Fisheries and Oceans Canada kimberly.howland@dfo-mpo.gc.ca

Licence Number: S-11/12-3025-YK

Species: Arctic grayling, dolly varden (searun)

Location: Firth River System, Little Fish River, Malcolm River System, Joe Creek fish hole (Ivvavik National Park), Babbage River System, Herschel Island

Population studies on Dolly Varden and biological sampling of Arctic grayling from the NT & Yukon N

The objectives of this study were: 1) to capture Dolly Varden <400 mm and Arctic grayling passing in front of a DIDSON camera during the summer in the Firth River, Joe Creek and the Babbage River in order to confirm species ratio; 2) to conduct mark recapture studies: recapture tagged Dolly Varden in the Big Fish and Babbage rivers, and then tag Dolly Varden from the Big Fish River (n= 500), Babbage River (n= 500), Firth River (n=1000) and Joe Creek (n= 500); 3) to investigate the presence (yet to be confirmed) of Dolly Varden in the Malcolm River and Fish Creek and possibly seine and tag 100 Dolly Varden from each river system for mark-recapture; 4) to collect samples of Arctic grayling from the Big Fish, Babbage and Firth rivers and Joe Creek during the fall in order to obtain biological information such as length, weight, age, sex and maturity and diet, and tissue samples; and 5) to sample subsistence catch of Dolly Varden for biological information at Herschel Island and Ptarmigan Bay, Yukon.

Howland, Kimberly

Fisheries and Oceans Canada kimberly.howland@dfo-mpo.gc.ca

Licence Number: S-11/12-3026-YK

Species: Arctic char Location: Hornaday River

Arctic char monitoring at Hornaday and Brock Rivers, NT, 2011

The objectives of this study were: 1) to maintain char monitoring project and continue to provide information on status and life history of Arctic char captured at the mouth of the Hornaday and Brock rivers; and 2) to provide important support information for the formulation, delivery and compliance of the Paulatuk Char Management Plan.

Jones, Paul

University of Saskatchewan, Toxicology Centre paul.jones@usak.ca

Licence Number: S-11/12-3046-YK

Species: All fish species Location: Slave River

Fish health study at the Slave River and Slave River Delta

The objective of this study was to study the health of the fish in the Slave River and at the Slave River Delta. A number of different fish species will be collected to study metals, organic contaminants, gene expression and histology.

Knopp, Jennie Trent University

Licence Number: S-11/12-3011-YK

Species: Arctic char, lake trout

Location: Raddi Lake, Fish Lake, Sachs River, Thesiger Bay, Sachs Harbour Area, Kuptan Lake, Middle Lake, Ulukhaktok Area – Freshwater Rivers/Lakes, Capron Lake

Baseline fish study and char community-based monitoring plan for Sachs Harbour

The objective of this project was: 1) to conduct baseline studies for freshwater fish and anadromous fish in the areas surrounding Sachs Harbour and Ulukhaktok to provide local HTC, FJMC and DFO fisheries managers with information on baseline conditions fish and changes to fish growth in the region. Arctic Char will be sampled from the Sachs Harbour and Ulukhaktok areas for analysis through a PhD thesis at Trent University (Jennie Knopp). Arctic Char (and potentially Lake Trout) otoliths, tissue samples and stomach contents will also be sampled for follow-on analysis. This data will provide information towards: 1) Length, weight and age frequencies for the thesis project analysis; 2) to obseve responses of char to variability in context of climate change; 3) to establish baseline conditions, current char biodiversity and provide a point of reference against which future changes can be compared; 4) to provide data and knowledge towards the creation and implementation of community-based monitoring plans in Sachs Habour; 5) to obtain information on contaminant levels in locally-consumed fish (as requested by the community); 6) to support future (follow-on) aassessments of genetic and morphological variation, genetic population structure and variation in the dynamics of char populations within the Sachs Harbour area; and, 7) to develop/provide information towards long-term community-based monitoring plans for char in the area.

Landry. François

Rescan Environmental Services flandry@rescan.com

Licence Number: S-11/12-3004-YK

Species: All fish species Location: EKATI Dimagnd Mine

EKATI Diamond Mine - Fish monitoring and sampling program 2011

Rescan Environmental Services Ltd. (Rescan) was retained by BHP Billiton Canada Inc., the operator of the EKATI Diamond Mine, to monitor fish populations on the EKATI claim block during the open-water season of 2011.

The objectives of this study were: 1) to monitor the use of the stream habitat in the Panda Diversion Channel by spawning fish, particularly Arctic grayling; 2) to compare the biological characteristics of fish populations in the PDC with those in nearby reference streams; 3) to determine if the handling associated with non-lethal tissue plug sampling of round whitefish increases mortality of these fish as compared to typical handling (length and weight measurements, fin clipping); and 4) to determine if slimy sculpin can be used as a sentinel species by comparing bile polycyclic hydrocarbons of sculpin to round whitefish as well as determining the distribution of metals in the various tissues of the sculpin.

This project is a continuation of the long-term monitoring program at EKATI. The PDC has been monitored every year since 1998, although the specific sampling objectives have varied among years.

Landry, Francois

Rescan Environmental Services flandry@rescan.com

Licence Number: S-11/12-3041-YK

Species: Lake trout Location: Counts Lake, Nanuq Lake,

Vulture Lake, Lac de Gras

Nitrate toxicity study using lake trout

The objective of this study was to collect gametes from lake trout (Salvelinus namaycush) either from Counts Lake, Nanuq Lake, Vulture Lake, Lac de Gras or Great Slave Lake. Gametes will be collected by hand-striping and then brought immediately to Vancouver, BC where they will be combined to create fertilized eggs. The eggs and resulting larvae will be exposed to various concentrations of nitrate to determine toxicity levels for long and short term nitrate exposure for developing lake trout larvae.

Leonard, Deanna

Fisheries and Oceans Canada Deanna.Leonard@dfo-mpo.gc.ca

Licence Number: S-11/12-3016-YK

Species: All fish species Location: Slave River and Delta

Fish health study at the Slave River and Slave River Delta

The objective of this study was to study the health of the fish in the Slave River and at the Slave River Delta. Six different fish species will be collected to study metals, organic contaminants, gene expression and histology.

Low, George Dehcho AAROM geobarbgeo@hotmail.com Licence Number: S-11/12-3033-YK

Species: Lake trout, walleye, burbot, whitefish Location: Greasy Lake, Yendo Lake, Long Lake, Trout Lake, Deep Lake Ekali Lake,

Sanguez Lake, Gargan Lake

Updating data on mercury levels in food fish species in lakes used by Dehcho communities

The objective of this study was to update data on mercury levels of certain fish in Dehcho lakes that have shown elevated levels in previous studies. This project will also determine if any other traditional fishing lakes are affected. The data from this project will also be used by scientists to help determine which factors are leading to increases of mercury in fish. It is also an objective of this study to provide traditional knowledge data on which lakes and fish species are being used for subsistence by the first nations.

Machtans, Hilary

Golder Associates Ltd hilary_machtans@golder.com

Licence Number: S-11/12-3040-YK

Species: All fish species Location: Yellowknife Bay, Yellowknife

River, Baker Creek, Trapper Creek

Giant Mine Remediation Project (GMRP) - Sediment gap analysis and sampling plan for Baker Creek

The objective of this study was to determine the magnitude and extent of sediment contamination and toxicity in Baker Creek, assess benthic invertebrate abundance and community composition, assess fish abundance and community composition in adjacent waters, and measure contaminant concentration in benthic invertebrate and periphyton tissues. The information collected will be used to provide a basis for evaluating the potential risk of historical sediment contamination in Baker Creek as outlined by the Framework for Addressing and Managing Aquatic Contaminated Sites under the Federal Contaminated Sites Action Plan (FCSAP). Benthic invertebrate and periphyton samples were collected from within Baker Creek and a reference area (the Yellowknife River). A fish survey will also be conducted in the adjacent Trapper Lake/Creek and Martin Lake.

Maier, Kris

Gwichya Renewable Resource Council kmaier@grrb.nt.ca

Licence Number: S-11/12-3036-YK **Species:** Dolly vardon, arctic grayling

Stony Creek Dolly Varden assessment

The objective of this study was to confirm the presence of northern form Dolly Varden char in the Stony Creek watershed and determine if they are anadromous and a distinct genetic population.

Mason, Kristine

Golder Associates Ltd. kristine mason@golder.com

Licence Number: S-11/12-3008-YK-A1

Species: All fish species Location: Waterbodies near the Gahcho

Kué Project

Location: Stony Creek

Gahcho Kué Project

De Beers Canada Inc. (De Beers) has been conducting fish and fish habitat baseline studies for a number of years in the area of the Gahcho Kué Project. The overall objectives of this year's

program are: 1) to collect additional fish and fish habitat baseline information on Lake N11 and the proposed Reference Lakes (Lake X6 and the Unidentified Lake); 2) to confirm northern pike presence/absence in the N watershed; 3) to evaluate fish passage at previously identified barriers between Area 8 and Lake 410 (Streams L1a, L1b, L1c, L3 and M4); and 4) to collect lower level trophic data at Lake N11, Reference Lakes (X6 and Unidentified Lake) and small lakes in the L and M watersheds downstream of Kennady Lake (L2, M1, M2, M3 and M4).

McPherson, Morag

Fisheries and Oceans Canada morag.mcpherson@dfo-mpo.gc.ca

Licence Number: S-11/12-2000-HR-A4

Species: All fish species **Location:** Baker Creek, Yellowknife River,

Yellowknife Bay

Baker Creek and fish use and habitat study

The objectives of this study were: 1) to observe and describe fish habitat use (spawning, feeding and rearing) by adult, juvenile and young-ofyear fish in Baker Creek; 2) to quantify the number of spawning fish in Baker Creek; 3) to determine fish use and assess habitat in Reaches 1, 2, 3, 4, 5 and 6 of the creek on the Giant Mine site, areas of the creek above the mine site (upper Baker Creek); and 4) to capture fish and benthic invertebrates and assess habitat in portions of Yellowknife River and Yellowknife Bay as a reference areas.

McPherson, Moraq

Fisheries and Oceans Canada morag.mcpherson@dfo-mpo.gc.ca

Licence Number: S-11/12-3017-YK

Species: All fish species Location: Waterbodies near Tundra Mine

Site

Fisheries survey and baseline construction monitoring program for Tundra Mine

Rescan Environmental Services Ltd. (Rescan) was retained by Contaminants and Remediation Directorate (CARD), Indian and Northern Affairs Canada (INAC), NT Region to conduct baseline fish and fish habitat surveys during the summer of 2011 for the Tundra Mine Project in the Northwest Territories. The objectives of the 2011 construction baseline monitoring program are to measure the potential effects on fish from 1) to monitor potential effects associated with treatment and discharge of arsenic-impacted tailings on fish spawning and feeding within lake tributaries; and 2) potential effects associated with treatment and discharge of arsenic-impacted tailings will be achieved through larval fish surveys in lake tributaries, which will also identify critical spawning habitat.

McPherson, Morag

Fisheries and Oceans Canada morag.mcpherson@dfo-mpo.gc.ca

Licence Number: S-11/12-3037-YK

Species: All fish species **Location:** Alpha Lake, Sherman Lake,
Gamma Lake, New Control Lake, Mill Lake

Rayrock Mine Site - Aquatic Health Monitoring Program

Under the Federal Contaminated Sites Action Plan (FCSAP), the Government of Canada, led by Indian and Northern Affairs, is conducting an aquatic health monitoring program in 2010 at the remediated Rayrock Mine Site located in the Tłįchǫ Region. This includes the collection of fish, fish tissue (muscle and organ), benthic invertebrates, and sediment samples for metals analysis. This program is designed to monitor the health of the environment as part of a Detailed (site-specific) Quantitative Environmental Risk Assessment, part of the Long-term Monitoring program at Rayrock Mine.

Mochnacz, Neil

Fisheries and Oceans Canada mochnacznj@dfo-mpo.gc.ca

Licence Number: S-11/12-3034-YK

Species: All salmon species **Location:** Mackenzie River, Beaufort Sea,

Mackenzie Delta Area

Pacific salmon distribution in the western Arctic

The objective of this study was to collect information on the distribution of Pacific salmon in the western Arctic. Once the study establishes a basic understanding of the distribution for each species this study will monitor annual catches to track dispersal. A major shift in these distributions could serve as an indicator of environmental change.

Mochnacz, Neil

Fisheries and Oceans Canada mochnacznj@dfo-mpo.gc.ca

Licence Number: S-11/12-3037-YK

Species: Bull trout, arctic grayling, slimy sculpin **Location:** Prairie Creek, Funeral Creek,

Nahanni River

Habitat use of bull trout in the Prairie Creek watershed

The objective of this study was to improve our understanding of habitat use for Bull Trout found in Prairie Creek and lower South Nahanni river system. Specific objectives are: 1) to document spawning and winter habitat use; 2) to document baseline habitat reference conditions which can be used to monitor change over time; and 3) to document seasonal movements by Bull Trout populations in the lower South Nahanni River Watershed.

Nicol, Sandra

Stantec Consulting Ltd. sandra.nicol@stantec.com

Licence Number: S-11/12-3023-YK

Species: All fish species **Location:** Ring Lake, Ball Lake, Buck Lake,

Drizzle Lake, Murky Lake

Fisheries baseline studies for Avalon Rare Metals Inc.

Det'on Stantec was retained by Avalon Rare Metals Inc. in 2008 to conduct an environmental baseline assessment of the Thor Lake area in support of their proposed Thor Lake Rare Earth Metals Project. The

specific objectives of the 2011 Fisheries field program are to collect a third year of data from five lakes to: 1) determine fish presence and distribution (if any); 2) establish relative abundance of fishes in fish-bearing waterbodies; 3) further characterize baseline water and sediment quality and the aquatic invertebrate community structure (phytoplankton, zooplankton, and macroinvertebrates)

Panayi, Damian

Golder Associates Ltd damian_panayi@golder.com

Licence Number: S-11/12-3042-YK

Species: All fish species **Location:** Bluefish Lake, Prosperous Lake,

Yellowknife River

NTPC Bluefish Hydro Repairs

The objective of this study was to describe the terrestrial and aquatic environment in the Yellowknife River between Prosperous Lake and Bluefish Lake. This may include investigating vegetation, water quality, benthic invertebrate populations, and fish. The results will be used to determine the most suitable means of fish habitat compensation, and to guide mitigation during construction of the new dam for Bluefish Hydro. Further, monitoring of methyl mercury in the aquatic ecosystem is required by water licence MV2009L4-004. Lethal sampling of small and large bodied fish is anticipated.

Parent, Bruce

Imperial Oil Ltd.

bruce.d.parent@esso.ca

Licence Number: S-11/12-3007-YK

Species: All fish species Location: Bosworth Creek

Bosworth Creek aquatics and fisheries monitoring program

The objective of this study was to determine current water and aquatic habitat quality, and establish baseline conditions for evaluation of future monitoring data. In order to achieve the project objectives, sampling stations within and downstream of the Imperial Oil Limited Norman Wells operations will be compared to upstream reference stations, located beyond the possible influence of the development, and against an appropriate range of regional data indicating natural variability (if available). Stations will be monitored seasonally to characterize natural temporal variability in conditions, and to identify potential trends in water quality over time. Regarding fish; the main objective is to collect information on fish species presence/absence, habitat suitability and potential seasonal habitat use within Bosworth Creek, as information pertaining to this watercourse is limited.

Pink. Melissa

Rescan Environmental Services Ltd.

Licence Number: S-11/12-3003-YK

Species: Slimy sculpin, lake trout, lake whitefish **Location:** Courageous Lake Project Area

Baseline fisheries surveys for the Courageous Lake Project

Rescan Environmental Services Ltd. (Rescan) was retained by Seabridge Gold Inc. to conduct baseline fish and fish habitat surveys during the summer of 2011 for the Courageous Lake Project in the Northwest Territories. The information collected during the baseline program will assist in preparing for an Environmental Impact Statement (EIS). The objective of the 2011 surveys will be to further characterize the streams and lakes in the area so that the effects assessment can be carried out and appropriate monitoring programs and fish habitat compensation plans can be developed. This includes an understanding of the structure and growth patterns of the fish community inhabiting the lakes and streams, the quality of fish habitat prior to development of the site, the current tissue metal concentrations of slimy sculpin in four potentially impacted and two reference lakes, as well as the diets (stomach contents) of those sculpin. As well, a human health study is proposed to sample edible species (lake trout and lake whitefish) for tissue metals in two lakes in the project area.

Pink, Melissa

Rescan Environmental Services Ltd.

Licence Number: S-11/12-3012-YK

Species: All fish species Location: Waterbodies in the Pine Point

Project area

Baseline fisheries surveys for the Pine Point Project

Rescan Environmental Services Ltd. (Rescan) was retained by Tamerlane Ventures Inc. to conduct baseline fish and fish habitat surveys during the summer of 2011 for the Pine Point Project approximately 42 km east of Hay River, Northwest Territories. The information collected during the baseline program will assist in preparing a Developers Assessment Report (DAR). The objective of the 2011 surveys will be to characterize the streams and lakes in the area with respect to fish community and fish habitat so that the effects assessment can be carried out and appropriate monitoring programs and compensation plans can be developed (if required). This includes an understanding of the structure and of the fish community inhabiting the lakes and streams and the quality of fish habitat prior to development of the site.

Pink, Melissa

Rescan Environmental Services Ltd.

Licence Number: S-11/12-3038-YK

Species: All fish species Location: L-Shaped Lake, Spanner Lake,

Paddle Lake

Post-discharge ecological assessment of the Dam 2 Discharge Pathway, Colomac Mine Site, NT

Rescan Environmental Services Ltd. (Rescan) was retained by the Contaminants and Remediation Directorate (CARD), Aboriginal Affairs and Northern Development Canada (AANDC), NT Region to conduct an ecological assessment of the Colomac Mine site, Northwest Territories. To assess the potential ecological impacts of the discharge from Tailings Lake to the receiving environment, a before-after-control-impact (BACI) design was designed. Pre-discharge baseline ecological data were collected by Rescan in 2005 for lakes along the discharge path (L-Shaped Lake, Paddle Lake, and Lake 315) and from a reference lake (Spanner Lake). It was intended that these baseline data be compared to post-discharge ecological data that would be collected in 2011 to determine whether the discharge of Tailings Lake water has had an effect on

downstream aquatic ecosystems. Potential effects to fish from effluent will be determined through sampling fish for biological assessments (internal and external) and fish tissue analyses.

Robertson, Dale

Golder Associates Ltd dale_robertson@golder.com

Licence Number: S-11/12-3039-YK

Species: All fish species Location: Slemon Lake

Tłycho Fish Camp

The objectives of this study were: 1) to document and combine traditional knowledge of fish in the Behchokö area with conventional fish sampling methods; 2) to obtain baseline data on mercury levels in fish consumed by Behchokö residents (SleemanLake and area); 3) to evaluate mercury levels found in fish in terms of species and size/age relationships, habitat use, trophic status; and 4) to establish a monitoring program to identify contaminant levels and changes in levels through time for fish in the Marian/Russell Lakes area and possibly as part of a larger watershed stewardship program.

Sawdon, Lorraine

Fisheries and Oceans Canada lorraine.sawdon@dfo-mpo.gc.ca

Licence Number: S-11/12-3005-YK-A1 **Species:** Lake trout, burbot, northern pike

Location: L-Shaped Lake, Spanner Lake,

Paddle Lake

Calibration and comparison of two standardized protocols: Nordic netting protocol

The objectives of this study were: 1) Conduct the BsM (Broad-scale Fish Community Monitoring) protocol on Alexie, Baptiste, Chitty and Drygeese lakes in the NWT. The netting effort will serve as the recapture event for the population estimate on Alexie lake; 2) To conduct a modified Petersen's population estimate on large-bodied fish species (Lake Trout, Burbot, Northern Pike and Lake Whitefish), on Alexie Lake in the NWT. Mark-recapture data will be used to calibrate the proposed 2011 BsM Protocol data and the existing 2008 Nordic Protocol data. Multiple gear types will be used in order to catch sufficient numbers of large-bodied fish; 3) Conduct comparisons using the 2008 Nordic Netting data and proposed 2011 BsM data a) comparison between the two protocols b) comparison among the four northern lakes, and c) comparison between northern (NWT) and southern (Ontario) regions.

Tonn, William

bill.tonn@ualberta.ca

Licence Number: S-11/12-3014-YK-A2

Species: All fish species **Location:** Lac de Gras Area Lakes

Improving habitat connectivity to enhance productive capacity of arctic freshwater ecosystems

The objective of this study was to study systems comprise four 'treatment' lakes (locally known as M1, M2, M3, W1), their outlet streams, and 3 'control' lake-stream ecosystems in the Lac de Gras watershed. Treatments will involve physical alteration of the lakes' outlets to improve their suitability for fish passage with the goal of thereby enhancing accessibility of the lakes to fish; one stream will also receive in-stream habitat enhancement to promote spawning and rearing. The

study is to evaluate the effectiveness of the habitat manipulations and assess the biotic and abiotic responses to these treatments. In 2011, the study will continue baseline, pre-treatment monitoring of the habitat and biota in the study systems.

Toyne, Melanie

Fisheries and Oceans Canada melanie.toyne@dfo-mpo.gc.ca

Licence Number: S-11/12-3013-YK

Species: Inconnu Location: Great Slave Lake, Buffalo River

Closed Area

Assessment of Buffalo River Inconnu

Buffalo River Inconnu have been the subject of concern for many years. Data for stock assessment purposes has been collected in periodic years for decades. The last sampling program took place in 2008. In 2011, the plan is to set gillnets and lethally sample up to 250 Inconnu for biological and CPUE information, to be used to update stock status.

Ware, Chaz

Triton Environmental Consultants Ltd. cware@triton-env.com

Licence Number: S-11/12-3032-YK

Species: Brook sticleback, walleye, burbot Location: Tributary to the Hay River 60° 10'

25.184" N & 116° 41' 40.632" W

Canadian National Railway Company culver replacement mile 316.7 Meander Subdivision The objective of this study was to conduct fish sampling to remove fish from the worksite prior to instream work associated with culver replacement. Fish will be salvaged from the worksite to

avoid fish stranding during isolation of the worksite.

Wiatzka. Gerd

SENES Consultants Limited gwiatzka@senes.ca

Licence Number: S-11/12-3021-YK

Species: All species (excludes marine mammals) Location: Great Slave Lake Area

Great Slave Lake Area Mines: Site assessment and remediation planning

This objectives of this study were: 1) to collect fish at two abandoned mine sites at Outpost Island and Blanchet Islands in Great Slave Lake as part of the human health and ecological risk assessments; and 2) to collect fish that are part of the traditional and commercial fishery to determine whether chemicals of concern are elevated in fish collected near the mines.

Wilcockson, John

Hatfield Consultants jwilcockson@hatfieldgroup.com

Licence Number: S-11/12-3035-YK

Species: Arctic grayling, slimy sculpin, bull trout **Location:** Prairie Creek

Fish habitat/fish usage assessment - Proposed Prairie Creek Mine Dilution Zone

The objective of this study was to assess fish utilization of a section of Prairie Creek that would contain the proposed mine efflient exfiltration trench (diffuser) and the downstream mixing zone. Study area would include the location of the trench, as well as 100m up and downstream. Of specific interest is the potential migration of bull trout past the mine site on their way to spawn in upstream creeks.

Zhu, Xinhua

Fisheries and Oceans Canada xinhua.zhu@dfo-mpo.gc.ca

Licence Number: S-11/12-3031-YK

Species: All fish species Location: Great Slave Lake

Monitoring and assessing the cumulative impacts on important fish population productivity and community integrity

This objectives of this study were: 1) to develop a standard monitoring framework addressing cumulative impacts on major fish population productivity and community integrity. To approach this goal, the team will create methodologies, in accordance with the Cumulative Impacts Monitoring Program Pathways Approach to Protocol Development, to quantify changes in the aquatic environment, fish population index, growth and productivity, species richness and community diversity, and association between abiotic factors and fish populations; set up a framework to integrate information on hydroclimatic dynamics, population biological characteristics, fishing effort, harvest, and social economy; and establish collaborative partnerships between researchers, resource users, Aboriginal community, and decision makers to ensure the effectiveness and representative of Great Slave Lake fisheries and ecosystem; and 2) to construct an operational assessment network for Great Slave Lake fisheries and ecosystem changes to ensure the sustainable exploitation of fisheries production. To accomplish this goal, the team will compare the mesh-specific gear selectivity between single- and multi-mesh gillnet to standardize species-specific population indices, model sustainable population productivity and management tactics for Lake Whitefish under the scenarios of changing exploitation and virgin population biomass, estimate quantities for Lake Whitefish fisheries management, such as total allocable catch (TACs), maximum surplus production (MSP), and precautionary biological reference points (PBRPs); 3) to conduct management strategy evaluation (MSE) for effective utilization while minimizing by-catch effects on Inconnu and Lake Trout; and 4) analyze the association of multi-species interactions (fish community diversity) and environmental parameters.

BIOLOGY 2012

Buddle, Chris McGill University Ste-Anne-de-Bellevue, QB chris.buddle@mcgill.ca

File Number: 12 402 861 Licence No: 15091 (Multi-year licence)

Region: GW **Location:** Dempster Highway, between kilometers 0 and 540 including Rock River; Mackenzie lowlands; Engineer

Creek

Ecological structure of northern arthropods: Assessing community structure and biodiversity across an environmental gradient

Northern parts of Canada are home to many different species of insects, spiders and other arthropods. Researchers from McGill University have been working to document the species of insects and arachnids living in the North, in an effort to benchmark northern biodiversity so that climate change and other environmental effects can be better understood. In 2012, a field team from McGill University travelled into the Northwest Territories (NT), between the Yukon-Northwest Territories border, but not further than the Peel River. The field team spent less than one day in the NT. The objective was to collect some insects and arachnids, and the bulk of the collections were done in the Yukon. However, about 50 pseudoscorpions (small arachnids, related to spiders), several dozen caterpillars, some wolf spiders and beetles were collected in the NT. These arthropods were collected by hand, and by searching in the tundra. Determining exactly what species were collected is a time-consuming task and the data are not yet fully processed. Preliminary observations, however, suggest the species are similar to what field teams collected along the Dempster Highway in past years.

Hansen, Ken

Husky Oil Operations Limited Calgary, AB ken.hansen@huskyenergy.com

File Number: 12 402 880 Licence No: 15160 (Multi-year licence)

Region: SA Location: Within the boundaries of EL 462/EL 463 in the

Central

Husky EL462/463 fisheries baseline study

Travel to the study area (near Norman Wells) occurred in September 2012. Baseline fish studies involving the electrofishing of several streams and the netting of several lakes were attempted (as outlined in the proposed scope of work). The studies were not completed due to increased water levels and flow rates in the streams from higher-than-normal snow fall in the area in mid-

September. Several of the proposed study sites were inspected, however, no activities associated with electrofishing or netting were performed. As such, no fish were removed from the streams or lakes of interest.

Harris, Allan

Northern Bioscience Thunder Bay, ON aharris@tbaytel.net

File Number: 12 402 876 Licence No: 15137 (Multi-year licence)

Region: DC Location: Hotsprings (approximately 60) in and around

Nahanni National Park

Nahanni Aster status report

A survey for Nahanni Aster (*Symphyotrichum nahanniense*) was conducted in 2012 in and around Nahanni National Park Reserve. Fifty springs were surveyed, including the four previously documented sites for this species, an undocumented site reported by Parks Canada staff, and 44 sites that had not previously been surveyed for this species. The species was rediscovered at all five known sites and newly discovered at a sixth site. The number of stems (mature individuals) ranged from 200 to over 1500 per site, with an overall total of more than 5000 stems.

Jones, Paul

University of Saskatchewan Saskatoon, SK paul.jones@usak.ca

File Number: 12 402 867 Licence No: 15014 (Multi-year licence)

Region: SS Location: The mouth of the Slave River Delta at Fort

Resolution; Slave River within the municipal boundaries of

Fort Smith

Fish health study in the Slave River and the Slave River Delta

In 2012, researchers continued fish sampling on the Slave River in collaboration with the Government of the Northwest Territories. These collections were carried out in conjunction with sampling on the Athabasca River. Fish were collected by gillnetting with the assistance of community members. A total of 185 fish were collected, including: 7 burbot, 57 goldeye, 49 jackfish, 45 walleye and 27 whitefish. After collection, the fish were subject to a detailed external and internal health examination and samples were collected for contaminant analysis. Samples are currently being analyzed for chemicals associated with oil sands activities in Alberta. These tested chemicals include: PAHs (polycyclic aromatic hydrocarbons), naphthenic acids and metals.

Krizan, Julia

IMG-Golder Corporation Inuvik, NT jkrizan@golder.com

File Number: 12 402 664 **Licence No:** 15097

Region: IN Location: From the Ikhil Production Facility to the proposed

South Parsons Lake well

Fish habitat survey and bathymetry survey for the South Parsons Lake gas supply project, NWT

The proposed work was not completed, and is not anticipated to be completed in the future.

Marken, Sandra

ConocoPhillips Canada Calgary, AB sandra.l.marken@conocophillips.com

File Number: 12 404 792 **Licence No:** 15100

Region: SA Location: Central Mackenzie Valley

Environmental assessment for ConocoPhillips EL470

Assessments of the EL470 study area were conducted to gather baseline information for the environmental assessment process, in support of the ConocoPhillips' 2012-2013 winter drilling program and future exploration activities. Biophysical components studied included: soils, vegetation, wildlife, species at risk and plants and animals of traditional interest. Bathymetric surveys, volumetric calculations and surface water sampling were conducted on 14 water bodies within the study area; however, only eight of these water bodies and the Mackenzie River were found to be suitable for potential project use. Four groundwater monitoring wells were installed, with groundwater sampling planned for 2013. The geotechnical activities originally planned for 2012 have been deferred to 2013. Community meetings were held in Tulít'a and Norman Wells during the week of May 22, 2012.

Muir, Andrew

Great Lakes Fishery Commission. Ann Arbor, MI United States amuir@alfc.org

File Number: 12 402 849 **Licence No:** 15078

Region: SA **Location:** Narakay Islands; Dease Arm; the eastern side of

McTavish Arm extending from the Doghead Peninsula in the north to the Superstition Islands in the south; Great Bear

Lake

Lake trout diversity in Great Bear Lake: Do deep-water forms exist?

A type of lake trout lives in the deep waters of several big lakes in Canada, including Great Slave Lake and the Great Lakes. The objective of this research was to determine if this deep-water lake trout also existed in Great Bear Lake. Fourteen gillnets in deep-water regions (> 50 m) southwest of the Narakay Islands and McTavish Arm were set, resulting in only 132 lake trout caught. This low number of caught fish suggests that very few fish live in the deep-water parts of the lake. No other kinds of fish were caught in the nets. Each fish was sampled to determine their age, growth, reproductive condition, body shape, and diet. These samples are still being processed. Preliminary information suggests that deep-water lake trout - similar to the ones in Great Slave Lake - do not occur in the areas of the McTavish Arm. Two fish that were caught near the Narakay Islands in the Dease Arm did resemble the deep-water lake trout form, but further analysis is ongoing.

Renaud. Claude

The Canadian Museum of Nature -- Research and Collections Division Ottawa, ON crenaud@mus-nature.ca

File Number: 12 402 873 Licence No: 15057 (Multi-year licence)

Region: DC, SS

Location: Lower Martin River; Mackenzie River Basin;
Harris River mouth opposite and north of Fort Simpson;
Mackenzie River near Fort Providence about 75 km

Mackenzie River near Fort Providence about 75 km downstream from the outlet of Great Slave Lake; Mackenzie River at the outlet of Great Slave Lake near Big Island; Lower Hay River in approximately the 16 km before it enters Great Slave Lake; upper Slave River at the water intake

reservoir of the water treatment plant in Fort Smith

Evolution of arctic lampreys

The purpose of this research was to study the two lamprey species found in Northwest Territories (NT). To date, there is very little information about both the arctic lamprey (Lethenteron camtschaticum, formerly known under the scientific name Lampetra japonica) and the Alaskan brook lamprey or darktail lamprey (Lethenteron alaskense). This study is part of a larger scale research initiative about lampreys of the genus Lethenteron across the northern hemisphere (Eurasia and North America). For the NT study, the objectives of this ongoing project are: (1) To collect a total of about 30 larvae (at least 60 mm in total length) for each of the two species: (2) To collect adults of each species, describe their gross morphology and pigmentation and produce an identification key. A total of about 10 adults of each species would desirable to corroborate the identity of the larval; (3) Compare the DNA sequences between the two lamprey species; (4) Perform a histological examination of lamprey larvae of different sizes (5 larvae per 10 mm increments from 30 mm up to 120 mm total length) for the two species to follow the development of their ovaries and determine if they contain one or two generations of eggs. This gives a combined total for the two species of 90 larvae. To date, researchers have done preliminary work based on a single mitochondrial gene on samples from Japan (L. camtschaticum) and Alaska (L. alaskense) and found no differences. Work is ongoing.

Robb, Tonia

Rescan Environmental Services Yellowknife, NT trobb@rescan.com

File Number: 12 402 766 **Licence No:** 14996

Region: NS Location: Waterbodies located within the EKATI claim

block

EKATI aquatic monitoring program, 2009-2013

In 2012, five monitoring projects were ongoing in the lakes and streams of the Koala, King-Cujo, and Pigeon watersheds, where EKATI mine infrastructure are located. The monitoring programs are: (1) Aquatic Effects Monitoring Program (AEMP); (2) Surveillance Network Monitoring Program (SNP); (3) Panda Diversion Monitoring (PDC) Program; (4) seepage monitoring program; and (5) Air Quality Monitoring Program (AQMP). The objective of the AEMP was to assess the current conditions in the lakes and streams of the Koala and King-Cujo watersheds to determine whether there have been any mine effects. The assessment incorporated meteorology, hydrology, water quality and physical limnology, phytoplankton, zooplankton, benthos and fish data. Data analyses for the 2012 are currently being completed and will be submitted in April 2013 to the Wek'èezhìi Land and Water Board (WLWB). The main objective of the SNP was to confirm EKATI's compliance with effluent quality criteria in its water license. The data are reported monthly to Aboriginal Affairs and Northern Development Canada and are available on the WLWB online registry. Results from the past 13 years have shown that the PDC is successfully providing fish

habitat and that vegetation is establishing itself along its banks. A full habitat assessment of the PDC was conducted during July 2012 to provide information for the design and installation of the habitat enhancements that were proposed prior to the completion of the program. Although many plants have successfully re-colonized riparian and bank slope areas, results indicate that the establishment of in-stream vegetation was occurring at a slower rate. In response, in-stream vegetation mats were transplanted to the PDC during the summer. Further analysis of this work is in progress. Monitoring of seepage from the waste rock storage areas (WRSAs) at Misery, Fox and Panda-Koala continued in 2012. Seepage samples were collected in June to correspond with snow melt, and again in September prior to freeze up. The data will indicate the extent of metal leaching from the WRSA. Results are reported to the WLWB. Air quality was monitored using high volume air sampling (HVAS), partisols, continuous ambient monitors (CAM) and dustfall measurements as a part of the AQMP.

Sibbald, Carey

Deton' Cho Stantec Yellowknife, NT carey.sibbald@stantec.com

File Number: 12 402 879 **Licence No:** 15155

Region: NS, SS Location: Within Avalon's Thor Lake Property; two

reference lakes

2012 baseline aquatic program for Avalon Rare Metals Inc. proposed Thor Lake rare earth element project

In 2012, Deton'Cho Stantec conducted two field programs for aquatics baseline studies of Avalon Rare Metals' proposed Thor Lake Rare Earth Metals Project. In September 2012, water, plankton (phytoplankton and zooplankton), sediment, and benthos sampling was undertaken. In October 2012, the fieldwork consisted of water sampling. Sampling was carried out at nine lake stations. Results of the 2009-2012 field programs indicate neutral to basic water and very low nutrient levels at all stations. There were large fluctuations in some general and metal parameters (primarily during winter in small, shallow lakes that developed low oxygen conditions under the ice). Sediment characteristics varied, though generally lake sediment had: (1) high phosphorus, nitrogen and organic carbon content; and (2) ranging metal levels (below the ability to detect higher than Canadian guidelines). Chlorophyll a levels varied among lakes and seasons, but in general, most of the lakes are considered oligotrophic (poor in nutrients and plant life and rich in oxygen). Biota data from 2012 is still under analysis.

Tonn, William

University of Alberta Edmonton, AB bill.tonn@ualberta.ca

File Number: 12 402 724 **Licence No:** 15051

Region: NS Location: Small headwater lakes and their outlet streams

into Lac de Gras

Improving habitat connectivity to enhance productive capacity of arctic freshwater ecosystems

Diavik Diamond Mines, Inc., located on Lac de Gras, has undertaken two habitat compensation projects on headwater lake and stream systems near the mine site. Lake outlet streams at two sites were modified to improve fish passage and thus the ecological "connectivity" among these headwater lakes and with Lac de Gras. One habitat manipulation - called M-lakes - occurred in

the fall of 2011, while the second habitat project - called West Island - occurred the year following. In 2012, ongoing sampling continued at all reference sites and the West Island site for hydrology, water quality, habitat characteristics, primary producers, invertebrates, and fish. Results from the habitat assessments are similar to previous years. Stream riparian zones are dominated by shrubs, forbs, grasses, mosses, and boulders, while streambeds are sparsely vegetated and composed predominantly of inorganic fines, boulders, and pebbles. Water quality is similar among all streams, but does show seasonal variation. Streams feature low, diffuse flows and cascades that obstruct fish movement from Lac de Gras. Stream electrofishing and hoop netting continues to document low abundances of slimy sculpin, juvenile burbot, and arctic grayling. All lakes are oligotrophic. Riparian zones of lakes are similar to those of streams. Littoral zones are dominated by boulders and inorganic fines. Lake fish communities and species abundances vary among lakes, but are made up mainly of arctic grayling, lake trout, round whitefish, burbot, longnose sucker, and slimy sculpin. Macroinvertebrates samples from the streams are still being analyzed. Further research examined the post-winter recolonization of streams by macroinvertebrates and lake trout feeding ecology. For macroinvertebrates, larger, more mobile taxa tended to colonize from downstream areas, while smaller, less mobile taxa drifted in from upstream or colonized from the zone below the stream bed. Few specimens colonized the streams from the air. Diets of lake trout were more specialized in lakes with fewer potential fish competitors, with lake trout mainly feeding on zooplankton. This is compared to the broader, more littoral-oriented diets of trout co-occurring with multiple competitor species. Growth rates did not differ between the lake types, but trout were in better body condition in lakes with fewer potential competitors. The preliminary evaluation of the M-Lake habitat manipulation indicated that two of three streams at the M-lakes site were ineffective at increasing fish passage and would likely need modifications. A third stream at M-lakes did improve fish passage and required no further work.

Trimble, Annika

Aurora Research Institute Inuvik, NT atrimble@auroracollege.nt.ca

Region: IN, GW **Location:** In and around Inuvik

Northern native seed development field trials

In 2005, three field trial sites were established in partnership with the Inuvik Community Greenhouse to assess Northwest Territories seed collections under natural conditions on disturbed sites. Each field site was established in a different type of habitat (i.e. gravel pit, clay slope, peat firebreak). Beyond the initial site clearing, seeding, and transplanting, these plots received no additional care. Each year the field sites are surveyed to assess the survival of transplants, germination of seeded plots, as well as the vigour and flower production of both. In 2012, the field plots continued to demonstrate the following trends: (1) higher survival and vigour in plots with transplants, as opposed to low germination in plots which were seeded; (2) increasing ground cover and recovery at clay and gravel pit sites, and low survival and vigour at the peat site; and (3) notable vigour in the legume species at the clay and gravel sites. These field plots will continue to be monitored for several more years to assess site recovery and long-term survival of the transplants.

CONTAMINANTS 2012

Blais, Jules University of Ottawa Ottawa, ON jules.blais@uottawa.ca

File Number: 12 404 800 Licence No: 15128 (Multi-year licence)

Region: DC **Location:** Tathlina Lake/Cameron River Delta; Lakes in and

around Yellowknife

The Arctic in flux: How has recent climate change affected contaminant transport and uptake in aquatic arctic systems?

The Cameron Hills, located along the Alberta/Northwest Territories border, are the largest actively producing oil and gas fields in the Northwest Territories, and yet the impacts of these activities, in particular the release of polycyclic aromatic hydrocarbons (PAHs) downstream, has not been investigated. Lake sediments are natural archives of physical, biological, and chemical information that can track environmental change in lakes, including contaminant histories. In September 2012, in collaboration with the Ka'a'gee Tu First Nation in Kakisa, sediment cores were obtained from four lakes in the Cameron Hills that are adjacent to oil and gas wells. Two sites were located downstream of the development in the Cameron River Delta (at Tathlina Lake), and three reference lakes. These cores were used to help determine whether oil and gas production in this region is releasing PAHs and other contaminants into the ecosystem, and whether contaminants are being transported downstream to Tathlina Lake. Samples of benthic invertebrates from streams in the Cameron Hills that are upstream and downstream of development are still being analyzed for PAH and metal contamination to assess whether contaminants are bioaccumulating in the food web.

Blowes, David

University of Waterloo Waterloo, ON blowes@uwaterloo.ca

File Number: 12 402 843 Licence No: 14993 (Multi-year licence)

Region: NS Location: Lac de Gras mine site at Diavik Diamond Mines

Waste rock studies at a diamond mine site

The objective of this ongoing research is to investigate the processes related to water quality and quantity draining from experimental waste rock piles that are located in areas of continuous permafrost. Waste rock piles are mounds of rock removed from open-pit and underground mines. The quality of water draining from a waste rock pile is determined by: (1) the combined effects of

oxygen transport in the air phase; (2) biogeochemical processes that control mineral weathering rates; (3) the release of heat and dissolved constituents due to sulfide mineral oxidation; and (4) hydrologic processes that control unsaturated water flow. The transport of dissolved constituents is further affected by the formation and subsequent dissolution of secondary minerals. Three instrumented experimental waste rock piles were constructed from 2004 to early 2007 at the Diavik Diamond Mine. Instruments in the pile include: basal lysimeters; basal drain; thermistors; time domain reflectometry probes and moisture sensors; tensiometers to measure near-surface infiltration; soil water solution samplers; air permeability probes; air pressure sensors; thermal conductivity access ports; gas sampling ports for oxygen and carbon dioxide; and microbiology access conduit and pyrite growth medium. In 2010, three 40 m deep boreholes were drilled into the operational waste dump and a series of instruments similar to those in the test piles were installed. In 2011, two additional 40 m boreholes and one 80 m borehole were installed and instrumented. Data from these instruments will be compared to data from the test piles to evaluate differences in measurement scale. Data collection and analysis, including modeling that incorporates climate change, continued in 2012.

Dewar, David

Northwest Territories Power Corporation Hay River, NT ddewar@ntpc.com

File Number: 12 402 878 Licence No: 15153 (Multi-year licence)

Region: NS Location: Bluefish Lake

Bluefish Lake mercury monitoring

No research was conducted under this research licence in 2012.

Evans, Marlene Environment Canada Saskatoon, SK marlene.evans@ec.gc.ca

File Number: 12 402 681 **Licence No:** 15129

Region: SS Location: Around Great Slave Lake (the East Arm near

Łutsel K'e; the West Basin near Fort Resolution; the West

Basin near Hay River)

Spatial and long-term trends in persistent organic contaminants and metals in lake trout and burbot from the Northwest Territories

This ongoing research project has been investigating the changing contaminant levels in Great Slave Lake fish since the early 1990s. In 2012, 20 lake trout from Hay River, 20 pike and 20 burbot from Fort Resolution, and 20 lake trout and 20 burbot from Łutsel K'e were shipped (frozen) whole by community members to Saskatoon where length, weight, and age were determined and samples submitted for metals (including mercury) and persistent organic contaminant analyzes; pike and Łutsel K'e burbot were analyzed for mercury only. Mercury concentration has been increasing, although levels remain below the 0.5 ppm guideline. Organic contaminant concentrations have not been increasing and some such as: HCH and DDT, have decreased in concentration due to their decreased usage. The results of 2012 findings will be presented in a 2012 Northern Contaminants Program (NCP) report. This is an ongoing study under the NCP.

Farrell, Rory

EBA, A Tetra Tech Company Yellowknife, NT rfarrell@eba.ca

File Number: 12 402 686 **Licence No:** 15144

Region: NS, SS **Location:** Approximately 70 km east of Yellowknife

EBA, Five Mine Sites ESA update and remedial action plan

No summary was submitted for this licence. This project is not in compliance with licencing requirements.

Gantner, Nikolaus (Klaus)

University of Victoria Victoria, BC gantnern@uvic.ca

File Number: 12 402 868 Licence No: 15046 (Multi-year licence)

Region: IN Location: In and around Inuvik and Tuktoyaktuk; Yaya

Lake; Noell Lake; Big Lake; Husky Lakes; 16 lakes along the 177 road; 4 lakes in the Tuktoyaktuk-Inuvik road corridor

Evaluation of hydro-climatic drivers of contaminant transfer in aquatic food webs in the Husky Lakes Watershed (Inuvialuit Settlement Region, NWT)

The goal of this ongoing research is to identify and quantify the physical, chemical and ecological processes that effect contaminant transfer in Arctic aquatic food webs in response to observed and predicted climate variability and changes in the Husky Lakes watershed. In May and August 2012, the research team completed two successful sampling trips to Husky Lakes, both based out of Inuvik and Tuktoyaktuk. During May, the research crew accessed the southern Husky Lakes to collect samples before continuing on to Tuktoyaktuk for further sampling. All fish samples were collected from fish caught at Husky Lakes by local fisherman. Less than 50 fishes were donated for sampling from 26 families in Tuktoyaktuk. In August, researchers returned to Inuvik to conduct open water sampling at all study lakes. The fieldwork was very successful, with collections of multiple fish species and near-shore invertebrate communities at all study sites. Fifteen traditional knowledge interviews were conducted in February through August 2012 in the region. All TK interviews are currently being translated and transcribed and verification through participants is planned for early 2013.

Geddes, Robert

AMEC Environment & Infrastructure Calgary, AB brian.geddes@amec.com

File Number: 12 404 862 **Licence No:** 15125

Region: SA **Location:** The south shore of the Great Bear River between

Bear River Landing and Lower Shipyard

Great Bear River summer of 2012 site assessments

No research was conducted under this licence in 2012.

Livingstone, Steve

Franz Environmental Inc. and SENES Consultants Ltd. Ottawa, ON

slivingstone@franzenvironmental.com

File Number: 12 402 811 **Licence No:** 15150

Region: SA **Location:** Along the Canol Trail

Environmental site assessments, Canol Trail, NT

SENES Consultants Ltd. and Franz Environmental Inc. were retained by Public Works and Government Services Canada – Northern Contaminated Sites Program to complete a case study for petroleum hydrocarbon stability and ecological integrity at discrete oil spill sites along the Canol Trail, in 2012. The purpose of the 2012 case study program was to characterize possible oil (petroleum hydrocarbon) related impacts to soil, groundwater and/or surface water at 11 selected spill sites identified during the Phase II Environmental Site Assessments. Each spill site was selected based on its suitability to characterize potential historical oil impacts within the surrounding geological and environmental area. The methods utilized to implement the field program included drilling boreholes and/or test pits (as soil conditions allowed), taking near surface and at-depth soil samples, and installing groundwater/active layer monitoring wells. A number of overall site characteristics including: oil spill morphology and mobility, petroleum hydrocarbon observations, site impacts and recovery were summarized in the report titled "Case Study Program for Petroleum Hydrocarbon Stability and Ecological Integrity" prepared by SENES/Franz and dated July 2, 2013.

Low, George

Dehcho First Nations Hay River, NT geobarbgeo@hotmail.com

File Number: 12 402 857 Licence No: 14999 (Multi-year licence)

Region: DC Location: Cli Lake; Little Doctor Lake; Sibbeston Lake;

Tsesto Lake: Blackwater Lake: Fish Lake: Jean Marie River-

McGill Lake; Deep Lake; Trout Lake

Updating data on mercury levels in food fish species in lakes used by Dehcho communities During 2012, collection of fish was carried out in the following locations: Sanguez Lake, Willow Lake, Trout Lake, Tathlina Lake and Big Island Lake. 20 fish were collected from each lake and biologically sampled. Muscle samples were taken and analyzed for mercury. The work and results have been presented to the community and community workshops have been set up to explain the research and encourage eating fish.

Macdonald, Colin

Northern Environmental Consulting Pinawa, MB northern@granite.mb.ca

File Number: 12 402 333 **Licence No:** 15172

Region: SA Location: Keith Arm of Great Bear Lake offshore from

Déline

The continuation of a community-led fish monitoring study in Déline, NT 2012

This study was designed to test for metals and radionuclides in fish species around Déline and to provide the people of Déline with advice on whether the fish are safe to eat. Major concerns are: (1) radionuclides (found at Port Radium and other mines in the eastern Sahtú); and (2) mercury (found in fish throughout the Northwest Territories). The high levels of mercury in some fish species can lead to health advisories, where Health Canada recommends that pregnant women and children eat less of the fish to avoid possible effects from mercury exposure. During November and December 2012, community members collected lake whitefish, lake trout and herring using gill nets in the waters off the eastern end of Déline. A total of nine lake whitefish, 17 lake trout and 10 herring were measured and aged, and muscle and liver tested for a wide range of chemicals. Data were combined with results from 2009 and 2010 to provide a comprehensive set of data to test differences between species and to determine if there were any causes for concern. In total, 39 trout, 26 whitefish and 26 herring were analyzed from 2009 to 2012, giving a solid basis to provide conclusions. The average concentration of mercury in all three fish species was well below Health Canada's guideline of 0.5 mg/kg for commercial sale. Only one lake trout sample (0.72 mg/kg ww), caught in 2010 was above the guideline. The mercury concentration in all whitefish and herring were far below the guideline. Mercury increased with the size of the fish in all three species tested in this study, but only larger lake trout approached the Health Canada quideline. Organochlorine pesticides, which move into the Northwest Territories from southern North America and Asia, were tested in 2010 and were found to be present in the fish of Great Bear Lake, but the concentrations are very low and are not a health concern.

McLachlan, Stephane

University of Manitoba Winnipeg, MB mclachla@cc.umanitoba.ca

File Number: 12 402 863 **Licence No:** 15036

Region: SS **Location:** Fort Resolution; Fort Smith

Multi-scale environmental health implications of the Athabasca oil sands for aboriginal communities in Alberta and Northwest Territories

The overall goal of this project was to better understand and communicate how the environmental and human health of downstream aboriginal communities are affected by industrial activity associated with the Athabasca tar sands and large-scale hydro developments. In June 2012, researchers conducted video interviews with community members in Fort Chipewyan (Alberta), Fort Smith and Fort Resolution about environmental changes and community concerns arising from industrial activities impacting the Slave River and Peace Athabasca River deltas. The research was shared with community collaborators through community newsletters. These newsletters were an attempt to create a communication network for communities to share their knowledge and concerns with each other and for the scientific community to communicate their research approaches and results. Two thousand copies of these newsletters have been printed for distribution to First Nation and Métis Nation research partners.

Nash, Tyler

Queen's University Kingston, ON tjnash@hotmail.com

File Number: 12 402 872 **Licence No:** 15055

Region: NS Location: Along Baker Creek near Giant Mine; Outside of

Yellowknife

An investigation of arsenic speciation and toxicity in Baker Creek sediments from Giant Mine in the Northwest Territories, Canada

The objective of this research project was to assess the sediment quality of Baker Creek. Researchers sampled arsenic contaminated dirt and water from the Giant Mine. Sampling was conducted in locations believed to be both highly and lightly contaminated. Sediments were dug up using a trowel. Water was filtered and collected into large jars. The research revealed that arsenic exists as highly toxic arsenic trioxide in some heavily contaminated areas. Many other forms of arsenic were also discovered, as the arsenic at Giant Mine is very complex. Liveorganism toxicity tests were conducted on all of the collected dirt and it was discovered that some of the dirt samples are very toxic to fish.

Osawa, Akira

Kyoto University Sakyo-Ku, Kyoto Japan aosawa@kais.kyoto-u.ac.jp

File Number: 12 402 492 Licence No: 15013 (Multi-year licence)

Region: GW, SS

Location: Forest stands adjacent to and along Highway #5;

around Wood Buffalo National Park; the Dempster Highway between the north shore of Campbell Lake and Rengleng

River

Structure, carbon dynamics, and silvichronology of boreal forests

The main objective of this ongoing fieldwork was to collect data in jack pine and black spruce forests on annual movement of organic matter and carbon. During the 2012 field season, a total of ten scientists participated in the study of carbon dynamics in the forests of black spruce, jack pine, or dwarf birch in the Inuvik and Fort Smith area. Soil and plant samples were collected allowing the researchers to examine: soils; fine roots; mycorrhizal hyphae (thin, white, thread-like things related to mushrooms); aboveground litter (amount of falling leaves and branches); tree growth and mortality; and growth history. Changing trends in the amounts of forest materials and forest ecosystems are crucial to understanding carbon dynamics of these forests. This in turn relates to the effect of forest vegetation on climate change (what scientists call feed-back). New efforts this year included: (1) collection of stem core or stem disc samples from two stands in Inuvik and two stands in Fort Smith areas for estimation of stands' growth history; (2) collection of soil samples for examining processes of phosphorous dynamics in permafrost soils; and (3) initiation of growth measurement in mycorrhizal hyphae (that may grow to mushrooms) in jack pine and black spruce stands near Fort Smith. Efforts remain ongoing to quantify the movement of these materials in northern forests.

Purdy, Colin

Queen's University Kingston, ON cpurdy.rocks@gmail.com

File Number: 12 402 871 Licence No: 15054 (Multi-year licence)

Region: NS Location: Thor Lake

Geochemical and mineralogical controls on the low-temperature aqueous mobility of rare earth elements (REE) in mine waste from the Nechalacho deposit, NWT

No summary was submitted for this licence. This project is not in compliance with licencing requirements.

Reimer, Kenneth

Royal Military College of Canada Kingston, ON reimer-k@rmc.ca

File Number: 12 402 528 **Licence No:** 15095

Region: NS **Location:** In and around Yellowknife, with a focus on mine

properties

Non-toxic arsenic in mushrooms and plants from Yellowknife

The purpose of this research was to determine and compare total arsenic and arsenic species in edible and non-edible fauna and soil from contaminated and uncontaminated locations (mine properties and roadsides/parks). In 2012, a total of 16 different mushroom species were collected from contaminated and uncontaminated locations. Mushrooms were analyzed for total arsenic and arsenic species to determine if the forms of arsenic are toxic or non-toxic. Puffballs, shaggy manes and meadow mushrooms from contaminated areas had total arsenic concentrations ranging from 2.8 to 36 mg/kg dry weight, mainly in the non-toxic form of arsenic, arsenobetaine. Associated soils were also collected and their microbial communities were characterized by DNA analysis, and are currently being compared. The arsenic distribution in a negus tailings mushroom was mapped and showed arsenobetaine is mainly found in the cap and the gills of the mushroom. A field bioaccessibility test was developed and performed on three species of plants collected from five locations. Results were lower than lab methods (research beginning in Fall 2013 to seek to explain this). Total arsenic has not yet been analyzed in all the sampled plants but of those analyzed so far (from Ndilo), the concentrations ranged from 0.04 to 70 mg/kg dry weight. Arsenic compounds present in the plants have not yet been determined, but future work will address this.

Reinfort, Breanne

University of Manitoba Freshwater Institute - Department of Fisheries and Oceans Winnipeg, MB b.reinfort@gmail.com

File Number: 12 402 870 Licence No: 15042 Region: IN Location: Sachs Harbour

Arctic contaminants: Exploring effective and appropriate communication between Inuvialuit communities and researchers

The purpose of this research was to understand indigenous perspectives on contaminants, contaminant research, and how research is communicated and made accessible within communities. From February to March 2012, four focus groups occurred with 12 individuals (six males and six females). Overall, 41% of the Sachs Harbour population contributed to this project since 2009. Early analysis suggests, a person's perceptions of communications processes about environmental hazards may influence their perceptions about the hazard itself. Examples of these processes are: communication methods (how) and communicators/sources (who). Perceptions affect an individual's reception, comprehension and compliance with messages about contaminants. Continuing analysis is investigating the emerging connections and importance between how risk communication is approached and carried out and how contaminants are perceived. In addition, the roles that trust and relationships play in message development, reliability and retention are also being studied. Focus group discussions highlighted the

importance of researchers spending time in the community to explain their research and connect with locals. This research identified a gap in community members' understanding as to why researchers are interested in studying contaminants. Participants identified an absence of basic background knowledge about mercury (the contaminant being discussed), which prevented them from engaging with research updates. Criteria were established for creating a community-based pamphlet about mercury, which is in progress. Fieldwork is complete, and interview participants continue to verify their transcripts.

Stern, Gary

Department of Fisheries and Oceans Canada - Freshwater Institute Winnipeg, MB gary.stern@dfo-mpo.gc.ca

File Number: 12 402 869 **Licence No:** 15041

Region: SA, DC **Location:** In and around Trout Lake; Kelly Lake

Impact of climate change on freshwater fish mercury levels

The purpose of the research was to collect fish from lakes and tributary rivers under consumption advisories due to mercury and compare results with historical record. This research also collected and analyzed sediment cores from these lakes to see how lake ecosystem changes relate to fish mercury levels. In 2012, researchers analyzed sediment cores from Trout Lake and Kelly Lake to see how mercury was deposited over time, and how this compares with mercury in lake trout that were collected and analyzed over the past three or four years. Fish from both lakes showed marked increases from earlier collections (1970s and 1980s). Increases over time were also observed in the sediment cores, suggesting that fish mercury levels were linked in some manner to in-lake processes. There was a link between algae-derived organic matter and mercury. This implies that a longer ice-free period (due to climate change) is leading to more mercury entering the lake, which in turn can enter fish. More fish collections are needed to improve the statistical power of these trends, which are currently weak.

Wiatzka, Gerd

SENES Consultants Ltd. Richmond Hill, ON gwiatzka@senes.ca

File Number: 12 404 778 **Licence No**: 15146

Region: NS, SS **Location:** The former Copper Pass Mine located on

Sachowia Lake

Copper Pass Mine: Environmental site assessment and remediation planning

The Copper Pass Mine is a former small-scale mine on Sachowia Lake, within the East Arm of Great Slave Lake. The nickel mine operated for only a few years and has since been abandoned for approximately 40 years. The objective of this research was to assist Aboriginal Affairs and Northern Development Canada (AANDC) with its ongoing work to clean up abandoned mines across the Northwest Territories. The site investigation (conducted August 1-3, 2012) involved the collection of soil, vegetation and waste rock. A detailed topographic survey was conducted using GPS equipment, as well as site reconnaissance to find roadways, hazards, borrow materials, etc. No mechanical equipment was used, excluding a float plane for site access. Results of the site investigation were incorporated with previous site assessment findings and confirmed there are aspects of the site that will require remediation. The finalized document is still being prepared, and will be provided to AANDC for ownership and distribution.

ENGINEERING 2012

Patterson, R. Tim
Carleton University
Ottawa, ON
tpatters@earthsci.carleton.ca

File Number: 12 406 054 Licence No: 15106 (Multi-year licence)

Region: NS Location: Along the length of the Tibbitt to Contwoyto

Winter Road

Paleoclimatological assessment of the central Northwest Territories: Implications for the long-term viability of the Tibbitt to Contwoyto winter ice road

The purpose of this research was to develop a comprehensive sub-decadal to centennial-scale late Holocene climatic history along a 2° latitudinal gradient in the central Northwest Territories. Over 100 lakes along the route of the Tibbitt to Contwoyto Winter Road were analyzed for water property data (e.g. pH, conductivity), substrate characteristics (e.g. LOI, grain-size, BSi), nutrient loading, water geochemistry (e.g. F/U, Fe/Mn, DIC/DOC) isotopes (C/N) and environmentally available metals. This dataset was used to develop training sets and transfer functions based on micropaleontological proxies (thecamoebians, diatoms and chironomids). A large number of freeze cores were also collected from these lakes. Use of a freeze core microtome has permitted subsampling of freeze cores to mm-resolution (2-5 years). Time series analysis results indicate that throughout the late Holocene there was been considerable climate variability with winter and summer signals often becoming decoupled. The Pacific Decadal Oscillation and North Atlantic Oscillation have contributed to step-wise temperature changes, as these phenomena vary between positive and negative phases. The computer model results indicate that the Tibbitt to Contwoyto Winter Road should remain viable for the coming decades as winter temperatures will only rise slightly. Precautions should be taken during El Niño years though. 2012 was the final year of this research project.

Trimble, Annika

Aurora Research Institute Inuvik, NT atrimble@auroracollege.nt.ca

File Number: 12 406 058 Licence No: 15151 (Multi-year licence)

Region: IN **Location:** Storm Hills (68°53'1.14"N, 133°56'55.65"W)

Wind energy monitoring at Storm Hills: 2012-2014

In October 2012, wind monitoring equipment was installed on an existing 150 foot communications tower in the Storm Hills area. The datalogger is being powered by solar panels,

and data are being transmitted by satellite. Wind speed data will be collected for two years and then used to verify the feasibility of developing a wind energy project in the area.

HEALTH 2012

Goodman, Karen

University of Alberta Edmonton, AB karen.goodman@ualberta.ca

File Number: 12 408 149 Licence No: 15009 (Multi-year licence)

Region: GW Location: Aklavik; Tuktoyaktuk; Sachs Harbour; Fort

McPherson

The Aklavik *H. pylori* project

The objective of the ongoing Aklavik *H. pylori* project is to develop a comprehensive approach to investigating community health problems related to Helicobacter pylori infection (a bacteria that can be found in the gastrointestinal tract) in Northwest Territories communities. The purpose of this research project is to identify public health solutions aimed at reducing related health risks. and to obtain representative data from diverse settings in northern Canada for informing regional public health strategies for reducing health risks from *H. pylori* infection. From fall 2011 to summer 2012, research team members followed up with the Aklavik *H. pylori* project participants to assess the onset of new infections and re-infections. Additionally, microbiology results were shared with Aklavik community members. Since April 2012, 16 new participants joined the Inuvialuit Settlement Region (ISR) H. pylori Project in Tuktoyaktuk. Project team members and the team's lead gastroenterologist have made two additional trips to offer medical consultations to project participants and initiate the treatment component of the project. Planning for expansion to other communities in the ISR is underway. In addition, the Fort McPherson H. pylori project was launched in June 2012. As of September 2012, 179 residents of Fort McPherson have joined the project and completed a urea breath test. Recruitment is still ongoing, and the endoscopy and treatment components of the project are being planned. These are expected to be held in March 2013.

Hammond, Merryl

Consultancy for Alternative Education Baie d'Urfé, QB merryl.hammond@videotron.ca

File Number: 12 408 148 Licence No: 14997 (Multi-year licence)

Region: IN Location: Aklavik; Ulukhaktok

Changing the "culture of smoking": Community-based participatory research to empower Inuvialuit communities

The purpose of this research was to empower local Inuvialuit community members to better understand tobacco and its health and other effects and to motivate them to consider quitting or

reducing their use of tobacco. This community-based participatory research project began in Aklavik and Ulukhaktok in 2007. 2012 was the final year of the project. At the beginning of the projects, participants completed baseline surveys that were analyzed and presented to communities in 2010. The first smoking cessation challenge, called the *Quit to Win Challenge* (open to smokers and non-smokers aged 8 and up) took place in January 2009. It was very successful with a total of 34% of eligible community members signing up. The second challenge, the *Be Smoke-free Challenge* took place in November 2010. Community responses were excellent, again with 34% of the eligible population of the two communities signing up. The third challenge, the *No Tobacco Week Challenge* was held in June 2011 in Aklavik and November 2011 in Ulukhaktok. This time, participation rates went up to 43% for both communities. More non-smokers than smokers entered (an equal number of girls and boys, and more women than men). In 2012, the final year of the project, participants completed an exit survey.

Hannon, Judy

Canadian Blood Services Edmonton, AB judy.hannon@blood.ca

File Number: 12 408 142 Licence No: 15029 (Multi-year licence)

Region: IN, GW, DC, SA, NS, SS **Location:** At health centres, hospital and clinic laboratories

throughout the Northwest Territories

RHD alleles in prenatal patients from northern Canada

The objective of this research is to ensure that current prenatal testing methods are appropriate for prenatal patients in the northern regions of Canada, and to learn more about the RHD genotype in northern populations. This project was initiated in 2006, but getting participants has been slow despite repeated recruitment efforts. Much is known about the RHD genetic make-up of ethnic groups world-wide but this information has never been compiled for the indigenous populations of northern Canada. The information is important because the test reagents - used for prenatal testing - are largely developed based on a Caucasian population. To date 72 maternal blood samples have been collected, DNA extracted and frozen in a molecular laboratory. A minimum of 80 samples are required to make the results statistically significant. This study is likely to continue at least until the minimum 80 samples are collected.

Janssen, Patricia

University of British Columbia Vancouver, BC patti.janssen@ubc.ca

File Number: 12 408 187 Licence No: 15162 Region: SS Location: Fort Smith

Outcomes of primary maternity care in Fort Smith, NWT

The purpose of this research was to look at outcome data that is available from a perinatal database developed during the last two years in Fort Smith and to describe the experiences of women who have given birth using the community-based midwifery service. Data were collected from medical charts at the Fort Smith Midwifery Program to evaluate the birth outcomes of moms and babies in Fort Smith from 2005-2011. Researchers collected information including: who the birth attendant was; if the woman transferred out of Fort Smith for the labour/delivery; if it was a preterm birth; if the woman had a Caesarean section; if the woman had any tearing from delivery; and gestational age and weight of the baby. The dataset is now being analyzed and compared to outcomes from the midwifery-led maternity centres in Nunavik, Quebec. Focus groups were also

held with several women who had used the Fort Smith Midwifery Program to discuss their experiences and the impacts that the program has had on their own and their family's lives.

Kuhn, Karen University of Bath Victoria, BC karen.kuhn@telus.net

File Number: 12 408 184 **Licence No:** 15058

Region: DC, NS Location: Yellowknife Primary Care Clinic and Stanton

Territorial Hospital; Fort Simpson Clinic; Behchokò Clinic;

Dettah Clinic

Evaluation of the electronic health record (EHR) system used in the Northwest Territories No research was conducted under this licence in 2012.

MacLeod, Martha

University of Northern British Columbia Prince George, BC macleod@unbc.ca

Region: IN, GW, SA, DC, NS, SS Location: Registered Nurses, Nurse Practitioners,

Licensed Practical Nurses and Registered Psychiatric

Nurses across the NWT

Nursing practice in rural and remote Canada II

The survey instrument was developed and revised during this year. It was tested outside of the Northwest Territories. No data were collected in the Northwest Territories in 2012.

Mitton, Craig

University of British Columbia Vancouver, BC craig.mitton@ubc.ca

File Number: 12 408 183 Licence No: 15010 (Multi-year licence)

Region: NS **Location:** Within the Stanton Territorial Health Authority

Achieving high performance in health care priority setting

The intent of this project was to develop an evaluative framework that identifies how health care organizations can be transformed to achieve excellence in priority setting and resource management. As part of this project, the researchers interviewed five members of the management team in Stanton Territorial Health Authority. This was one of six case studies conducted across Canada, in a range of large and small, urban and rural sites. Managers were asked to define the concept of 'high performance' in a priority setting and what resource allocation looked like to them. Then they were asked to assess their own organization's performance in relation to those ideas. Gathering these results, the researchers developed a framework for high performance, including 19 elements, within four broad domains. They represented areas that organizational managers should address in order to improve priority setting and resource allocation practice in the health care sector. None of the case study sites were identified by name in public presentations of the findings.

Scott, Shannon

University of Alberta Edmonton, AB shannon.scott@nurs.ualberta.ca

File Number: 12 408 186 **Licence No:** 15099

Region: NS **Location:** Stanton Territorial Hospital

TRanslating Emergency Knowledge for Kids (TREKK)

In Canada, the majority of children requiring emergency care are treated in general emergency departments. Evidence shows however that as many as 40% of children cared for in general emergency departments do not receive treatments for which clear evidence exists, and up to 20% of these children receive a treatment which has been shown to provide no benefit or in some cases even harm. This study, Translating Emergency Knowledge for Kids (TREKK), is a Pan-Canadian initiative aimed at ensuring that the latest research in pediatric emergency medicine is applied within general emergency departments. TREKK's long-term vision is to efficiently and effectively improve the outcomes of acutely ill and injured children cared for in all Canadian emergency departments. During this phase, TREKK will determine the knowledge needs of health care providers working in general emergency departments, and the families seeking care within these facilities. Through an established partnership with 32 general emergency departments across Canada (including at the Stanton Territorial Hospital), TREKK will work with staff, administrators and consumers to understand the existing knowledge gaps, needs and priorities in pediatric emergency medicine. Understanding the needs and preferences for knowledge mobilization among receptor communities is central to the mandate of TREKK. Phase one work is ongoing.

Young, Barbara

Stanton Territorial Hospital Chelsea, QB barbara.youngmd@gmail.com

File Number: 12 408 185 **Licence No:** 15074

Region: IN, GW, SA, DC, NS, SS Location: Database study using the Non-Insured Health

Benefits prescription drug database

Anticoagulation in Canada's north: A cost-effectiveness analysis of point-of-care INR testing at remote sites in the Northwest Territories and Western Nunavut.

No research was conducted under this licence in 2012.

PHYSICAL SCIENCES 2012

Anderson, Natalie

Colorado State University Fort Collins, CO United States n.kramer.anderson@gmail.com

File Number: 12 404 789 Licence No: 15093 (Multi-year licence)

Region: DC, SS **Location:** Along the Mackenzie drainage basin at the Slave

River Rapids

Big river wood dynamics in the Canadian subarctic

The primary objectives of this ongoing research are to: (1) evaluate fundamental controls on wood dynamics within the Mackenzie; and (2) develop an empirical predictive model to estimate future wood dynamics. In the summer of 2012, preliminary research was conducted on the Slave River near Fort Smith, Great Slave Lake, and Fort Simpson. This work mainly focused on networking, reconnaissance and gaining familiarity with the region. Data on log jam structure and size were collected from islands in Mountain and Cassette Rapids (on the Slave River), Paulette Island (near the Slave Delta) and along the shores of the Mackenzie and Liard (near their confluence). Time-lapse photography was taken at the Slave R. Fitzgerald gage to capture wood moving downstream during high flows. Repeat analog aerial photographs of the Pelican Islands (data usage allowed from the Department of Environment and Natural Resources and the Pelican Advisory Circle) from 1978 to 2004 were scanned and digital photographs from 2004 to present were acquired. Work will continue in 2013.

Armstrong, Terry

Government of Northwest Territories - Environment & Natural Resources Fort Smith, NT terry armstrong@gov.nt.ca

File Number: 12 404 750 Licence No: 15059 (Multi-year licence)

Region: DC, NS

Location: The Mackenzie Bison Sanctuary and areas to the west to Mills Lake; Birch Lake; Fawn Lake; Sharun Lake and

Second Lake. Tree core sampling took place south of

Behchokò, primarily along Highway 3

Landscape scale flooding in the Great Slave Lake Plain

This ongoing research project is designed to study the Great Slave Lake Plains. In 2012, researchers collected nine sediment cores from the bottom of eight lakes within the Mackenzie Bison Sanctuary. In addition, water samples were collected from those eight lakes plus three others in the area. Water samples were analyzed for a variety of water chemistry parameters,

including major ions, mercury and dissolved carbon. Analysis of the water chemistry was undertaken at Taiga Laboratory in Yellowknife. Sediment cores are now being analyzed to determine the algae (diatoms) living in lakes and how their populations may have changed as the lakes have expanded. Sediment analysis is being done to determine if mercury is becoming more abundant in the lake systems as the lakes have grown in size and flooded the forests and shrubs surrounding the lakes. In addition, the research team also traveled to Fort Smith where they gave presentations and took students from Aurora College out into the field to explain the research in Fort Providence.

Barber, David University of Manitoba Winnipeg, MB dbarber@cc.umanitoba.ca

File Number: 12 404 371 Licence No: 15044 (Multi-year licence)
Region: IN Location: In and around Sachs Harbour

An integrated sea ice project for BREA: Detection, motion, and RADARSAT mapping of extreme ice features in the Southern Beaufort Sea

Three groups are represented in the integrated sea ice project for BREA. The University of Manitoba contributions to the integrated sea ice BREA project focuses on: (1) the identification of hazardous ice features using remote sensing techniques; (2) understanding the ice motion relative to winds and ocean currents; and (3) obtaining seasonal temperature profiles of extreme ice features (>4m thick). In April 2012, the research group deployed 13 GPS beacons, 11 on multi-year ice floes located 80-100 nautical miles (NM) west of Banks Island and two GPS beacons on an ice island (glacial ice) located 48 NM west of Sachs Harbour. These provided hourly position data from April 9 onward through the summer period. To help understand the nature of ice movement, instruments were deployed on and through the ice to measure surface winds and ocean currents. The ocean current profiler measured speed and direction of currents at 2 metre (m) intervals to 60 m depth. Two ice mass balance (IMB) buoys measured ice temperature profiles from April 10 to July 31. Ice temperature has a direct bearing on ice strength and will be used in modeling studies. All instruments deployed transmitted data back to the University of Manitoba at 0.5-1 hour intervals. This dataset is currently being analyzed. EM Induction surveys were conducted on multi-year ice to obtain local ice thickness data. Average ice thickness measured on the multi-year floes ranged between 4 to 7 m. The ice island measured 33 m thick.

The second research group was focused on characterizing the thickness and strength of hummocked multi-year ice. In May 2012, researchers used new equipment to measure the temperature, salinity and strength of a 12 m thick multi-year ice hummock. A total of 10.8 m of ice core was removed to provide information about the temperature and salinity of the ice. The ice was coldest near the top layer of ice (-15.1°C) and increased towards the bottom of the ice (to -1.5°C). The ice salinity ranged from 0.1 to 3.7%. Strength tests were performed with an instrument that is lowered into the borehole in ice. A series of strength tests were conducted in each borehole until the bottom of the ice has been reached. For the first time, researchers provided data about strengths in the bottom portion of a thick, multi-year ice hummock. The measurements confirmed that not all hummocks are the same. Some hummocks are old and fully consolidated, with high strengths throughout their full thickness. Other hummocks may have formed recently, and because they still have seawater-filled cavities, they will be weaker overall. Researchers did not encounter any cavities in the hummock that were sampled in May 2012. Tests showed several weak layers within the ice but, overall, the hummock was strong. Researchers intend to use this equipment to sample several more hummocks in the Beaufort Sea in March and April 2013.

The third research group focused on airborne observations of the distribution, thickness, and drift of different sea ice types and extreme ice features in the Canadian Beaufort Sea. Extensive airborne electromagnetic (EM) ice thickness surveys have been performed in April 2009, 2011, and 2012 over the Canadian Beaufort Sea with a long-range airplane. Results show that the location of the multi-year ice edge can be very variable from year to year. Multi-year ice modal thicknesses ranged between 3.0 and 3.7 m. The seasonal ice zone had very variable ice thicknesses depending on the amount and age of ice formed in coastal polynyas and leads throughout the winter. However, researchers gathered enough data to show that modal first-year ice thicknesses of 2.0 to 2.2 m emerge if profiles are long enough. This can be considered the most representative first-year ice thickness estimate in the Canadian Beaufort Sea in April. However, in the seasonal ice zone also regions with heavily deformed ice thicker than 10 m, and occasional multi-year hummock fields of similar thicknesses occur. Results suggest that multiyear hummock fields may not comprise the thickest ice as they are affected by melt during the summer. Two ice islands had thicknesses between 20 and 30 m. Ice thickness surveys were complemented by the analysis of satellite radar data and tracking of ice features by means of GPS beacons.

Bartlett, Mike

WorleyParsons Canada Services Ltd. Calgary, AB mike.bartlett@worleyparsons.com

File Number: 12 404 793 **Licence No:** 15101

Region: SA Location: Bosworth Creek

Bosworth Creek surface water monitoring program

No summary was submitted for this licence. This project is not in compliance with licencing requirements.

Bierwirth, Eike

University of Leipzig Leipzig, SN Germany eike.bierwirth@uni-leipzig.de

File Number: 12 404 780 **Licence No:** 15045

Region: IN Location: Canadian Beaufort Sea

Study on the vertical distribution of ice in arctic clouds (VERDI)

The purpose of this research was to improve the understanding of the cloud-related processes in the Arctic atmosphere and to use these measurements to improve the performance of regional and global climate models of the arctic. This research was designed to obtain a snapshot of ice thickness in this region of the arctic and create an inventory of arctic sea ice volume. Fieldwork was successfully conducted out of Inuvik between 20 April and 20 May, 2012. A total of 13 research flights were undertaken. Most of the flights targeted clouds over the sea ice of the southern Beaufort Sea. No wildlife was seen on the ice. Researchers observed low, stratiform clouds. Diameters of sampled cloud droplets were typically 20 micrometre at cloud top and 10 micrometre at cloud bottom. Ice crystals were also observed in the clouds in low concentration. A cloud with snowfall was sampled at multiple altitudes and is a useful example to validate modelling of arctic precipitation. The cloud reflectivity was measured to support algorithms for arctic cloud

detection from satellites. Several scientific publications are currently in preparation to shed more light of the role of clouds in the energy budget of the arctic atmosphere.

Bishop, Nicole

Rescan Environmental Services Ltd. Yellowknife, NT nbishop@rescan.com

File Number: 12 404 752 Licence No: 15071 (Multi-year licence)

Region: DC, NS, SS **Location:** Within the Seabridge Gold Inc. Courageous

Lake lease area

Courageous Lake Project

Data collection at Courageous Lake took place for several technical disciplines between May and September 2012. Water quality was sampled at 27 lakes and 12 streams up to three times over the field season. Some of these lakes were also sampled in March for water quality and in August, for sediment quality, primary and secondary producer communities. Soil and vegetation surveys were completed to verify and confirm existing vegetation and terrain mapping based on imagery. Vegetation and soil sampling was also done to collect baseline metal concentration distribution and patterns. Soil surveys were completed to assess the salvage suitability of the soil. Fish communities and fish habitat were sampled within the regional study area of the Courageous Lake Project. Sampling methods to assess community composition included: minnow trapping, gillnetting, and electrofishing. Sensitive habitat inventory mapping and littoral zone habitat mapping were used to assess fish habitat. Two fish fences were installed on Matthews Creek to assess arctic grayling and longnose sucker use of the creek for spawning. Most water bodies sampled were classified as non-fish bearing, and those water bodies that contained fish had communities comprised of arctic grayling, longnose sucker, lake trout and northern pike. During the 2012 open water field season, the hydrology group monitored eight stream discharge locations in order to develop stage-discharge relationships. Five lakes were monitored for stage height changes across the same time period. Bathymetric mapping surveys were conducted in five small lakes in the local study area as well. In June 2012, the meteorological station had the winter precipitation adapter removed and a diagnostic test performed on the current set of sensors on the station tower. In anticipation of the winter period, the power system for the station was updated and the winter precipitation adapter was reinstalled in September 2012.

Bottenheim, JanEnvironment Canada

Toronto, ON jan.bottenheim@ec.gc.ca

File Number: 12 404 729 Licence No: 15111 (Multi-year licence)

Region: IN Location: Arctic Ocean

O-buoy measurements of ozone, carbon dioxide and bromine oxide over frozen surface of Hudson Bay and Arctic Ocean

The purpose of this research was to take part in the establishment of long-term observation network to measure concentrations of ozone, carbon dioxide and bromine oxide. An autonomous ice tethered platform known as "O-buoy 4" started measuring the concentrations of ozone, carbon dioxide and bromine monoxide in the air over the ice of the Arctic Ocean (88.15°N and 157.49°W) on September 5, 2011. The deployment of the instrumentation package was conducted by the CCGS Louis S. St-Laurent Icebreaker. O-buoy 4 drifted with the ice in the wind driven-ice circulation known as Transpolar Drift Stream. This circulation moves ice from the Siberian Coast

of Russia across the arctic and exits predominantly into the North Atlantic through the Fram Strait on the east coast of Greenland. The O-buoy 4 was successfully recovered on August 25, 2012. It collected data on air composition, meteorological variables, the ice drift and the ice conditions through its 355-day journey across the High Arctic. It was the first O-buoy that recorded ozone and carbon dioxide concentrations over the ice of the Arctic Ocean with the first parallel measurements of bromine monoxide. It was also collected a complete suite of the surrounding environmental parameters throughout all four seasons in the High Arctic, providing details on regional and seasonal variability. It recorded carbon dioxide variations through the ocean freezing and the ice melt periods and reported data for the Marine Weather forecasting. Collected data are to be used for wind direction and wind speed validation of the forecast models, as well as by scientists to better understand the atmospheric processes in the Arctic. This will help develop advance forecasting capacity for the future developments and scientific understanding of the climate. Daily updated information was posted on open to the public web site: http://obuoy.datatransport.org/monitor.

Budziak, Jerry

Seaway Energy Services Inc. Calgary, AB jerry.budziak@seawayenergy.com

File Number: 12 402 475

Region: SA

Licence No: 15002 (Multi-year licence)

Location: The Nota Creek C-17 wellsite

Phytoremediation study on the CDN Forest et al Nota Creek C-17 Wellsite

Phytoremediation is a remediation strategy involving the use of plants to remove contaminants. In theory, plants uptake the contaminant from the soil, are harvested and then removed from the site. This process is repeated until the impacted soil is remediated to applicable guidelines. Phytoremediation activities on the Nota Creek C-17 wellsite progressed to full site planting in 2009 and 2010. Remediation results were encouraging enough to support excavating impacted soil still buried on the site and integrating it into the phytoremediation process. Soil laboratory results from fall 2011 indicated that the impacted soil still required further phytoremediation. Over the course of the 2012 summer, a full phytoremediation cycle of planting, growth and harvesting was undertaken. Laboratory results from the collected samples indicate further soil remediation progress and support continuing with the application of phytoremediation technology to the wellsite.

Burgess, David

Natural Resources Canada Ottawa, ON david.burgess@nrcan.gc.ca

File Number: 12 404 707 Licence No: 15008 (Multi-year licence)
Region: IN Location: South Melville Ice Cap

Melville Island South Ice Cap mass balance and snow pollution

The objective of the multi-year project is to measure the changing volume of the South Melville Ice Cap using an automatic weather station and sampling techniques. Measurements of snow accumulation and ice melt were performed at 21 pole locations on the ice cap by researchers on April 16 and 17, 2012. The South Melville Ice Cap is a small plateau ice cap (76 km² in size) that is located on the western portion of Melville Island. Over half of the mass balance poles planted in the Melville ice cap were bent in a southwesterly direction. Bending of these poles was most likely caused by heavy loading of rime (frozen mist) and subsequent strong wind events that

occurred in the late fall of 2011. Pole measurements indicate that the ice cap has thinned by ~134 cm as a result of warm temperatures during the summer of 2011 (this equates to a loss of 0.1 km³ [or 100,000 metric tonnes] of water to the ocean over the past year). These melt rates are the highest since records began in 1963, surpassing the previous record in 2007 by ~15 cm water equivalent mass loss. Temperature data downloaded from the automatic weather station on the ice cap indicate that the 2011 summer melt season extended from early June to early September with an average summer temperature of +3.5°C. Continued monitoring of the Melville South Ice Cap is important as its rapid changes are providing valuable insight into long-term climate change over the western Canadian arctic region. The accelerating melt trends observed from this ice cap are consistent with those from the other glacier monitoring sites located across the Canadian high arctic. Ongoing measurements are essential for improved estimates to global sea-level rise and documenting climate change across this region. The only wildlife sightings included an Arctic fox that came within 20 meters of the Melville Hut.

Burn. Chris

Carleton University
Ottawa, ON
christopher_burn@carleton.ca

File Number: 12 404 325 Licence No: 15040 (Multi-year licence)

Region: IN Location: Garry Island; Illisarvik, Inuvik, near the

Dempster highway; Paulatuk; Red Lake; Bar C; Seal Lake;

Dennis Lagoon

Permafrost and climate change, western arctic Canada

The objective of this ongoing project is to understand how climate change is affecting permafrost in the western arctic, particularly in the outer Mackenzie Delta. In 2012, investigations continued on ground ice and near-surface permafrost at Herschel Island. Samples were collected for analysis from the alluvial fan at Pauline Cove and the surrounding hills. Researchers continued to collect ground temperatures near Inuvik and at both Garry Island and Illisarvik on Richards Island. A new investigation of the carbon content of permafrost at Illisarvik began this year. Overall, researchers continued to monitor ground temperatures in the region to see how they are warming up as the climate changes. Last May, a multi-disciplinary book on Herschel Island based this long-term research project was published by the Wildlife Management Advisory Committee (North Slope). The publication was sponsored by several Inuvialuit agencies, the Aurora Research Institute and several Yukon agencies. Copies of the book are available from the researchers.

Coulton, Daniel

Golder Associates Ltd. Yellowknife, NT daniel_coulton@golder.com

File Number: 12 404 763 Licence No: 15027 (Multi-year licence)

Region: NS **Location:** Fortune Mineral's NICO property; along the route of a proposed all-weather access road from the

proposed Tłycho Road

Environmental baseline surveys of the Fortune Minerals Ltd. NICO Project

A plankton baseline field program was completed during the 2012 open water season (July to September) at the Fortune Minerals Limited – NICO project. The program was designed to collect additional plankton data in response to a request made during the February 2012 technical

meetings. Phytoplankton, chlorophyll a and zooplankton samples were collected at five stations within Nico Lake, Peanut Lake, Burke Lake and Reference Lake, as well as from three stations within Little Grid Pond. Due to logistical issues, Reference Lake was only sampled in August and September. Samples have been submitted for analysis and results are pending. The plankton dataset will be summarized in a baseline report, which will be submitted to the Wek'èezhìi Land and Water Board. This report will be available on the Public Registry. Additional water quality sampling was completed at eight stations in the ephemeral stream between Little Grid Pond and Nico Lake. Samples were submitted for analysis and results are pending. Leachate samples from the on-site waste rock and/or tailings field cells were collected on a monthly basis between May and October, 2012. These samples were submitted for chemical analysis and will be used as additional information in the project design.

Dahl, Mark

Environment Canada Winnipeg, MB mark.dahl@ec.gcca

File Number: 12 404 788 **Licence No:** 15088

Region: IN **Location:** Within a 5 km radius of Sachs Harbour

Sachs Harbour disposal at sea follow-up study

No research was conducted under this licence in 2012. The project cancelled due to inclement weather and unsuitable ice conditions.

Dallimore, Scott

Geological Survey of Canada Sidney, BC sdallimo@nrcan.gc.ca

File Number: 12 404 359 Licence No: 15004 (Multi-year licence)

Region: IN, GW **Location:** Outer Mackenzie Delta in Camp Farewell and Mallik areas; Richards Island, Tuktoyaktuk Peninsula

Mackenzie Delta shallow gas and permafrost studies

No research was conducted under this licence in 2012.

Derksen. Chris

Environment Canada Toronto, ON chris.derksen@ec.gc.ca

File Number: 12 404 641 Licence No: 15169 (Multi-year licence)

Region: IN Location: Trail Valley Creek (40km north of Inuvik)

Airborne SAR and passive microwave measurements over snow covered tundra for CoReH20 retrieval validation and land surface model testing

The objective of this ongoing research project is to describe the distribution and physical properties of snow cover in the Trail Valley Creek watershed. This will be used to assess estimates of snow cover properties obtained from a distributed hydrological model, and airborne radar remote sensing. The proposed European Space Agency Earth Explorer 7 CoReH20 satellite mission has the primary objective of retrieving snow water equivalent (SWE) over land at a high spatial resolution (200-500 metres). This would make high resolution, satellite SWE measurement

available for monitoring and modeling applications. The Trail Valley Creek project is a test project for: (1) the CoReH20 satellite SWE retrieval approach; (2) the potential use of CoReH20 SWE retrievals as observational inputs to various environmental prediction models (e.g. land surface data assimilation systems and distributed hydrological models used at Environment Canada). During December 2012, the airborne radar instrument did not function properly, and so usable measurements were not collected. Regardless of these issues, a comprehensive dataset of snow properties was collected during a time period when field observations are typically not made in the subarctic or arctic due to the limited light and cold conditions. Long transects (~15 km total ground distance covered) of snow depth, bulk density, and SWE were collected along predetermined sampling lines. Detailed observations of snowpack stratigraphy (layering, grain type and size, density) were collected at twelve sites (selected to include the variability in snow properties). Snow grain specific surface area measurements were also acquired at these sites. Ground based LiDAR surveys of three snow drift sites were performed in order to estimate the volume and mass of snow storage in these features. Collectively, these measurements represent an important baseline from which the snow measurements in March and April will be compared. There are three phases for the data collection, further measurements are planned for March and April 2013.

Desrosiers, Sarah

University of British Columbia Vancouver, BC desrosie@gmail.com

File Number: 12 404 794 **Licence No:** 15113 (Multi-year licence)

Region: NS **Location:** Daring Lake

Impacts from climate change on berry productivity in the Canadian Arctic: Integrating community participation with science

The objective of this research was to establish a long-term community-based monitoring program in arctic communities using culturally important berry species as indicators of climate change. Sample were collected from the berry-monitoring site (established in 2008) from July 23 to August 5, 2012 in Daring Lake. Changes in berry productivity (weight, abundance and ripeness levels) were record and compared with past data. The influence of environmental change on berry productivity may have serious implications for wildlife that depend on berry crops for survival and for humans who view berries as culturally important. Data collected from this site will also be compared with berry data collected from sites located around Kugluktuk, Nunavut. Cranberries and blueberries were also harvested from around the experimental research valley to analyze berry productivity under certain environmental conditions. Samples are still being analyzed.

Duffe, Jason

Environment Canada -- National Wildlife Research Centre Ottawa, ON jason.duffe@ec.gc.ca

File Number: 12 404 743 **Licence No:** 15081 (Multi-year licence)

Region: IN **Location:** The coastline of the Beaufort Sea, from the

Alaska/Yukon border to the Northwest Territories/Nunavut border. Ground-based measurements using remote sensing were collected at: eastern Ivvavik National Park; Herschel Island; Northern Richards Island; north of

Tuktoyaktuk; Anderson River Delta

Assessing the potential for Environmental Sensitivity Index mapping in the arctic using Synthetic Aperture Radar

The objective of this project was to create an Environmental Sensitivity Index map using helicopter videography. In late July 2012, researchers collected geo-tagged video and audio commentary of approximately 2500 km of Northwest Territories coastline around Banks Island and the east and middle channels of the Mackenzie River Delta (north of Inuvik). Radar (RADARSAT-2 and TerraSAR-X) and optical (SPOT) data are being analyzed to determine the most suitable techniques and optimal datasets to differentiate shoreline types (e.g. intertidal and supratidal zones). Textural measures and polarimetric parameters will also be generated and used with supporting datasets (e.g. bathymetric/elevation data and slope grids) to extract the shoreline types. In addition, satellite imagery and ground measurements (GPS points, ground photos, spectrometer and chlorophyll measurements) were acquired over and in the Aulavik National Park, Mackenzie Delta and Herschel Island study sites. Comparisons between traditional mapping approaches (manual interpretation of shoreline videos) and remote sensing techniques are being performed to verify if the satellite products are as reliable as traditional approaches. Data from the ground campaign are currently being analyzed and will be supplemented by further measurements and image acquisitions next summer. The shoreline videos are currently being interpreted and input into a Beaufort Coastal Sensitivity Atlas.

England, John University of Alberta Edmonton, AB john.england@ualberta.ca

File Number: 12 404 141 **Licence No:** 15082

Region: IN **Location:** Duck Hawk Bluffs, southwest Banks Island;

Bernard Island

Environmental change at Duck Hawk Bluffs, SW Banks Island: From a forested to glaciated arctic

The purpose of this research was to re-log the stratigraphic units that comprise the length of Duck Hawk Bluffs, southwest Banks Island. Research was conducted at the bluffs from June 20 to July 30, 2012. The site includes 12 kilometers of coastal cliffs reaching 50 meters in height and extending 12 km from Mary Sachs Creek to Kellett Point, at the southwest tip of the island. The bluffs provide an important source of sand and gravel for the airstrip and local roads around Sachs Harbour and are well known for their fossil tree stumps. The most important new conclusion from this research is that the lowermost unit of sand and gravel (called the Mary Sachs gravel) were not deposited by rivers before the first arrival of glaciers, but rather came from the glacier that advanced from the mainland, that also picked up the tree fragments along its way there. Above the sand and gravel, two-thirds the way up to the top of the bluffs, is a beautiful, buried tundra surface from an ancient time that was similar to today. The buried tundra has fossil tundra polygons and well-preserved mosses that grew around ponds, more than 800,000 years ago. Above the buried tundra are later deposits from at least two younger glaciations that extend to the top of the bluffs. The last glaciation dates from about 25,000 to 14,000 years ago when Banks Island became ice-free like today. These studies show the importance of Banks Island is to understanding how the Arctic has changed.

English, Michael

Wilfrid Laurier University Waterloo, ON menglish@wlu.ca File Number: 12 404 555 Licence No: 15016

Region: NS Location: Around Wekweeti in the boreal forest and east-

north-east into the tundra towards the Coppermine River

Evolution of the snowpack and snowmelt chemistry in the boreal forest and tundra ecosystem

The objective of the snow survey portion of this research has been accomplished within the constraints of a one year field study. With the existing field data it is possible to compare the pattern of snow accumulation over the snowyear between the data generated by the Environment Canada algorithm using the special sensor microwave/imager satellite data and the field data. From these data it has been demonstrated that within the constraints of this one field season, that utilizing the algorithm approach solely would lead to overestimations of Snow Water Equivalent (SWE) on the Earth surface in this subarctic region by approximately 24%. The researchers' data indicate that this error decreases as the snowpack increases in SWE. The water chemistry portion of the research has yet to be achieved. The sampling procedures are in place for extracting surface water samples from the large catchment draining into Snare Lake once snowmelt starts. This data will be added to the report once the sample collection and chemical analysis is complete. The data recorded in a single year of this study provides the researchers with an indication of the relationship between the algorithm produced SWE values and those determined by fieldwork during the snowyear. However, given snowpack variability from year to year, this relationship may change from low to high snowfall years. Adding to this data base by continuing this type of study would provide that level of understanding. Competence in using satellite data to accurately discern patterns of SWE would help biologists interested in understanding the role of the snowpack in the biology and ecology of ungulates and small mammals. As gathering snowpack data is very expensive having a remote sensing tool to do this over large areas with confidence would benefit not only biologists but those government officials charged with determining the probability of forest fires (areas with lower winter snowfall) and those managing hydroelectric reservoirs who are very interested in annual recharge to those reservoirs from the snowpack.

Fortier, Martin

ArcticNet -- University Laval Quebec, QB

martin.fortier@arcticnet.ulaval.ca

File Number: 12 404 652 **Licence No:** 15070 (Multi-year licence)

Location: The Beaufort Sea; Mackenzie Shelf; Amundsen Region: IN

Gulf region

ArcticNet: an integrated regional impact study of the Coastal Western Canadian Arctic.

In 2011, the ArcticNet Network of Centres of Excellence of Canada and IMG-Golder Corporation established two Marine Observatories under the Beaufort Regional Environmental Assessment (BREA) framework. The first observatory consists of one oceanographic mooring located north of the Mackenzie Trough, at the western limit of the Canadian Beaufort Sea and the second observatory consists of three moorings located roughly 100 nautical miles northwest of Tuktoyaktuk. The purpose of the observatories is to obtain continuous sea ice, ocean circulation and biogeochemical flux data in the Beaufort Sea over a four-year period. This results are likely to contribute to a better understanding of the different ecosystem processes operating in this region of primary importance for potential future oil and gas exploration. In 2012, mooring operations in the Beaufort Sea/Mackenzie Shelf/Amundsen Gulf region were carried out from the Canadian Coast Guard icebreaker CCGS Sir Wilfrid Laurier from September 18 to October 5. During the 17 days at sea, the four moorings deployed in 2011 were serviced and redeployed. A

new mooring was also deployed further north of the Mackenzie Trough, in deeper water. Operations at sea also included CTD-Rosette deployments to obtain the physical-chemical profile of the water column at each of the mooring sites. The two Marine Observatories address monitoring needs for oceanic and sea-ice conditions in the Beaufort Sea, providing essential information for the corroboration of ocean circulation models, oil spill trajectory models and for future environmental (biophysical) assessments at local and regional scales.

Grogan, Paul

Queen's University Kingston, ON groganp@queensu.ca

File Number: 12 404 687 Licence No: 15073 (Multi-year licence)

Region: GW, NS Location: Daring Lake Terrestrial Ecosystem Research

Station

Controls on carbon and nutrient cycling in arctic tundra

The objective of this ongoing research is to substantially advance the understanding of how Canadian arctic tundra ecosystems function, and therefore how it are likely to be affected by perturbations such as climate change, resource development and extraction, and atmospheric pollution. In 2012, researchers worked at Daring Lake with collecting soil and gas flux samples from the snow fence experiment. The samples are currently being analyzed to determine the potential for legacy effects of deeper snow on carbon release and soil nutrient mobilization. In 2012, results were published that demonstrate birch shrub apical growth at the Daring Lake research site is limited as much by the availability of phosphorus as it is by nitrogen.

Guest, Bernard

University of Calgary Calgary, AB bguest@ucalgary.ca

File Number: 12 404 790 **Licence No:** 15094

Region: IN, GW **Location:** Areas north and south of the town of Inuvik

including the Caribou Creek quarry

Thermochronologic approaches for fundamental energy exploration (TAFFEE)

No research was conducted under this licence in 2012.

Haas, Claudia

Environment and Natural Resources -- Government of the Northwest Territories Yellowknife, NT claudia haas@gov.nt.ca

File Number: 12 404 774 **Licence No:** 15000 (Multi-vear licence)

Region: DC Location: Ekali Lake (Ezáa Łue Túe); Sanguez Lake

(Tł'onie Túé); Gargan Lake (Tłįtętįį); Deep Lake (Dechį

Ná?a); McGill Lake (Tthets'éhk'e')

Ecological assessment fieldwork for Łue Túé Sųlái (the Five Fish Lakes) Candidate Cultural Conservation Area

The final phase of the candidate area ecological assessment was conducted in August 2012. McGill and Deep Lakes were sampled and inflow, centre and outflow sites were established for

each lake. Field measurements (temperature, pH, DO, conductivity) and water samples were collected and analyzed by Taiga Labs, Yellowknife. The water quality was good with temperature uniformly warm, pH slightly basic, major ion concentrations moderate, with nutrient and metal concentrations low. Dissolved oxygen was uniformly high even in the hypolimnion (bottom layer at the lake below the thermocline) of Deep Lake. Sonar measurements were taken across the Deep Lake basin and a maximum depth of 37 meters was recorded. Shoreline observations were made next to each lake station and vegetation included: black spruce, tamarack, birch, aspen, willow, alders, dogwoods and rosehips. Emergent vegetation was more common in epilimnetic (above the thermocline) areas of McGill Lake because Deep Lake contours were very steep. Sediment ranged from organic, woody debris to sandy and small rocks in both lakes but limestone cobble was very common along the Deep Lake shoreline. The survey was conducted as part of a community Youth, Culture and Archaeology initiative.

Hansen. Ken

Husky Oil Operations Ltd. Calgary, AB ken.hansen@huskyenergy.com

File Number: 12 404 797 Licence No: 15118 (Multi-year licence)

Region: SA **Location:** 39 surface water bodies within (and near) the EL462 and EL463 parcels; Bog Creek; Slater River; Little

Bear River

EL462 and EL463 - Regional hydrology and surface water quality sampling

The objectives of this project were to establish baseline water quality, water levels and flow conditions and to evaluate potential changes in water quality and quantity before and after the start of site operations. The regional hydrology survey of EL 462 & EL 463 was conducted between June and October 2012 and included: (1) installing 3 hydrometric stations on Little Bear River, Bogg Creek and Slater River; (2) stream flow measurements on Little Bear River, Bogg Creek and Slater River; and, (3) one surface water quality sampling program of 39 locations across EL 462 and EL 463. The results of the hydrometric stations and stream flow measurements were used to calculate seasonal changes of water flow during the open water season. The surface water quality program provided baseline water quality data prior to additional exploration activities. The hydrology and surface water quality study will continue throughout the open water season of 2013.

Hansen, Ken

Husky Oil Operations Limited Calgary, AB ken.hansen@huskyenergy.com

File Number: 12 404 797 **Licence No:** 15121

Region: SA **Location:** The west side of the Mackenzie River southeast

of Norman Wells 40 km along the proposed year-round

access route

EL462 and EL463 aggregate and permafrost mapping - geophysical survey and confirmatory drilling

The geophysical survey was conducted in July and August, 2012 and employed ground penetrating radar and ohm mapper surveys. The combination of these two survey methods were successful in identifying areas of permafrost, ice lenses and bedrock that will assist in the location

of road construction materials. Hand augering in areas of interest provided supporting information to validate the interpretation of the geophysical survey responses.

Harris, Katherine Golder Associates Ltd. Yellowknife, NT kharris@golder.com

File Number: 12 404 796 **Licence No:** 15117

Region: NS **Location:** Mouth of Baker Creek; Yellowknife Bay;

Yellowknife River upstream of bridge; Yellowknife River in

the Tartan rapids area; Horseshoe Island Bay

Giant Mine phase 4 environmental effects monitoring

Two fish surveys were conducted for the Giant Mine Phase 4 Environmental Effects Monitoring Program to examine the health of two small-bodied fish species exposed to treated effluent from Giant Mine. Fish were captured from two exposure areas (Baker Creek and Yellowknife Bay) and two reference areas (Yellowknife River and Horseshoe Island Bay). A non-lethal survey focused on ninespine stickleback was conducted in July 2012. Fish were captured using seine nets, assessed for non-lethal health endpoints and released live. A sub-set of ninespine stickleback were lethally sampled for age verification. A lethal survey focused on slimy sculpin was conducted in September 2012. Fish were captured using a backpack electrofisher and assessed for lethal health endpoints. Additional supporting water quality and sediment quality information was collected during both fish surveys. During both fish surveys, any young-of-year and bi-catch captured were measured for length and weight prior to release, with the exception of nine burbot and 54 slimy sculpin. These additional fish were sacrificed and archived for potential tissue analysis. All samples have been submitted and results are pending. A final report will be submitted to Environment Canada and will be available through the Giant Mine Remediation Project Public Registry.

Haugaard, Rasmus

University of Alberta Edmonton, AB rasmus@ualberta.ca

File Number: 12 404 787 **Licence No:** 15087

Region: NS Location: Russell Lake; Labrish; Point Lake

Petrology and geochemistry of the late Archaean banded iron formation and associated turbidites, western Slave Craton: constraints on palaeoenvironment

The recent discovery of new occurrences of 2.9 - 2.6 billion years old banded iron formations in the Northwest Territories have opened an exceptional opportunity to study ancient ocean chemistry. These banded iron formations (BIF) provide important environmental information regarding these ancient (late Achaean) oceans including: sea-level fluctuations, volcanism, biology, oxygen level and climate. From July 1–14, 2012, fieldwork was conducted at various locations throughout the central Slave Craton, which is north of Yellowknife. Two different field relations were found for the BIF within the Slave Craton: a 2.6 billion year old BIF associated with shales and turbidites, and a 2.85 billion year old BIF associated with chromium rich quartzites. Both types were documented and samples were obtained. These two different setting may indicate that these rock formations were deposited in two different types of ocean waters. Furthermore, a three-day visit to a drill camp in the eastern Slave Craton were carried out and very pristine core samples were selected and shipped home. Analysis of the samples is ongoing.

Few preliminary geochemical results from Dwyer and Bell Lake show very interesting chromium signatures. Chromium is very important in the enzymes of oceanic microorganism. A high content of graphite were also found within some of the BIF sequences, which could indicate organic production within the water column.

Hicks, Faye

University of Alberta Edmonton, AB faye.hicks@ualberta.ca

File Number: 12 404 493 Licence No: 15023 (Multi-year licence)

Region: SS Location: Along the Hay River from the NWT/Alberta

Border to Great Slave Lake

Hay River ice jam study

Because of limited research funding, only a modest field campaign was conducted in 2012. The University of Alberta field team came to the Town of Hay River just prior to breakup (April 14 to 18, 2012) and placed time-lapse cameras and water level data-loggers along the river upstream of Alexandra Falls. These instruments supplemented those already deployed by the Town of Hay River Flood Watch Committee. Several ice jam release waves were documented with the remote instruments. Operational testing of the ice jam flood forecasting models continued during breakup 2012. The timing of the onset of breakup, the expected peak snowmelt runoff streamflow and the time of arrival of the ice runs from were all predicted with reasonable accuracy. Photos documenting the 2012 this public web site: breakup are posted http://www.riverice.ualberta.ca/breakup/hayriver/pub/HR%20photos%202012.html Additional information about Hay River breakup and this research can be viewed at this site: http://www.riverice.ualberta.ca/breakup/hayriver/pub/Hay River Breakup Study.htm

Holmes, R. Max

Woods Hole Research Center Falmouth, MA United States rmholmes@whrc.org

File Number: 12 404 713 **Licence No:** 15032 (Multi-year licence)

Region: IN, GW **Location:** The Mackenzie River near the Tsiigehtchic ferry

crossing

The arctic great rivers observatory

This five-year project studies the six largest rivers that flow into the Arctic Ocean (in North America the Mackenzie and Yukon Rivers, and in Russia the Ob', Yenisey, Lena, and Kolyma Rivers). In the first year (2012), researchers measured the concentration of naturally occurring chemicals (e.g. carbon, nitrogen, and phosphorus) in these rivers, to obtain baseline information about the flow of these chemicals to the ocean. Sampling began in May of 2012 and was conducted every second month. Sampling was conducted from a motorized boat, just upstream of the Tsiigehtchic ferry crossing. For each sampling trip, 8 litres of water were collected, and processed in the Inuvik labs. Researchers also used a hand-held water meter to measure water temperature, pH, conductivity, and dissolved oxygen concentration. Laboratory analysis is underway. These measurement will help show how climate change is impacting Arctic rivers. All data from this project are posted on a public website (http://arcticgreatrivers.org) and are available for free download by the public.

Hood, Alexandra

De Beers Canada Inc. Yellowknife, NT alexandra.hood@ca.debeersgroup.com

File Number: 12 404 728 **Licence No:** 15025 (Multi-year licence)

Region: NS, SS Location: Snap Lake

De Beers Snap Lake Mine - 2011-2014 environmental monitoring program

In 2012, De Beers undertook a number of studies to meet requirements from the Environmental Agreement, Water License and Land Use permit, including studies on aquatics (specifically, water, benthos, plankton, sediment and fish sampling), air (specifically dioxins and furans, NO₂, SO₂ and dust sampling), vegetation, and Wildlife (caribou, grizzly and black bears, wolverine and raptors observation). These studies were carried out as per all license and permit requirements and all reports were submitted to the appropriate parties for review and comment.

Hoos, Rick

EBA Engineering Consultants Ltd. Vancouver, BC rhoos@eba.ca

File Number: 12 404 677 **Licence No:** 15090

Region: DC **Location:** 5 locations along the Flat River at Tungsten

Aquatic environmental effects monitoring - Flat River at Cantung Mine

EBA conducted aquatic sampling on the Flat River from August 30 to September 4, 2012. A total of 512 slimy sculpin, 8 bull trout and 4 Arctic grayling were captured and all but 25 sculpin (sacrificed for tissue metals analysis) and two bull trout (mortalities) were released alive. The Third Interpretive Report for EEM Studies will be submitted to Environment Canada in early March 2013.

Kanigan, Julian

Indian and Northern Affairs Canada Yellowknife, NT julian.kanigan@inac.gc.ca

File Number: 12 404 661 Licence No: 15015 (Multi-year licence)

Region: DC **Location:** Sampling sites will be located on historic seismic lines near the Mackenzie Highway (between 61-

62°N and 120°30'-121°30'W); Scotty Creek Research

Basin

Investigating the effects of winter overland travel in sub-Arctic Boreal Forest

No research was conducted under this research licence in 2012.

Kerr, Jeremy University of Ottawa Ottawa, ON jkerr@uottawa.ca

 Region: IN, GW **Location:** Roadsides along portions of the Dempster

Highway just outside the road bounds of Inuvik and

Tuktoyaktuk

Canadian global change transect: Northern transect No research was conducted under this licence in 2012.

Kershaw, G. Peter University of Alberta Edmonton, AB peter.kershaw@ualberta.ca

File Number: 12 404 116 Licence No: 15089 (Multi-year licence)

Region: SA **Location:** Along the Canol Heritage Trail between mile post 55 and Macmillan Pass on the Yukon border

Long-term ecological and geomorphological investigations in the alpine tundra of the Mackenzie Mountains, NWT

This ongoing research aims to: (1) determine the status of permafrost landforms; (2) determine long-term recovery after abandonment of the Canol No. 1 project; and (3) determine the status of treeline. In August 2012, four automated microclimate stations were serviced and data retrieved. The station damaged by bears in 2010 was returned and reinstalled after repairs. Thaw depth measurements were completed on the nine features at the eight monitoring sites. Permafrost warming is in the order of 0.75 to 1.25°C despite atmospheric cooling in the past few years. Permafrost landforms continue to shrink in area at a rate of ~1% each year. Lack of change in thaw depth on the top of permafrost features confirms they are shrinking from their edges. A survey was conducted on disturbances originating from the CANOL Project (mainly the road right-of-way) to locate colonizing coniferous trees (spruce and subalpine fir). Over 100 were found and most appear to have started growth within the past 30 years (30.78 SD±11.93). A resurvey of CANOL crude-oil spills in tundra environments was also completed. Previous botanical surveys were conducted in 1977-9 and 1997-8. Comparisons of species composition, cover and frequency of occurrence will be included in the analysis.

Kjarsgaard, Bruce

Geological Survey of Canada Ottawa, ON bkjarsga@nrcan.gc.ca

File Number: 12 404 455 **Licence No:** 15061

Region: SS **Location:** The area from 62° 30' to 63° 30'N 106° 15' to 104° 30'W (approximately 270 km northeast of Łutsel K'e)

Heavy mineral indicator tracing in glaciated terrains

The objective of this study was to improve understanding of transportation and deposition of heavy minerals in surficial materials (till and esker systems). 181 samples surficial sediment samples were collected in 2012, of which 76 were from eskers and 105 were from tills (diamicton). A till sample was taken approximately every 100 square km, based on a 10 km x 10 km grid. An esker sample was taken approximately every 10 km along selected/major esker ridge crests. Geochemical results are published in the Geological Survey of Canada (GSC) Open File 7351 with additional portable x-ray fluorescence spectrometry results published in GSC Open File 7408.

Kokelj, Steven

Aboriginal Affairs and Northern Development Canada -- Renewable Resources and Environment Directorate Yellowknife, NT steve.kokelj@inac.gc.ca

File Number: 12 404 545 Licence No: 15052 (Multi-year licence)

Region: IN, GW **Location:** Permafrost Monitoring: Mackenzie Delta
Active layer freezeback: Harry Channel; Fish Island

Shrub removal: Taglu Island

Contaminant movement: Navy Road Quarry; Tundra Lakes

Stream crossings: Ya Ya Crk, Holmes Creek; Jimmy

Creek; Cabin Creek

Environmental studies across treeline

This research investigated permafrost conditions across treeline between Inuvik and the Beaufort Sea coast. The most recent activities included evaluation of ground temperature and environmental conditions at several stream crossings from Inuvik to the coast. Preliminary results show that valley bottoms with tall-shrub riparian vegetation have much higher ground temperatures than those measured on the adjacent tundra hill tops due to wet soils and thick snow cover in the valley-bottoms. Tall shrubs were also removed from a drilling-mud sump near Taglu Island in the outer Mackenzie Delta to study the impacts of shrub removal on ground temperatures. Winter ground temperatures in the top 2 meters of the sump cover decreased by several degrees because shrub removal led to a reduction in winter snow cover. The thinner snow promoted ground cooling in winter. The results suggest that shrub removal may be a useful management technique to maintain permafrost in the sump cap.

Kokeli, Steve

Aboriginal Affairs and Northern Development Canada -- Renewable Resources and Environment Directorate Yellowknife, NT kokeljsv@inac.gc.ca

File Number: 12 404 545

Region: GW

Licence No: 15072 (Multi-year licence)

Location: Dempster Highway corridor

Evaluating the environmental impacts of permafrost mega-disturbances along the Dempster Highway, NWT

In 2012, the field team collected water, permafrost and benthic samples from streams impacted by thaw slumps. The field team also collected ground temperature data to determine permafrost conditions and downloaded cameras that were tracking the growth of large thaw slumps along the Dempster Highway corridor. Lake sediment cores were also collected from Husky Lake. The analysis indicates that large thaw slumps have a major impact on both the water quality and ecology of streams draining the Peel Plateau. In summer 2012, a high intensity rainfall event of 95 mm was recorded at a remote weather station on the Peel Plateau. This extreme event accelerated the movement of debris flows downslope of thaw slumps and caused significant erosion along stream channels. The impacts of extreme events continue to be investigated. Mapping of thaw slumps also continues and air photo and remote sensing techniques are being used to map the extent of disturbances and to identify impacted watersheds across the Peel Plateau.

Krizan, Julia IMG-Golder Corporation Inuvik, NT jkrizan@golder.com

File Number: 12 404 803 **Licence No:** 15136

Region: IN **Location:** In the corridor for the proposed Inuvik to

Tuktoyaktuk Highway between kilometre 105 and kilometre

120

Lake bathymetry survey for the Inuvik to Tuktoyaktuk Highway, NWT

On behalf of the Government of the Northwest Territories Department of Transportation, IMG-Golder completed a lake bathymetry survey for lakes suitable to be used as water sources for winter access road construction between km 105 and km 120 of the proposed Inuvik to Tuktoyaktuk Highway. The objectives of the fieldwork were to identify potentially suitable lakes. collect bathymetry data, estimate water volumes and allowable water extraction volumes for each lake. A field investigation was completed in August 2012 over a three-day period. A crew of two specialists and one Inuvialuit Wildlife Monitor accessed the lakes daily from Inuvik via helicopter and a 14-foot aluminum boat to survey the lakes. Four potentially suitable lakes were identified in proximity to the proposed highway and subsequently surveyed using continuous depth recordings, which were geo-referenced with a Global Positioning System (GPS) inside the bathymetry recorder. The bathymetry data from these lakes were analyzed to estimate the lake profiles, and from that total lake water volumes were calculated. In a final step, a two metre thick ice cover was subtracted from the total water volume to allow for winter conditions. From the remaining available winter water volume, the allowable 10% water withdrawal amount was calculated as per DFO Protocol for Winter Water Withdrawal from Ice-covered Waterbodies in the Northwest Territories and Nunavut.

Lacelle, Denis University of Ottawa Ottawa, ON dlacelle@uottawa.ca

File Number: 12 404 782 Licence No: 15060 (Multi-year licence)

Region: GW Location: Watersheds of Stony Creek; Vittrekwa Creek

The cumulative impacts of rapid environmental change in the northwestern NWT: Investigating the impacts of mega-slump disturbances on terrestrial and aquatic ecosystems in the lower Peel watershed, NWT

From May 28 to June 18 and August 2 to 17, 2012, fieldwork was undertaken in the Stony Creek watershed to study the impacts of landscape disturbances (specifically retrogressive thaw slumps) on the terrestrial and aquatic ecosystems. Eight shallow permafrost cores (50 to 100 cm depth) were collected above the headwall of the slumps. The cores were divided into 1-2 cm vertical sections and will be used to determine the ice content (volumetric ice content and excess ice), as well as the isotope geochemical composition of shallow permafrost. Four hydrological sondes were installed in streams above and below two thaw slumps to monitor and quantify the water quality (water-level, conductivity and turbidity) of clear tundra streams and streams impacted by slump runoff. At these two sites, researchers performed in-situ measurements of stream velocity, to calculate ionic and sediment fluxes and the contribution of slump runoff to the overall stream flow. A survey of stream water quality in the Stony Creek watershed was performed during both field campaigns. Here, stream water samples were collected at the mouth of Stony

Creek, above and below thaw slumps, and from clear tundra streams. The samples will be analyzed for major dissolved ions and trace metals and suspended sediments. The data will allow assessing the impacts of slumps on streams water quality at various sub-catchment scales and will be reported on a GIS platform.

Lafleur, Peter Trent University Peterborough, ON

plafleur@trentu.ca

File Number: 12 404 621 Licence No: 15001 (Multi-year licence)

Region: NS Location: Near Daring Lake Terrestrial Ecosystem

Research Station (within 2 km)

Exchange of carbon gas fluxes over low arctic tundra

Carbon flux measurements at four arctic tundra sites near Daring Lake, continued in 2012. Instruments were set up in early May and continued operating until late August. The overall objective is to see if the tundra is taking more carbon dioxide out of the atmosphere by plant photosynthesis than it is releasing by respiration. If more is taken up by the tundra than released, the tundra is called a sink for carbon and if the opposite is true, it is a source. Researchers hypothesized that the tundra is a sink, which indeed what the 2012 results seem confirm. In the summer of 2012, researchers focused on tundra vegetation growth and measured the amount of leave on the surface and how the amount of leaves changes over the growing season. Results show that in years when more leaves are present, the tundra is a larger carbon dioxide sink. This research helps to show how arctic tundra will influence the amount of carbon dioxide in the atmosphere and thus how it might influence the climate today and into the future.

Laidlaw, Shawn

Ka'a'gee Tu First Nation Kakisa, NT ktfnenvironmental@gmail.com

File Number: 12 404 795 Licence No: 15115 (Multi-year licence)
Region: SS Location: Tathlina Watershed near Kakisa

Investigating the cumulative effects of environmental change and human activity in the Tathlina watershed

Fieldwork has been completed in the Tathlina watershed as part of the Ka'a'gee Tu First Nation (KTFN)-led Cumulative Impact Monitoring Program project in the region. In March, Carleton University researchers and KTFN personnel traveled to Tathlina Lake to collect core sediment samples to determine historical biotic and abiotic conditions at the lake. The cores were collected successfully and are pending analysis in the laboratory. In March, hydrological sondes were retrieved and water grab samples were collected as part of an ongoing regional water quality monitoring program. Further water samples were collected in late May and hydrological sondes were redeployed. In September, Canadian Aquatic Biomonitoring Network (CABiN) protocol work and invertebrate contaminant testing was completed and additional core sediment samples were collected at Tathlina Lake and throughout the watershed. These sediment samples are aimed to test for signs of historical contaminants in the area and to determine baseline conditions.

Lamoureux, Scott Queen's University

Kingston, ON

scott.lamoureux@queensu.ca

File Number: 12 404 567 Licence No: 15067 (Multi-year licence)

Region: IN Location: Unnamed lake, near Chevalier Bay, Melville

Island

Long term river flow and climate conditions reconstructed from lake sediments

No research was conducted under this licence in 2012 due to logistical constraints (lack of aircraft flight time).

Landry, Francois

Rescan Environmental Services Ltd. Vancouver, BC flandry@rescan.com

File Number: 12 404 767 Licence No: 15019 (Multi-year licence)

Region: SS **Location:** In and around the former Pine Point Mine

Pine Point project (N-204)

The main objective of this study was to continue to collect baseline data to characterize the environmental setting (physical and biological) for the proposed Pine Point Project. Much of the originally proposed work was not completed in 2012. Surface water quality samples were collected from the Buffalo River, Paulette Creek, Twin Creek and Birch Creek. Water quality samples were analyzed for standard parameters including total and dissolved metals. Hydrometric monitoring stations were installed along the Buffalo River, Twin Creek and Paulette Creek to record surface water elevations and discharge measurements over high and low flow periods. Wildlife call surveys were completed for amphibians, owls and waterbirds at the project sites. Several incidental wildlife observations were also noted for raptors, landbirds and mammals (moose, deer and bat). Remote cameras were also installed at several locations to collect data on wildlife habitat use.

Langhorne, Amy

Golder Associates Ltd.
Saskatoon, SK
amy_langhorne@golder.com

File Number: 12 404 733 Licence No: 15024 (Multi-year licence)

Region: SS **Location:** Kennedy Lake watershed and surrounding

watersheds

De Beers - Gahcho Kué environmental monitoring program

The purpose of this ongoing research is to build upon current knowledge of the existing environment around the Gahcho Kué Project site including aerial, aquatic and terrestrial baseline conditions. Studies at the project site in 2012 included collection of baseline meteorological data, hydrology, soil, water/sediment quality, and fish and aquatic resources data. Air temperature, rainfall, wind speed/direction and relative humidity data were collected from the site weather station. Fifteen streams were surveyed for navigability in September, with supplemental assessment to characterize the hydrological regime of the surrounding watershed and quantify annual/seasonal water yields and lake water levels. The water/sediment quality component included collecting physico-chemical profile data and water/sediment samples in five potential reference lakes during winter and summer and in five small lakes adjacent to Kennedy Lake during summer. Water quality parameters sampled included major ions, nutrients, chlorophyll a,

trace metals, and sediment quality parameters (i.e. trace organics and trace metals). Two continuous data logging multi-parameter sondes were used to collect physico-chemical water quality in two lakes during the freshet period. Fisheries work included downstream flow monitoring, fish/plankton sampling and fish tissue collection at two new potential reference lakes, and spring sampling in small lakes in the surrounding watershed. Sampling methods included small-mesh gill netting, angling, baited minnow traps and shoreline electrofishing. Physico-chemical water quality, stream discharge/flow measurements and fish habitat assessments were recorded during sampling.

Lantz, Trevor
University of Victoria
Victoria, BC
tlantz@uvic.ca

File Number: 12 402 712 Licence No: 15123 (Multi-year licence)

Region: IN, GW **Location:** Throughout the Mackenzie Delta; Peel Plateau; 16 sites along the Dempster Highway; 5 Aklavik area sites

A multi-scale assessment of cumulative impacts in the Northern Mackenzie Basin

Since 2010, Aboriginal Affairs and Northern Development Canada scientists have been working with researchers at the University of Victoria, and Hunters and Trappers Committees (HTCs) in the Mackenzie Delta to develop a vegetation and permafrost monitoring protocol that can be implemented by a range of users. The long-term goal of this program is to establish and maintain a network of sites to characterize regional environmental variability, and serve as a baseline against which to measure changes resulting from the cumulative impacts of multiple natural and human-caused disturbances. In 2012, measurements of vegetation structure, plant community composition, tree density, the productivity of edible berries, active layer depth, and near surface ground temperatures were made at all 60 sites across 9 terrain types. At a set of core sites, measurements from meteorological stations, frost tubes, and deep ground temperature cables were also recorded. Over the course of this project, the researchers have worked with a range of participants including: local youth, HTC wildlife monitors, graduate and undergraduate research assistants, and government scientists. Based on simulations of statistical power and preliminary comparisons of disturbed and undisturbed sites they are confident that their protocol can be used to detect temporal changes in variables of ecological relevance. Data on vegetation, active layer, and snow are added to the NWT Discovery Portal on an ongoing basis.

Machtans, Hilary

Golder Associates Ltd. Yellowknife, NT hmachtans@golder.com

File Number: 12 404 799 **Licence No:** 15127

Region: NS Location: Great Slave Lake at: Yellowknife Bay; Jackfish

Bay; Kam Bay; Horseshoe Island Bay

Con Mine Phase 4 EEM - periodic monitoring

Golder Associates Ltd. was contracted by Miramar Northern Mining Ltd. to collect field environmental, fish, and invertebrate data for the Phase 4 Environmental Effects Monitoring (EEM) program for Con Mine as required under federal Metal Mining Effluent Regulations. The field program was conducted in and around Yellowknife Bay of Great Slave Lake, between July and September 2012. Sampling areas included an exposure area (Jackfish Bay) located downstream of the Con Mine outfall. Two reference areas in Great Slave Lake were also sampled:

Horseshoe Island Bay was sampled for fish and Kam Bay was sampled for invertebrates. Ninespine stickleback were captured using seine nets and were processed for lethal health endpoints, as such gonad, liver, and stomach samples were collected. Bi-catch were measured for length and weight prior to release. Invertebrates were sampled in Jackfish Bay and Kam Bay using Hester-Dendy Artificial Substrates and an Ekman Dredge. Additional supporting water quality and sediment quality information was also collected. All samples have been submitted and results are pending. A final interpretative report will be submitted to Environment Canada by June 6, 2013.

MacNaughton, Robert

Geological Survey of Canada Calgary, AB robert.macnaughton@nrcan-rncan.gc.ca

File Number: 12 404 529 **Licence No:** 15105 (Multi-year licence)

Region: SA Location: Mackenzie Mountains near Norman Wells and

Tulít'a

Geological Fieldwork in Mackenzie Plain and adjacent mountains

A team of three scientists from the Geological Survey of Canada undertook geological fieldwork based out of Norman Wells for two weeks in July 2012. The work involved helicopter visits or overland hiking to 170 rock outcrops on ridges and streams from the eastern Mackenzie Mountains to the Franklin Mountains. Locations and rock descriptions were recorded, photographs taken, and rock thicknesses and orientations were measured. Approximately 30 rock samples were collected, varying in size from a fist to slightly larger than a loaf of bread. Samples were shipped to labs at the Geological Survey of Canada in Calgary where they are undergoing paleontological and organic chemistry analyzes. Data are being used to produce new geological reports and maps of bedrock geology for the Norman Wells and Tulít'a region.

Marsh, Philip

National Water Research Institute -- Environment Canada Saskatoon, SK philip.marsh@ec.gc.ca

File Number: 12 404 378 Licence No: 15007 (Multi-year licence)
Region: IN, GW Location: Trail Valley Creek; Havikpak Creek

Hydrological studies, Mackenzie Delta Region

With a changing climate and increasing development there is an urgent need for appropriate hydrological information (e.g. snow cover, soil moisture, soil temperature, stream discharge) in the western Canadian arctic. For example, the design of roads and pipelines requires estimates of maximum stream discharge, while rules controlling land access in the fall require estimates of snow cover and whether the soil is frozen. However, with a changing climate, the recent past may not be a reliable guide to the hydrological conditions in the near future. As a result, in order to limit the environmental impact of development, better methods to predict future conditions are needed. This research program aimed at developing such improved methods. In 2012, researchers: (1) collected hydrologic data at two study sites in order to extend a 20+ year data set; (2) began enhanced studies of snow accumulation in Trail Valley Creek (50 km north of Inuvik) which included extensive and frequent snow surveys, use of a laser terrain scanner to determine volume of snow held in large valley-side drifts, the addition of a new snowfall precipitation gauge in a forested site, added infrastructure at several instrument sites to allow the upcoming installation of new sensors for measuring the amount of water held in snow (not just snow depth), and the

continuation of an experiment monitoring multiple snow depths at shrub and tundra locations; (3) continued development of better methods to predict future changes in snowcover, soil moisture, ground thaw, and streamflow; (4) rejuvenation of main meteorological measurement location in anticipation of the addition of new instrumentation to be installed in April 2013 to monitor fluxes of carbon dioxide and energy in the basin. Recent results consider the factors controlling the thaw of the upper layer of the ground over the summer period. This is an important step towards better predictions of the impact of a changing climate and developments on the hydrology of the region. Other results have considered the role of lakes on the hydrology and ecology of the Mackenzie Delta, combined with the continued analysis of highly detailed maps showing land and water elevations at four large transects across the Mackenzie Delta, with the intent of improving the understanding of arctic river delta water level regimes.

Messmer, David

University of Saskatchewan Saskatoon, SK dave.messmer@ec.gc.ca

File Number: 12 402 874 Licence No: 15066 (Multi-year licence)

Region: GW, NS

Location: The wetlands along the current and old
Yellowknife Highway between the communities of
Yellowknife and Rae; the wetlands along the Dempster

highway between Inuvik and the ferry crossing north of

Tsiigehtchic

Effect of spring and summer temperatures on amphipod reproduction (Gammarus lacustris)

No summary was submitted for this licence. This project is not in compliance with licencing requirements.

Miles, Warner

Geological Survey of Canada, Natural Resources Canada Ottawa, ON wmiles@nrcan.gc.ca

File Number: 12 404 718 Licence No: 15026 (Multi-year licence)

Region: SS Location: South Rae

South Rae, NWT aeromagnetic survey

The objective of this research was to acquire high-resolution aeromagnetic data in an area centred of the South Slave region of Northwest Territories. Aeromagnetic surveys measure magnetic properties of bedrock and are one of the tools used in geological mapping. Understanding these magnetic properties will help geologists map the area, assist mineral exploration activities, and provide useful information necessary for communities, aboriginal associations, and government to make land use decisions. The survey was flown between January 4, 2012 and March 23, 2012. It collected approximately 108,000 line km of data flown along parallel lines spaced 400 m apart. The flying height was at a terrain clearance of about 150 m. The intensity of the total magnetic field was measured from the aircraft. The contractor monitored interactions with large mammal species concentration. Final data were accepted for the survey and preparation of maps for publication is complete. The data and maps were published on September 7, 2012. Copies of all maps were sent to the NWT Métis Nation, the Athabasca Denesuliné, and the Aurora Research Institute. The data are available for free download from the Geoscience Data Repository for Aeromagnetic and Electromagnetic Data

(http://gdr.nrcan.gc.ca/aeromag) and digital versions of the maps are similarly available from MIRAGE (http://gdr.nrcan.gc.ca/mirage). The Open File numbers are: 7120 7121 7122 7123 7124 7125 7126 7127 7128 7129 7130 7131 7132 7133 7134 7135 7136 7137.

Moore. Kristin

Diavik Diamond Mine Inc. Yellowknife, NT krisitn.moore@riotinto.com

File Number: 12 404 766 Licence No: 15048 Region: NS Location: Lac de Gras

Diavik aquatic effects monitoring program 2012

Diavik Diamond Mines Inc. conducts environmental monitoring programs under the terms and conditions of the Territorial Water Licence (W2007L2-0003) issued for the Diavik Diamond Mine and the Fisheries Authorization (SC98001) issued by Fisheries and Oceans Canada. The principal objective of the Aquatic Effects Monitoring Program is to monitor the Mine's water discharge and other potential stressors. Specifically, this ongoing program's goals are: (1) to determine the short and long-term effects in the aquatic environment resulting from the Diavik Diamond Mines; (2) test impact predictions; (3) measure the performance of operations; and (4) evaluate the effectiveness of impact mitigation. In 2012, samples of water chemistry, sediment quality, lake productivity, planktonic and benthic invertebrate communities, dust deposition; fish, fish habitat, and the use of fisheries resources in Lac de Gras were collected. Overall, samples show that nutrients (nitrogen and phosphorus) released from the treated mine water discharge are causing low to moderate enrichment-effect in Lac de Gras. Dust deposition rates in 2012 were consistent with previous years (whereby deposition rates were highest immediately adjacent to the project infrastructure and decreased with distance from the Mine). The analysis of effluent and water chemistry data collected during the field program and from relevant sites from the Water Licence SNP program stations indicated similar trends as observed in 2011, including an increase in arsenic and iron concentrations. Effect levels will be determined during 2013. Results, to date, of the plankton monitoring program - which examines changes in the amount, number and types of tiny animals (zooplankton) and algae (phytoplankton) that live in the water of Lac de Gras indicate a pattern consistent with weak nutrient enrichment from Mine effluent. Higher amounts of phytoplankton (chlorophyll a) and total phosphorus were measured. The observed enrichment effect has been given a "moderate" effect level designation. Zooplankton biomass resulted in a "low" effect level designation.

Narbonne, Guy Queen's University Kingston, ON

narbonne@geol.queensu.ca

File Number: 12 404 783 **Licence No:** 15062

Region: SA Location: Mackenzie Mountains; June Lake (63°31'10"N,

128°38'00"W); 63°20'04"N, 128°19'16"W); 63°15'56"N,

128° 37'19"W)

Behavioural analysis of trace fossils at the Ediacaran-Cambrian boundary

Soft worm-like animals capable of movement appeared suddenly worldwide 555 million years ago. The purpose of this research was to examine fossils of these early animals to study the evolution of their muscles and brains, as well as their communities. Researchers spent one-month taking pictures and collecting samples of fossilized burrows in Sekwi Brook North, Sekwi Brook

South, and Ingta Ridge. The formations that were studied were chosen because they represent the transition from Ediacaran to Cambrian assemblages. It was observed that burrows collected in the Blueflower formation showed simpler and more random horizontal movement patterns, while burrows of organisms collected in the Ingta and Backbone Ranges formations were more diverse, with many burrows displaying more organised search behaviour and/or a vertical component. Some of the younger burrows demonstrate the ability to sense and avoid previous burrows, creating crude spiral and meandering patterns. These patterns become more refined over geological time. Other younger burrowers probed the sediment, either horizontally or vertically. The abilities to backfill ones burrow and to leave scratch marks on the sediment surface were also observed. These observations and the data collected are being used to categorize behavioural changes and correlate them with stratigraphy in order to learn more about the development of brains and muscles across the Ediacaran-Cambrian boundary.

O'Neill, Brendan

Carleton University Ottawa, ON brend.esq@gmail.com

File Number: 12 404 738 Licence No: 15114 (Multi-year licence)

Region: IN, GW **Location:** Peel Plateau - along the Dempster Highway between 67°27'11"N, 134°47'43"W and the Yukon Border

Permafrost conditions and terrain stability considerations along the Dempster Highway, NWT

The Dempster Highway from the Yukon to the Mackenzie Delta area is the only road to the western arctic. The highway crosses several distinct landscapes from the mountains at the Yukon border to the Peel Plain lowlands at Fort McPherson. The objectives of this study were: (1) to examine the influence of drainage hydrology and snow cover on permafrost temperatures in the area, and hence vulnerability to future warming, (2) to assess the geomorphic, climatic and permafrost conditions in the study region and their influence on hillslope stability; and (3) to determine the ground temperature changes next to the highway in order to assess its impacts on permafrost from the roadway. This study is ongoing and will enable an assessment of the controls on permafrost conditions and slope stability in the Peel Plateau region. Though this research is physical in nature, the results will be important to many northerners as they may facilitate management strategies to cope with potential impacts of climate change on this critical transportation route.

Osinski. Gordon

University of Western Ontario London, ON gosinski@uwo.ca

File Number: 12 404 701 Licence No: 15083 (Multi-year licence)

Region: IN Location: The Collinson structure (72°30' N, 114°0'W), a

suspected meteorite impact crater located on the

northwest portion of Victoria Island

Investigation of a possible impact structure on Victoria Island, NWT

Meteorite impact structures represent sites where asteroids or comets have struck the surface of the Earth in the past. They are one of the most common geological landforms in the Solar System. On Earth, approximately 180 impact craters have been documented to date, of which around 30 are in Canada. The goal of this project was to confirm the impact origin of an unusual structure

near Collinson Inlet in northwestern Victoria Island. This site was visited for a two-week period in July 2012. This expedition was successful in confirming the impact origin of this structure through the discovery of shatter cones – distinctive fracture surfaces with a conical shape. While in the field, this team carried out mapping of the structure and collected samples for follow-up laboratory work. This mapping showed that the original impact structure was about 28 km across, making it the largest impact crater to be discovered in recent years. Other findings included evidence for hydrothermal activity – this occurs when ground waters are heated by the impact event and then flow through the fractured rocks, potentially forming hot springs, the likes of which can be found in places like Iceland and Yellowstone National Park in the U.S.A.

Panayi, Damian

Golder Associates Ltd. Yellowknife, NT damian panayi@golder.com

File Number: 12 404 779 Licence No: 15034 (Multi-year licence)

Region: NS **Location:** Bluefish Lake; Prosperous Lake; Yellowknife
River between Prosperous Lake and Bluefish Lake

NTPC Bluefish Hydro repairs

The objective of this study was to describe and monitor the aquatic environment in the Yellowknife River between Prosperous Lake and Bluefish Lake during the construction of a new dam and spillway for the Bluefish Hydro Plant. In 2012, there was regular monitoring of water quality near construction areas to confirm the efficacy of mitigation. Efforts were made to document current fish use of the existing dam, to guide monitoring of fish use of a new spawning shoal to be built as fish habitat compensation. Samples were collected from large and small-bodied fish to assess current methyl-mercury levels. Finally, monitoring was undertaken to document water flows in the Yellowknife River and fish migration up from Prosperous Lake. Results from all monitoring will be presented in annual reports to the Mackenzie Valley Land and Water Board.

Paradis, Suzanne

Geological Survey of Canada Sidney, BC suparadi@nrcan.gc.ca

File Number: 12 404 772 Licence No: 15006 (Multi-year licence)

Region: SA, DC **Location:** Howards Pass deposit; Prairie Creek deposit;

Gayna River deposit

Hydrothermal event recognition and vectoring to SEDEX ore system in shale basins, Yukon and NWT

No research was conducted under this licence in 2012.

Pehrsson, Sally

Geological Survey of Canada Ottawa, ON pehrsson@nrcan.gc.ca

File Number: 12 404 504 **Licence No:** 15084

Region: SS **Location:** Parts of NTS Map sheets 65E, 65L, 75B, G, H, I,

J

Reconnaissance geology of the South Rae region

The Geological Survey of Canada (GSC) is upgrading geoscience knowledge of the north through its Geo-mapping for Energy and Minerals Program, with the aim of understanding the geologically least known parts of Northwest Territories in the Akaitcho region (parts of NTS 75A, 75B, 75H, 75G, 75J). Fieldwork was conducted July 22 to August 7, 2012. Work included an airborne geophysical survey in the Abitau-Rennie Lake area, used to identify bedrock features beneath glacial deposits. Researchers also reanalyzed rock and sediment samples from previous mapping campaigns. Helicopters were used to access remote regions. Fuel caches were established at Manchester and Labyrinth Lakes and empty drums were removed. Rock and soil samples (< 10 kg), have been sent for laboratory analysis to aid preparation of new geological maps. Once produced, these maps will be archived at the GSC in Ottawa. The data, maps and reports will be used in the longer term for land-use decision-making. Initial results show potential for mineralization, particularly nickel and palladium, not recognized previously. Research results will become available via the internet through publications of the GSC.

Pickart, Robert

Woods Hole Oceanographic Institution Woods Hole, MA United States rpickart@whoi.edu

File Number: 12 404 742 Licence No: 15110 (Multi-year licence)

Region: IN **Location:** The shelf edge, in the region from the US/Canada border to the entrance of Amundsen Gulf

Assessment of the western Arctic Boundary current

The purpose of this research is to characterize the western Arctic Boundary current (which flows at the edge of the shelf), in order to understand its role in dictating shelf-basin exchange of water and materials, as well as how it impacts the ecosystem of the region, including the occurrence of marine mammals. Cruise HLY1203 of the US Coast Guard Cutter Healy took place from October 5 to 25, 2012. Researchers used a combination of year-round subsurface moorings in the boundary current (deployed in US and Canadian waters), and seasonal (summertime) shipboard observations, including measurements downstream in Canadian waters. During the fieldwork, all of the moorings were successfully deployed. Researchers carried out a hydrographic survey of the boundary current from Barrow Canyon along the continental slope into Canadian waters to the mouth of Amundsen Gulf. The survey consisted of six cross-slope transects measuring the ocean's conductivity, temperature, and depth. This instrumentation used was equipped with a transmissometer, fluorometer, and oxygen sensor. Niskin bottles were used for taking water samples to measure salinity, dissolved oxygen, nutrients, dissolved inorganic carbon, total alkalinity, oxygen isotopes, and chlorophyll. Velocity measurements were made using the hullmounted acoustic Doppler current profiler (ADCP). This project is an ongoing collaboration between American and Canadian scientists.

Pisaric, Michael

Carleton University
Ottawa, ON
michael pisaric@carleton.ca

File Number: 12 404 640 **Licence No:** 14995 (Multi-year licence)

Region: IN, GW Location: Blueberry site (68°27'95" N 133°50'47" W);

lakes north of Fort McPherson on the western edge of the

Mackenzie Delta

Examining the impacts of climate change on aquatic and terrestrial ecosystems of the Mackenzie region, NWT

The objective of this ongoing research is to document the impacts of changing climate on Husky Lakes, especially the impact of thawing permafrost. Up in the mountains the permafrost is melting and the ground is collapsing. Streams flowing into Husky Lake are carrying sediment from these slumps. In 2012, researchers collected lake sediment cores from the bottom of Husky Lake, as well as deployed sediment traps in Husky Lake to record how much sediment was entering the lake from mountain streams. These sediment traps consisted of a plastic tube (about 20 cm in length) attached to the top of a pop bottle. The bottom of the pop bottle was cut off, creating a large funnel to collect sediment. Several of these traps were left floating beneath the water surface in Husky Lake from June until August. Amazingly, the sediment traps collected much more sediment than anticipated. Sediment filled the entire plastic tubes and in some cases filled part of the pop bottles as well. It is hypothesized that the large amounts of sediment entering Husky Lake this summer can be attributed to one particular rain event this summer. Colleagues who have positioned weather stations, with rain gauges, nearby in the Mackenzie Mountains recorded a single rain event this summer approaching 100 mm. It is likely that much of the sediment that made it into the traps arrived during this rain event.

Pratt, Brian

University of Saskatchewan Saskatoon, SK brian.pratt@usask.ca

File Number: 12 404 167 **Licence No:** 15145

Region: SA **Location:** Headwaters of Ravens Throat River, near

Redstone and Natla rivers

Cambrian stratigraphy

The purpose of this research was to study a unique fossil deposit of the ancient sea floor at the headwaters of Ravens Throat River, north of the South Nahanni River. This fossil deposit has exposed deeper water sedimentary rocks belonging to the Rockslide Formation, which are over 500 million years old. It exhibits elements of soft-bodied invertebrate fauna, plus a form interpreted to be a type of seaweed. Most sedimentary rocks preserve only shells and skeletons, the hard parts that escaped rotting on the sea floor. Over three weeks of fieldwork in July and August, researchers documented the geological setting, measured the stratigraphic section, and made collections of rocks and fossils, mainly specimens from the scree slope below the mountainside exposure. During the fall, the researchers organized and curated the entire collection of fossils, and made the necessary preparations for assigning each specimen a unique number provided by the Royal Tyrrell Museum of Paleontology (where the material will be archived). Specimens were prepared and photographed in the standard ways, that is, under air, under alcohol, and after dusting with ammonium chloride to bring out certain details. The fossilized species were measured and counted to provide population statistics. The rock samples were also sent to the in-house X-ray diffraction laboratory and the Saskatchewan Research Council microprobe laboratory for mineral and chemical analysis. Samples were sent to the in-house petrology laboratory for the preparation of transparent thin sections for optical analysis.

Prinsenberg, Simon

Fisheries and Ocean Canada -- Bedford Institute of Oceanography Dartmouth, NS simon.prinsenberg@dfo-mpo.qc.ca

Response of coastal sea ice properties of the Mackenzie Delta to climate change

No summary was submitted for this licence. This project is not in compliance with licencing

requirements.

Quinton, William Wilfrid Laurier University Waterloo, ON wquinton@wlu.ca

File Number: 12 404 570 Licence No: 15005 (Multi-year licence)

Region: DC **Location:** Within the Scotty Creek drainage area

Understanding and prediction of permafrost thaw impacts on northern water resources

In 2012, research at Scotty Creek has focussed on 1) understanding the rates and patterns of permafrost thaw, and the physical and biological processes that control it; 2) developing sciencebased tools to predict the rate and pattern of permafrost thaw; 3) understanding and predicting the impact of permafrost thaw on ecosystems and water resources; and 4) developing appropriate mitigation strategies. Significant progress was made in all four areas. For example, it was shown that over the last half century, the permafrost cover at Scotty Creek has declined from about 72% to about 40%, and that the rate of permafrost disappearance is accelerating. It was also shown that permafrost thaw results in conversion of forests to tree-less wetlands. Although these changes are driven by a warming climate, it was found that permafrost thaw is also initiated where trees are removed by fire, disease or human disturbance. Another study examined the impacts of seismic lines on permafrost thaw, and the associated publication recommended appropriate mitigation strategies. Current research is focussed on understanding the integrated ecohydrological behaviour of ecosystems in the context of thawing permafrost so that predictions of the rate and pattern of thaw and associated land-cover change (e.g. loss of forest) can be predicted with confidence. The ongoing research objectives are to: (1) develop fundamental knowledge of the major ecosystems and estimate the amount of water present. The watershed responses to changes in permafrost regime and the rate and trajectory of such changes will also be examined; (2) develop and test a new suite of eco-hydrological predictive tools for simulating the responses of ecosystems to permafrost thaw; and (3) apply the new integrated ecohydrological models to predict terrestrial and aquatic ecosystem responses to permafrost thaw extending to the year 2062. Implications of permafrost thaw on water quality are also being examined, including the impact on methyl mercury concentrations in wetlands, lakes and streams.

Reimink, Jesse University of Alberta Edmonton, AB

reiminkjesse@gmail.com

File Number: 12 404 761 Licence No: 15116 (Multi-year licence)

Region: NS Location: Acasta River Region

Petrogenesis of the Acasta Gneiss Complex: Ancient rocks revisited

The purpose of this research was to further study a set of ancient rocks called the Acasta Gneiss Complex. Fieldwork during the 2012 field season took place from July 14 to 29. During this time researchers used previously published geologic maps and samples collected during the 2011 field season to locate and document further samples of interest. A small (1 km x 1 km) area -

containing units of lower strain gradient than many areas with the Acasta Gneiss Complex - was identified and subsequently mapped in detail. Rock units of various ages (3.96-3.5 billion years) are suspected to occur within this area. Geochemical analysis is ongoing but preliminary results suggest multiple age components of sufficient quality as to be beneficial for the study of the rock's formational environments. Geochemical analysis will continue in the following months.

Sachs, Torsten

GFZ German Research Centre for Geosciences Potsdam Germany tsachs@gfz-potsdam.de

File Number: 12 404 784 Licence No: 15063 (Multi-year licence)

Region: IN Location: Mackenzie Delta - Along coastline between

Demarcation Point and Inuvik

Airborne measurements of methane (AIRMETH)

The objectives of this ongoing study are to quantify the surface-atmosphere methane emissions over large areas, and analyze the influence of different surface and vegetation characteristics on these emissions. Quantifying present methane emissions from the vast arctic permafrost wetlands and shelf areas is an essential prerequisite for identifying possible warming-induced future changes to the arctic carbon cycle, as well as for accurate model representation of regional to global methane contributions from the arctic. Current estimates are highly uncertain - as measurements are sparse and very localized. Methane emissions are known to be extremely variable over space and time. Researchers use the research aircraft Polar 5 for measurements across the entire North Slope of Alaska and the Mackenzie Delta. These measurements were made between July 4 to 10, 2012. Researchers observed clearly increased methane concentrations throughout the entire atmospheric boundary layer with a sharp drop to background levels above. Despite not having seen the peak of the growing season in 2012, strong regional differences were visible both on the North Slope and in the Mackenzie Delta. In the Mackenzie Delta, a generally higher level of methane concentrations was observed in the boundary layer, as well as clear signs of night time built-up of methane close to the surface. Methane fluxes were about three times higher in the Mackenzie Delta than on the North Slope and higher in the outer Delta than in the southern parts.

Schroder-Adams, Claudia

Carleton University
Ottawa, ON
csadams@earthsci.carleton.ca

File Number: 12 404 786 **Licence No:** 15085

Region: IN Location: Horton River; Smoking Hills Formation (70°02'

06.62"N, 126°56' 26.3" W); Boundary Creek; tributary to

Big Fish River (68°30' 30" N, 136°23' 50" W)

Upper Cretaceous paleoenvironmental reconstruction of the gateway between the Boreal and Western Interior Seas: The Boundary Creek and Smoking Hills Formations, NWT

The purpose of this research was to gain a better understanding of the paleoenvironmental setting of a Cretaceous-aged ocean gateway between the Boreal Sea and the Western Interior Sea that flooded large parts of North America. In July of 2012 a successful field season was conducted addressing Cretaceous-aged sediments at Horton River and along Boundary Creek and Big Fish River, east and west of the Mackenzie Delta, respectively. A total of 117 sediment samples were

collected and several sedimentary sections were measured. Samples for micropaleontological analysis were processed in order to extract marine microorganisms that provide information about the age of the sediments and the paleoenvironment of these ancient marine basins that occupied these regions between 70 to 100 million years ago. This was a time of frequent volcanism witnessed by ancient ash layers within the sediments. Analysis of these ancient ash beds revealed that their sources are volcanic arc systems; these were located along the tectonically active western margin of North America. In contrast, similar aged ashes from the Queen Elizabeth Islands have a different volcanic source. Samples are also being processed for carbon isotope stratigraphy - a method that detects times of major carbon burial and phases throughout earth's history when the seafloor had no oxygen and large amounts of organic matter gets buried in the sediments. These phases were relatively common during the Cretaceous Period, when the earth was considerably warmer than today. Micropaleontological and carbon isotope analyzes are ongoing.

Smith, Sharon

Geological Survey of Canada Ottawa, ON sharon.smith@nrcan.gc.ca

File Number: 12 404 657 Licence No: 15053 (Multi-year licence)

Region: IN, GW, SA, DC **Location:** In and around Jean Marie River; Fort Simpson;

Wrigley; Tulít'a; Norman Wells; Fort Good Hope;

Tsiigehtchic; Tuktoyaktuk

Permafrost monitoring and collection of baseline terrain information in the Mackenzie Valley Corridor, NWT

Permafrost monitoring sites throughout the Mackenzie corridor (Inuvialuit, Gwich'in, Sahtú, Dehcho regions) were visited in August and September 2012 to acquire ground temperature and active layer data. Data records for 40 monitoring sites were extended to: better characterize of the permafrost conditions; facilitate an understanding of the natural variability in permafrost thermal and active layer conditions; ensure availability of adequate baseline permafrost information to support land management decisions in the region. Permafrost in the discontinuous permafrost zone, which covers a large portion of the corridor, is generally warmer than -2° C. Permafrost temperatures generally continue to increase in the region and the overall range in temperature is decreasing. Analysis is in progress to characterize the variability of the onset of seasonal ground freezing and thawing. Ongoing collection of data from monitoring sites is planned to better assess the impact of climate change on the permafrost environment. A detailed report, including graphical and tabular summaries of data, is in preparation and will be sent to relevant organizations in the region.

Snyder, David

Natural Resources Canada Ottawa, ON dsnyder@nrcan.gc.ca

Region: SA, NS, SS

File Number: 12 404 548 Licence No: 15050 (Multi-year licence)

Location: Maintained stations at: Hepburn Lake: Sulky Lake; Lac des Bois; Simpson Lake; Colville; Kugluktuk; near Ulukhaktok. New stations: Johnson Point on Banks Island;

near Thor Lake

Teleseismic studies in the Wopmay

This ongoing study investigates the structure and composition of the Earth's crust and mantle. The researchers are seeking better methods to characterize diamond reservoirs, in order to make exploration more efficient and low impact. Three seismic stations were removed and one new one established in the quarry at Norman Wells in August. Eight stations remain in the field recording earthquakes. Recorded earthquakes have continued to be analyzed and used to map three major seismic boundaries at 40, 100 & 150-km depths beneath central Northwest Territories. The middle boundary is related to eruption of the kimberlite field at Lac de Gras and its diamond mines and helps to locate the diamond source region.

Sofko, George

University of Saskatchewan Saskatoon, SK george.sofko@usask.ca

File Number: 12 404 636 Licence No: 14994 (Multi-year licence)

Region: IN, GW Location: LOT 2, Block 107 PLAN 4166, Inuvik

PolarDARN (The northern hemisphere polar portion of the international SuperDARN (Super Dual Auroral Radar Network)

The goal of this long-term and ongoing project is to measure voltage patterns several hundred kilometers above the ground and out into space along the Earth's magnetic field lines. In 2012, the Clyde River radar (on Baffin Island) was added to the existing Inuvik and Rankin Inlet radars. This provides a unique opportunity for Inuvik and Clyde River to work together, as signals can be transmitted from Clyde River and received at Inuvik, or vice versa. This kind of signal geometry is used to measure signals called "forward scatter" and is expected to be quite strong. The data would provide new opportunities to study up-and-down motions near the midpoint between Inuvik and Clyde River. This unique combination of the PolarDARN radars provides an opportunity for space science projects at high latitudes that has never before been possible. For example, the Earth's magnetic field changes from being attached strictly to the Earth (closed magnetic field) to being partly attached to the solar wind and partly to the Earth (open magnetic field). The transition lies in the region examined by the three PolarDARN radars. Although there are now 29 radars in the international SuperDARN project, the three Canadian PolarDARN radars offer many new opportunities not possible with the other 26 radars. The year ahead should see some exciting new results from the PolarDARN radars at Inuvik, Rankin Inlet and Clyde River.

Spence, Christopher

Environment Canada Saskatoon, SK chris.spence@ec.gc.ca

File Number: 12 404 535 Licence No: 15107 (Multi-year licence)
Region: NS Location: The Baker Creek Basin

Investigations of the water cycle and hydrological processes of the subarctic Canadian Shield

The objective of this ongoing project is to determine the relationships between the climate, streamflow, water chemistry and permafrost in the subarctic Canadian Shield. Field activities in 2011 in the Baker Creek research catchment began with spring snow surveys and the activation of climate towers and water level stations in April. There were no people living in the research catchment in 2012. Remote measurements of meteorological conditions, evaporation, soil moisture, hydrochemistry and streamflow continued through 2012. This program involved

sampling streamflow in tributaries and at lake outlets along Baker Creek bi-weekly. Groundwater was also sampled. Samples were analyzed for ions, pH, metals, nutrients and carbon and nitrogen. This work is in support of determining how stream chemistry and frozen ground react to wetter autumn conditions during freeze up. These research questions are in response to observations that have shown streamflow in small subarctic Canadian Shield catchments changed from a predominantly nival (snowmelt) to a combined nival/pluvial (snowmelt and rainfall) regime in the late 1990s. The autumn of 2012 was dry and Baker Creek remained in a near zero flow condition, which is more typical of the pre-1995 period. Hydrochemistry and streamflow data during this type of freeze-up event were collected and are now being analyzed.

Tank, Suzanne York University Toronto, ON tanks@yorku.ca

File Number: 12 404 785 Licence No: 15069 (Multi-year licence)

Region: IN, GW

Location: Point Separation; East Channel, Inuvik; Middle

Channel, mid-Delta; East Channel at mouth; Reindeer

Channel; Napoiak Channel

Solar degradation of dissolved organic carbon in Mackenzie Delta lakes and river channels

This study examined the solar degradation of riverine dissolved organic carbon to carbon dioxide throughout the Mackenzie River Delta. The preliminary fieldwork involved 10 days of field sampling during early June 2012. Water was collected from river channel locations throughout the Mackenzie Delta and from a select series of lakes that are accessible by boat from Inuvik. Water samples have been analyzed for dissolved organic carbon concentration and composition, and some early analysis has been done to assess the photodegradation of the collected dissolved organic carbon samples.

Trimble, Annika

Aurora Research Institute Inuvik, NT atrimble@auroracollege.nt.ca

File Number: 12 404 720

Region: DC

Licence No: 15020 (Multi-year licence)

Location: Jean Marie River; Fort Providence

Solar irradiance monitoring in Jean Marie River and Fort Providence

The objective of this ongoing research project is to measure solar irradiance levels (the strength of sunlight over a certain area) in Jean Marie River and Fort Providence, in order to support prefeasibility studies on the use of solar energy in those communities. Data collected were used to support the pre-feasibility study completed in 2012 for Jean Marie River (available at www.nwtresearch.com). Should the community be interested in developing renewable energy projects, preliminary results indicate that solar would be a better option than wind energy. Analysis for Fort Providence has not yet been completed. However, due to the proximity of the two communities, it is probable that solar readings will be similar those of Jean Marie River.

Trimble. Annika

Aurora Research Institute Inuvik, NT atrimble@auroracollege.nt.ca

File Number: 12 404 720 Licence No: 15022 (Multi-year licence)

Region: NS Location: Wekweètì

Wind energy monitoring in Wekweètì: 2010 - 2012

The objective of this wind monitoring project was to quantify Wekweètì's wind energy resources. The project technician travelled to Wekweètì in July, 2011 to hire and train a new wind monitor and to perform maintenance on the two wind monitoring towers there. Data collection has been a challenge during this project, but wind speeds continue to be measured. Analysis of Wekweètì's wind resource is ongoing, and will be shared in a report in 2012/13.

Trimble, Annika

Aurora Research Institute Inuvik, NT atrimble@auroracollege.nt.ca

File Number: 12 404 720 **Licence No:** 15033

Region: NS **Location:** 62°4' 11.95"N, 112°37'10.7"W

Wind energy monitoring at Thor Lake 2012

The objective of this wind monitoring project was to quantify Thor Lake's wind energy resources. Due to data loss at the Thor Lake site over two winters (2009/10 and 2010/11), it was decided that the wind monitoring tower should be left up another year instead of being taken down in the fall. To prevent further data loss, equipment upgrades were made to the wind monitoring tower. Analysis of Thor Lake's wind resource is ongoing, and will be shared in a final update in 2012/13.

Trusler, Scott

Minerals and Metals Group (MMG) Canada Vancouver, BC scott.trusler@mmg.com

File Number: 12 404 781 Licence No: 15056 (Multi-year licence)

Region: NS **Location:** South of the proposed Izok project along and in

the vicinity of the Coppermine River

Biophysical baseline studies in support of the Izok Project

The purpose of this project was: (1) to refine understanding of baseline conditions in the vicinity of the Izok Project; (2) to provide input to design mitigation; and (3) to evaluate potential changes in the environment in relation to the Izok Project. Surface water and sediment quality surveys, vegetation and soil sampling, and hydrological surveys took place in the vicinity of the Coppermine River in 2012.

Turner, Elizabeth

Laurentian University Sudbury, ON eturner@laurentian.ca

Region: GW, SA **Location:** West side of the Arctic River

Stratigraphy of the Misty Creek Embayment

The purpose of this research is to understand the geological evolution of a deep-marine environment that existed in the northern Mackenzie Mountains between 500 and 450 million years

ago. Fieldwork for this project took place from July 29 to August 20, 2012 at one of the planned sites: 404000E/7155500N (west side Arctic Red River). A two-person camp was established by helicopter and moved to another part of the same slope (about 2 km away) halfway through the work. Work took place on foot along the nose of one slope and up a creek in another. The work was predominantly measuring and describing rocks in stratigraphic succession. A small number of rock samples were also collected (using a rock hammer) for later analysis in the lab. Since completion of this year's fieldwork, the data collected in the field is being analyzed. The collected material will be prepared for geochemical analysis starting in December, 2012. Similar work is likely to be completed in 2013 at other sites.

Urbanic, Jane Challen

Environment Canada Burlington, ON jane.challen-urbanic@ec.gc.ca

File Number: 12 404 741 Licence No: 15049 (Multi-year licence)

Region: IN, SA, DC, SS **Location:** Rae; Paulatuk; Fort Providence; Edzo;

Tuktoyaktuk

Arctic Wastewater Research

No summary was submitted for this licence. This project is not in compliance with licencing requirements.

van der Sanden, Josephus

Natural Resources Canada -- Canada Centre for Remote Sensing Ottawa, ON sanden@nrcan.qc.ca

File Number: 12 404 709 **Licence No:** 15018

Region: IN, GW Location: Mackenzie River Delta Middle Channel (from

Point Separation to just North of Oniak Island); selected

locations in the outer Delta

RADARSAT observations of river ice and flood patterns in the Mackenzie River Delta

The principal objective of the research carried out during 2012 was to assess and develop the utility of radar satellite images for the mapping of characteristics of winter ice cover and lakes and rivers, such as: ice type, ice thickness, and bottom fast vs. afloat. Fieldwork facilitates the correct interpretation of available radar images and the generation of reliable information products. In 2012, fieldwork was carried out at selected locations in the Mackenzie Delta from February 22, 2012 to March 9, 2012. The data collected capture properties of the ice cover (e.g. ice type, layering, thickness, roughness, air inclusions), as well as the overlying snow cover (e.g. snow type, layering, thickness). During the winter of 2011/2012 radar images were acquired by the Canadian RADARSAT-2 (R2) and the Italian Cosmo-Skymed (CS) satellites. R2 images were acquired every 24-days from freeze-up to breakup; a series of three CS images was acquired on March 8, 9, and 12, 2012. Analysis of the R2 and CS images focus on their potential for the mapping of ice types and ice thickness, respectively. The analysis of these data is ongoing. A report summarising the researchers' R&D to date can be found at:

http://geoscan.nrcan.gc.ca/starweb/geoscan/servlet.starweb?path=geoscan/downloade.web&search1=R=291867

Vavrek, Matthew

Royal Ontario Museum Toronto, ON matthew@matthewvavrek.com

File Number: 12 404 801 **Licence No:** 15132

Region: SA **Location:** Headwaters of the East Little Bear River

Palaeontology of the Summit Creek Formation, western Northwest Territories, Canada

From August 20 to September 1, 2012, the Royal Ontario Museum conducted a palaeontological survey of the Summit Creek Formation, a rock unit that is exposed approximately 100 km south of Norman Wells. This work was done to search for any fossils, in particular those of dinosaurs. The field crew flew in to Tulít'a, via Yellowknife and Déline. From Tulít'a, the crew was taken by helicopter to the first campsite along the East Little Bear River for five days. During the time there, the field crew searched any exposed rock face for fossils, and found a number of fossils at several different sites. The fossils that were recovered were from both hadrosaur (duckbill) and ceratopsian (horned) dinosaurs, as well as a large carnivorous dinosaur. Because all of the fossils were fragmented, it is not possible to know exactly what types of dinosaurs were found. The second field site was approximately 10 km to the north. After two days of searching this area, no fossils were found. After this, the helicopter was used on a final day to search several additional spots that were inaccessible by foot, however no more fossils were found.

Wang, Zhaohui (Aleck)

Woods Hole Oceanographic Institution Woods Hole, MA United States zawang@whoi.edu

File Number: 12 404 740 Licence No: 15098 (Multi-year licence)
Region: IN Location: Mackenzie River near Inuvik

Towards long-term monitoring of the CO₂ system in arctic rivers

The short-term goal of this research was to initiate time-series measurements of the carbon dioxide (CO₂) system in the Mackenzie River. This work will serve as the initial step towards long-term measurements and studies of the impacts of global warming on the CO₂ systems in Mackenzie River, its estuary, and adjacent coastal waters. This year's monthly sampling from the river continued near Inuvik. All sampling was very successful, with a total of 18 water samples collected. All samples will be measured for total dissolved inorganic carbon (DIC) concentration and alkalinity. About two thirds of the samples have been processed and is ongoing. Seasonal changes of DIC and alkalinity are different this year, as compared to the last two years. Future efforts will be made to investigate the mechanisms that caused these changes. From DIC concentrations and alkalinity, pH and partial pressure of carbon dioxide (pCO₂) in water can be calculated. The results also show large seasonal changes of pH and pCO₂. Monthly sampling will continue into 2013.

Whalen, Dustin

Geological Survey of Canada Dartmouth, NS dwhalen@nrcan.gc.ca

File Number: 12 404 798 Licence No: 15119

Region: IN Location: Mackenzie Delta

Beaufort Sea coastal geoscience research

The goal of this research was to improve the understanding of physical conditions in the coastal zone in order to monitor, manage and respond to changes in the coastline. A total of twelve coastal monitoring sites were visited between the Alaska/Yukon border and McKinley Bay in August 2012. Some coastal monitoring sites along the Yukon Coast and Tuktoyaktuk Peninsula were visited for the first time in twenty years. Preliminary analysis show the Alaska border continues to erode with rates exceeding one metre per year. Parts of the coastline have receded 25 m since 1991. In comparison, Tuktoyaktuk Island continues to erode at alarming rates with no signs of slowing down receding 140 m since 1950. The fieldwork also provided the first look at a number of coastal and near shore issues with Tuktoyaktuk Harbour. Preliminary analysis of the sub-bottom stratigraphy in and around the harbour give indication that the approaches to Tuktoyaktuk Harbour are in-filling with sediment. This new information provided critical present day observations of extreme coastal change that affect a number of key sites.

Wilcockson, John

Hatfield Consultants North Vancouver, BC jwilcockson@hatfieldgroup.com

File Number: 12 404 791 **Licence No:** 15096

Region: DC **Location:** Small lakes along a proposed winter road

Prairie Creek mine towards Nahanni Butte

Fish presence and lake bathymetry

Bathymetry data were collected at nine lakes located along the proposed winter road from the Prairie Creek mine towards Nahanni Butte. This dataset will be used to determine the suitability of each lake for water withdrawal for strengthening the road base of the winter road (following the Department of Fisheries and Ocean's guidance document for winter water withdrawal). At one lake, located at kilometre 95 from the mine, attempts were made to catch fish (lake was located at UTM 10V0465089 6812893). Fishing approaches attempted included: gillnet (soaking time 3 hours, checked regularly), electrofishing (239 seconds), minnow traps (soak time 4 hours) and angling (just under one hour). No fish were caught and no fish were seen.

Williams, Mathew

University of Edinburgh Edinburgh United Kingdom mat.williams@ed.ac.uk

File Number: 12 404 802 Licence No: 15133 (Multi-year licence)

Region: NS Location: The Ingraham Trail

Carbon cycling linkages of permafrost systems [CYCLOPS]

No research was conducted under this licence in 2012.

Wolfe, Stephen

Natural Resources Canada Ottawa, ON swolfe@nrcan.gc.ca

File Number: 12 404 549 Licence No: 15003 (Multi-year licence)

Region: NS

Location: Along the highway route west of Yellowknife and along the Ingraham Trail east of Yellowknife; the

Baker Creek Watershed north of Yellowknife; along the Tibbit to Contwoyto Winter Road based out of the Lockhart

and Lac de Gras facilities

North Slave permafrost study: Characterizing and predicting discontinuous permafrost for climate change adaptation

Fieldwork was conducted between June and November, 2012 in the Great Slave region along Highways 3 and 4, and the Tibbitt to Contwoyto Winter Road. Permafrost cores ranging from 1.0 to 8.2 metre depth were obtained from one study sites in peatland and birch forest settings. Thaw depths, soil types, visible ice moisture contents, and bulk densities were determined, with grain size, water geochemistry and geotechnical tests still to be determined. These data study geotechnical conditions associated with permafrost soils in the area. Temperature data continues to be collected and monitored at a number of sites including: active layer temperatures from birch, spruce forest and four peatland sites; ground temperatures from three burn sites, and eight peatland, birch and spruce forest sites; and ten air, ten active-layer, seven shallow-water, and four lake-bottom temperature monitoring sites. These data are used to understand potential climatic gradients and the effects of water on local permafrost conditions. Four new thermal monitoring sites were installed on the Tibbitt to Conwoyto Winter Road. Helicopter surveys were undertaken to validate remote sensing interpretations of surficial geology and vegetation cover mapping in NTS map sheets 85J, and P. Samples were collected for dating.

Wrona, Frederick

University of Victoria Victoria, BC wrona@mail.geog.uvic.ca

File Number: 12 404 711 **Licence No:** 15108 (Multi-year licence)

Region: IN Location: Noell Lake

Noell Lake ice study - Hydro-ecological responses of arctic tundra lakes to climate change and landscape perturbation

The objective of this research is to improve knowledge on lake ice and its effect on food webs and productivity in arctic tundra upland lake systems. A better understanding of this, will help produce better predictions about the changes that could occur under changing climate. In late-September 2011, prior to freeze-up, an automated ice buoy and subsurface mooring system was re-deployed successfully (after removal for servicing of the instruments) in Noell Lake for continuous monitoring of weather conditions, lake ice cover (e.g. formation, growth over winter, breakup in spring), light penetration into the lake (through ice in winter), and water quality. The monitoring system collected data over the 2011-2012 winter period. After spring breakup in 2012, the buoy was removed from the lake for servicing and replaced with another instrumented buoy. The buoy and mooring system then continued to collect water quality information through the open-water season. In addition, manual grab samples for standard water quality and aquatic biological parameters were taken seasonally during 2012 to compare with and validate measurements made by the instrumented buoy/mooring system. These data are allowing researchers to examine lake ice and its effects on the food web/productivity through the winter, and the character of food webs/productivity during the ice-free season.

SOCIAL SCIENCES 2012

Anakin, Megan
University of Otago
Dunedin, Otago
New Zealand
megan.anakin@otago.ac.nz

File Number: 12 410 931 Licence No: 15154 (Multi-year licence)

Region: SS Location: Schools in the South Slave Divisional Education

District, specifically, elementary and high schools and teachers who have been involved with the SmartLearning approach prior to the commencement of the 2012-2013

school year

Assessing the SmartLearning Project

The purpose of this ongoing study is to identify the impact of the SmartLearning Project on student outcomes. The SmartLearning Project gathers quantitative and qualitative data at classroom, school, and school district levels annually as a part of regular instructional and assessment practice. Its assessment involves action research conducted by classroom teachers and inquiry leaders in schools in British Columbia, Alberta, and the Northwest Territories. The South Slave Divisional Education Council authorized consent for their staff to participate in the study on September 15, 2012. Three staff members at Princess Alexandra School, Hay River have indicated interest in participating in the Assessing the SmartLearning Project during the 2012-2013 school year. No other data have been collected by the research time at this time.

Bathe, Adam

Blyth & Bathe Inc. Fort Smith, NT adam@blythandbathe.com

File Number: 12 410 915 **Licence No:** 15075

Region: DC **Location:** Within Wrigley townsite; Wrigley Property

Devonian Metals Wrigley Property TEK assessment

Located in the traditional territory of Pehdzeh Ki First Nation (PKFN) near the community of Wrigley, Devonian Metals Inc. is currently exploring their Wrigley Property for zinc, lead, and silver. This study described the pattern of land use created by the PKFN and the environmental knowledge that they have developed over the centuries in this region. Specifically, the intent of the study was to document the historic, present, and traditional use of the land, water, and wildlife in the Wrigley River, Moose Pasture Creek, and Fish Trap Creek watersheds by PKFN. This study was designed collaboratively with the participants and PKFN. Interviews of seven participants

took place from May 22 to 27, 2012. The interviewees included a cross section of knowledgeable elders and harvesters. Several types of data were collected during the study. The primary sources of data included the audio recordings and the maps from the interviews with the study participants. In addition to the audio recordings, notes were taken to facilitate the conversion of interview map data into a GIS. During this type of study, two classes of information are generally produced. The first class of information is typically more spatial in nature and oriented towards the traditional use and occupancy of the land. Some examples of this would be the description for the location of a cabin or a good lake for duck hunting. Additionally, in many cases traditional ecological knowledge is also mapped spatially (e.g., seasonal caribou habitat). The other class of information collected during the interviews can be categorized as traditional knowledge and is often (although not always) disassociated from specific locations and is not spatial in nature. Ethics surrounding respect for the land would be an example of traditional knowledge collected that could not be mapped. Initial interviews were verified and clarified by participants. The information from this study will be used by PKFN and Blyth & Bathe Inc. to inform Devonian Metals about potential impacts to environmental and cultural resources by the development activities.

Baumann, Britt

University of Waterloo Kitchener, ON bbaumann@uwaterloo.ca

File Number: 12 410 895

Region: IN, NS

Licence No: 14998 (Multi-year licence)

Location: Inuvik; Yellowknife; Tuktoyaktuk

The Impact of the Priest Decline on the Canadian Roman Catholic Church

No summary was submitted for this licence. This project is not in compliance with licencing requirements.

Bishop, Nicole

Rescan Environmental Services Ltd. Yellowknife, NT nbishop@rescan.com

File Number: 12 410 891 Licence No: 15080 (Multi-year licence)

Region: DC, NS, SS **Location:** Communities associated with the Courageous

Lake Project

Courageous Lake Project - Social and economic sciences research

Socio-economic and land use work in 2012 entailed updating community profiles through desk-based research. Data sources included recent documents released on the Mackenzie Valley Environmental Impact Review Board's website, as well as results from the 2011 Census (including population and demographics, age, gender, households and dwellings). Site visits to the Courageous Lake exploration site were held the week of August 27, 2012 with participants from the Yellowknives Dene First Nation, Łutsel K'e First Nation, North Slave Métis Alliance and Northwest Territories Métis Nation. Site visits provided a detailed overview of current activities and plans based on current project design, with representatives from each Aboriginal group providing feedback. A meeting was held in September 2012 with members of the Tłįchǫ Government and the Kwe Beh Working Group to provide scope and direction on the socioeconomic research activities to be carried out within the Tłįchǫ communities in the near future. Preliminary contact information was exchanged in order to identify potential dates for data gathering.

Borowitz, Michelle

University of Alberta Edmonton, AB borowitz@ualberta.ca

File Number: 12 410 873 Licence No: 15012 (Multi-year licence)

Region: SS Location: Fort Resolution

Human dimension of river resource development and transboundary water security in the Peace-Slave River Basin

The goal of this research was to document how issues and practices of trans-boundary water security and river resource developments affect local aboriginal communities in the South Slave region and the Peace Region. As part of the ongoing project, the researcher returned to Fort Resolution between May 28 and June 15, 2012. During this time they followed up with research participants who were interviewed in 2011. In addition, interviews were complete with those individuals originally missed in August 2011. The fieldwork is not complete, as such there are no results yet to report. Research will be ongoing on 2013.

Callingham, Christina

University of Ottawa Ottawa, ON christina.callingham@gmail.com

File Number: 12 410 935 **Licence No:** 15159

Region: IN, NS Location: Phone interview with individuals who are from

Paulatuk (currently living in Alberta) and Whatì (currently

living in Yellowknife)

Aboriginal youth involvement in a youth-driven program: A narrative exploration of the process and impact of engagement

The question guiding this research was: What narratives emerge, and are co-constructed, when exploring the experience of Aboriginal youth who engage in youth development programs aimed at fostering youth engagement? Two individuals from Northwest Territories were approached to participate in the study. One person participated in an interview about their experience of youth engagement through the ACTIVATE program. Results suggest the ACTIVATE was a positive experience that helped build confidence and contributed to goal building.

Cash, Penny

University of British Columbia, Okanagan Kelowna, BC penny.cash@ubc.ca

File Number: 12 410 886 Licence No: 15028 (Multi-year licence)

Region: NS Location: Yellowknife

Quality Workplace Environment

No summary was submitted for this licence. This project is not in compliance with licencing requirements.

Chugh, Pawan

NWT Business Development and Investment Corporation Yellowknife, NT

pawan chugh@gov.nt.ca

File Number: 12 410 926 Licence No: 15141 (Multi-year licence)

Region: GW, DC **Location:** Fort Liard; Fort Providence; Fort McPherson

Assessing Process and Outcomes of Government Funded Projects in Northern Canada Aboriginal Communities

No research was conducted under this licence in 2012.

Daitch, Sarah University of Victoria Fort Smith, NT s_daitch@hotmail.com

File Number: 12 410 937 **Licence No:** 15164

Region: IN, GW, SA, DC, NS, SS Location: Katlodeeche First Nation; Hay River; Ndilo

Yellowknife; Déline; Aklavik; Fort McPherson; Tuktoyaktuk, Inuvik; Behchokò; Fort Simpson; Fort Providence; Fort

Smith; Fort Resolution

An ethical space for dialogue about difficult history: Program evaluation of a residential school education module in Canada's Northwest Territories and Nunavut Territory

The guiding question for this research was: How can Northern Canadian youth connect difficult history with their identity, and become capable and committed to their communities? This study was conducted in collaboration with the Territorial Departments of Education in the Northwest Territories and Nunavut. The territorial education departments developed a mandatory curriculum module regarding the history and impacts of residential schools, piloted in high schools during the 2012-2013 academic year. The curriculum aimed to teach difficult history of the attempted assimilation of Indigenous students through residential schools. This research explored how the new education module influenced student thinking and behaviour, including the development of critical thinking skills and sense of community involvement and leadership. Because it is a region undergoing rapid development, fostering critical citizenship amongst students is vital not just to the North, but to all of Canada. Findings indicate that students are developing empathy, understandings of history, critical thinking, ethical decision making strategies, and hope for the future through the new course. After teaching the course, teachers had greater awareness of intergenerational effects of the schools. They also felt more confident in supporting students towards learning objectives. However, the new course appears to have had limited effectiveness on empowering students to take active roles in shaping their communities and connect history to their identities. These, and other findings were used by the respective Territorial curriculum teams as they created a second version of the new course for 2013-2014. Changes made as a result of this study include: reduction of materials and adjustment of time spent on each activity to enable completion of all activities; addition of a new section to the module on students' final project options, with examples provided; and the development of a video of the Health Canada Support session, to assist teachers in managing emotions in the classroom and to provide guidance for self-care for teachers and students. The results of this study will better position the Departments of Education to optimize student learning.

Douglas, Vasiliki

University of Northern British Columbia Prince George, BC douglasv@unbc.ca **File Number:** 12 410 869 **Licence No:** 15134

Region: IN Location: Aklavik; Ulukhaktok; Paulatuk; Tuktoyaktuk

Climate change impacts on Inuit food security in Canada's western arctic: Constructing a comparative anthropological model to guide adaptation planning

This project examined the critical impacts of environmental, economic and social change on traditional food availability to help determine the range of socially, culturally and physically acceptable adaptations. Researchers conducted a workshop in Inuvik on July 11 to 12, 2012. Results of contaminants research, food security research and Inuit Health Survey research were presented to participants from each of the participating communities. Participants provided feedback on fieldwork conducted in 2011 and suggested directions for future research and action in the light of results obtained from field research. Results of the workshop were collated, analyzed and incorporated into the final reports submitted to each community.

Fraser, Gail York University Toronto, ON gsfraser@yorku.ca

File Number: 12 410 913 **Licence No:** 15064

Region: IN, NS Location: Yellowknife; Inuvik

The environmental assessment process of Canadian 'frontier' oil and gas

No research had been conducted under this licence in 2012.

Gordon, Dylan

University of Toronto Toronto, ON dylan.gordon@utoronto.ca

File Number: 12 410 920 **Licence No:** 15092

Region: DC, NS, SS Location: Outside Behchokò; Yellowknife

Ethical value and market value in a Canadian Wild Food Network

Research was conducted from June 2 to July 1, 2012. The main research method was participating in and observing the life and work of a camp of mushroom buyers and pickers from British Columbia. They were picking wild mushrooms (called morels) in an area burned by a forest fire the year before. It was located about 100 km south of Behchokò on the highway. Interviews were also conducted with business people in Yellowknife who had experience harvesting, purchasing and selling wild food products from the Northwest Territories. Analysis of the results are now taking place.

Hall, Karen

Walter & Duncan Gordon Foundation Victoria, BC doxhae@uvic.ca

File Number: 12 410 916 **Licence No:** 15086

Region: NS Location: Yellowknife; Behchokò

Jane Glassco Arctic Fellows

No summary was submitted for this licence. This project is not in compliance with licencing requirements.

Hampton, Mary University of Regina Regina, SK mary.hampton@uregina.ca

File Number: 12 410 906 Licence No: 15103 (Multi-year licence)

Region: IN, GW, SS, NS **Location:** 2009 - 2012 Statistical data from the RCMP on all NWT communities based on GIS mapping results of

incidents and services for women who experience intimate

partner violence

Rural and northern community response to intimate partner violence

The goals of this ongoing research is to: (1) integrate several sources of data to create an action plan that maps the problem of intimate partner violence; (2) create narratives describing community response in to this violence; and (3) to generate a grounded theory as a practical tool to create and sustain non-violent communities. In 2012, researchers collaborated to complete an environmental scan and resource document at all project sites across Canada. In addition, they worked with the RCMP national office in Ottawa to collect information on incidents of intimate partner violence in the rural and northern regions. The resources from the environmental scan and the incidents of intimate partner violence are being uploaded into GIS maps that will allow further analysis of collected data. Team members from all regions were involved in an additional interviewing to capture qualitative information and stories that will contribute to the research questions. Researchers have continued to work with the Coalition against Family Violence, local RCMP, the Department of Justice and Health and Social Services.

Hanson, Dorie University of Alberta

Hay River, NT dhanson@ssdec.nt.ca

Region: SS Location: Princess Alexandra Middle School, Hay River

Incentive-based home reading challenges

By exploring the experiences of three participants of incentive-based home reading challenges three themes emerged: motivation, self-esteem, and social interactions. These themes developed through the interconnections between categories: incentives, home involvement, and academic achievement. By using incentives in the reading challenges, the participants were motivated to read at home. Home reading occurred on a daily basis and the participants identified academic benefits from their daily reading efforts. Improved reading ability, writing skills, increased vocabulary and improved comprehension were identified as some of the academic benefits realized by the participants. The participants' self-esteem increased as their reading improved and academic performance increased, which further motivated their continued participation in the reading challenges. All three participants indicated they preferred reading challenges that allowed for classroom discussions and involvement with other students. The participants frequently earned the final reading challenge incentive, which provided further opportunities to interact with other students. The final incentive included pizza luncheons and school-outings, which further provided opportunity for social interactions with other students. Through their increased motivation to read, increased self-esteem and opportunities for interactions with others, the

participants indicated their desire to reading increased significantly. After leaving the middle school, students were intrinsically motivated to read often, preferring reading over other activities.

Hay, Amie

Beaufort Delta Health and Social Services Authority Inuvik, NT amie hay@gov.nt.ca

File Number: 12 410 911 **Licence No:** 15043

Region: IN, GW **Location:** Kindergarten classroom in the following

communities: Fort McPherson; Tsiigehtchic; Aklavik;

Tuktoyaktuk

How standard is standard? Comparing culturally sensitive language norms versus CELF P2 in NWT

The aim of the project was to determine how to average kindergarten student in four selected communities of the Beaufort-Delta region score on the most frequently used 3-6 year test of language development. The purpose was to help the Speech-Language Pathologists gain more understanding of the scores that are being achieved during testing (e.g., "Johnny has scored this on the test but how is Johnny scoring against his peers?").

Henderson, Joanna

Centre for Addiction and Mental Health Toronto, ON joanna_henderson@camh.net

File Number: 12 410 916 Licence No: 15076 (Multi-year licence)

Region: DC Location: Fort Simpson; Fort Liard; Fort Providence

Enhancing youth-focused, evidence-informed treatment practices through cross-sectorial collaboration

The overall objectives of this ongoing national project are: (1) to provide training in the use of a youth screening tool for mental health and substance use concerns to service providers from agencies who have agreed to all use the same screening tool; (2) to understand the impact of training and project participation on service providers; and (3) to better understand youth mental health and substance use needs through their responses on the screening tool. This project is being implemented in 7-10 communities across Canada. To date, the following aspects of the project have been completed in the Northwest Territories: community visits; service provider training and data collection completed; and the youth data collection initiated. This research will continue in 2013 and include: completing the youth data collection; data analysis with community; and the report preparation.

Hodgkins, Andrew

University of Alberta Edmonton, AB hodgkins@ualberta.ca

File Number: 12 410 649 Licence No: 15120 Region: IN, GW Location: Inuvik

A comparative analysis of vocational education and training programs in Northern Canada

No research was conducted under this licence in 2012.

Horowitz, Wayne

The Hebrew University
Jerusalem
Israel
whorowitz@mscc.huji.ac.il

File Number: 12 410 903 **Licence No:** 14986

Region: IN, GW Location: In and around Inuvik; Tuktoyaktuk

Constellations and astral Lore: Far north and near east

In February 2012, the researcher conducted a sequence of astronomical observations, both in Inuvik (within the town site and in rural areas beyond the light pollution), and Tuktoyaktuk. The research observed the apparent behaviour of stars and constellations, in particular the fact that the northern stars appear to rotate around the sky rather than to move in an east-west direction as observed from more southern latitudes, planetary and lunar phenomena, as well as the *aurora borealis* (the northern lights). These observations provide context for materials found in the written sources, which include materials ranging from modern ethnographic research to the diaries of early European explorers in the arctic in the 19th century. This methodology will now be applied to study both the modern Arctic sky and the Ancient Mesopotamian sky, correlating a database of written sources, astronomical cuneiform texts in the case of Ancient Mesopotamia, with observations of astronomical phenomena in the sky for the Ancient near east and the sky over Israel.

Jaker, Alessandro

Goyatikö Language Society New Brighton, MN United States amjaker@gmail.com

File Number: 12 410 648 Licence No: 15122 (Multi-year licence)

Region: NS, SS Location: Dettah; Ndilo; Yellowknife; Łutsel K'e

Teaching our Yellowknives Dene languages

The ultimate goal of this research is to produce an intermediate-level reader, and a verb dictionary in Dogrib and Chipewyan that teachers can use in the classroom to teach Dene culture and languages. In 2012, researchers made very good progress towards the verb dictionary and reader in both Yellowknives Dene languages: Weledeh (Dogrib) and Chipewyan. Researchers collected approximately 100 verbs in Chipewyan and 200 in Weledeh. In Chipewyan, they recorded and transcribed two stories about fish, while in Weledeh they finished several stories about muskrat, arctic hare, and arctic fox. The researchers have shared some the current drafts of the Chipewyan materials with the language programs in Deninu Kue and Łutsel K'e, who were very interested. The final version is not yet ready to publish and work will continue in 2013 and 2014. The end of 2014 is the current goal for final publication (most likely through Alaska Native Language Center publications).

Jardine, Cindy

University of Alberta Edmonton, AB cindy.jardine@ualberta.ca

File Number: 12 410 882 Licence No: 15163 (Multi-year licence)

Region: SS Location: Ndilo

Engaging aboriginal youth in tobacco prevention using social media

This ongoing research explores if a social media intervention developed by Aboriginal youth (specifically videos to be available through YouTube) using a participatory approach can be an effective means for encouraging smoking prevention and/or cessation amongst youth and others in Aboriginal communities. To date, three teams involving a total of twelve high school students from the K'alemi Dene School in Ndilo, have completed their social media videos aimed at tobacco prevention and cessation. The youth worked with a local filmmaker to create story boards that represented their health messages. Each team produced, directed filmed and edited their own videos. Group interviews were conducted with the youth at the start of the project and after the videos were completed to determine their expectations and experiences. One-on-one interviews were also conducted with the K'alemi Dene School administrative staff participating in the project. The researchers are in the initial stages of interview analysis and will be able to provide preliminary results by the end of the year. The youth showcased their videos to other students, parents, elders and community members at their monthly circle ceremony at the K'alemi Dene School and during the schools' year-end celebration. They also came to Edmonton, Alberta to meet the Aboriginal students from the Queen Elizabeth School working on the same project, where they had an opportunity to learn about each other's cultures and view each other's videos. Together, the K'alemi Dene and Edmonton youth attended the Youth Day of the Dreamspeaker's Aboriginal film festival to learn more about film production.

Lizotte, Amy

Royal Roads University Yellowknife, NT amyruthlizotte@gmail.com

File Number: 12 410 921 **Licence No:** 15112

Region: IN, NS, SS **Location:** Yellowknife; Inuvik; Hay River

Backyard gardening - a viable food production solution for Yellowknife?

This community-based research project interviewed a cross section of local food growers to understand opportunities and challenges with producing local food, and also facilitated an online survey to assess food preference. This research was guided by the question: What would people buy if it were made available? Issues of agricultural management, sustainable development and food systems, and social strategies were examined to promote a slower, locally grown food system in Yellowknife. Data were also collected on economic profitability and the logistics of an urban farm business in Yellowknife. This research indicates Yellowknifers are ready for local food production. Over 99% said they would buy locally grown food if it was available, 88% would pay a premium for the food, and 96% said they would attend a weekly market. Salad mixes, and other high profit greens such as herbs, as well as vegetables like carrots, radishes and beans were identified in the top 60% of participant's responses. This data supports the feasibility of a small-scale garden business in Yellowknife. Also based on survey results, a sixth of an acre is currently available for commercial gardening in backyard spaces. However, over 7000 acres are potentially available under the municipality. Yellowknife is ready and willing to facilitate growth of an urban farming business - the city is just waiting for a creative entrepreneur to pave the way.

Luig, Thea

University of Alberta Fort McPherson, NT tluig@ualberta.ca

Region: GW **Location:** Fort McPherson

Collaborative research on community well-being

This project explores the role of volunteering, sewing, and human-land relationships in individual and community well-being. This project is in collaboration with the community of Fort McPherson. The researcher learned from Elders and others who are known for conducting positive lives and are respected as volunteers and seamstresses. In nineteen life-stories, twenty stories about strong Teetl'it Gwich'in of the past, and many informal interviews, people described what helped them through difficult times. Guided by Elders, the project is mainly concerned with documenting the history of the Peel River Alcohol Society, and the experiences of Elders who have volunteered in helping others by drawing on their own knowledge and experience. In many peoples' stories volunteering, sewing, and being out on the land are important to well-being. The stories relate what it is to experience oneself as a powerful and skillful person who is involved and needed in the community or on the land, and the importance of working for their own and others' well-being in the face of political and economic inequalities. All recordings were transcribed and transcripts verified by participants. Videos have been edited roughly and then approved by the storytellers.

Mair, Heather University of Waterloo Waterloo, ON hmair@uwaterloo.ca

File Number: 12 410 902 **Licence No:** 14971

Region: DC **Location:** Fort Simpson; Hay River

Exploring social support, sport participation, and rural women's health using Photovoice

This project sought to determine the nature of rural women's involvement in organized sport and recreation and to assess their perceptions of the influence of these activities on their individual and social health. In particular, the study: (1) examined the social lives and health of rural women within the contexts of curling and curling clubs; (2) explored the roles that these clubs play as community places for rural women; (3) utilized photovoice to help women photograph, document, and express their perspectives; (4) documented how these activities held different meanings for women across Canada; and (5) determined how health, sport, and recreation can be understood within the broader contexts of gender and rural community change. Specifically, researchers worked with members of the Hay River community in the Spring of 2012. Four women curlers were given cameras to take pictures to convey what the club meant to them. Follow-up interviews were conducted with the women (by phone) to gather deeper insight into the meaning of the club in their health and well-being. Research findings were shared with participants. The conclusions of the study suggest, curling clubs are significant community places that are deeply valued by women and girls. Specifically, clubs help women and girls: (1) establish new (and maintain) longstanding friendships; (2) increase physical exercise; (3) develop and improve curling expertise; (4) access opportunities for volunteering, mentoring, leadership and community engagement; and (5) share practical, emotional, and affirmational support (e.g. team members were frequently referred to as a "curling family"). The study demonstrates that curling clubs serve to enhance and sustain the physical, mental, and social wellbeing of rural women and girls.

Mitrovic, Inya

Royal Roads University Vancouver, BC inyamitrovic@hotmail.com

File Number: 12 410 917 **Licence No:** 15077

Region: IN, GW, SA, DC **Location:** Communities within each of the settlement regions and areas affected by the Mackenzie Gas Project

Addressing stakeholder interests through public consultation: A focus on natural resource exploration in the Canadian arctic

This research assessed the effectiveness of the public consultation processes utilized by the Mackenzie Gas Project's proponents in addressing and meeting stakeholder interests. Additionally, it examined how the requirements mandated by the National Energy Board affected the design of the proponents' public consultation processes and whether the use of mediation in public consultation could foster greater stakeholder outcome satisfaction. Understanding the gaps between the initial process design or intent of the proponent versus and the actual outcome satisfaction as experienced by the stakeholder is critical for determining the types of interests that proponents should focus on and emphasize during public consultation.

Pearce, Tristan University of Guelph Guelph, ON

Gueiph, ON

tristanpearce@gmail.com

File Number: 12 410 650 Licence No: 15139 (Multi-year licence)

Region: IN Location: Ulukhaktok

Inuit traditional knowledge for adapting to the health effects of climate change (IK-ADAPT)

This research is part of the Inuit Traditional Knowledge for Adapting to the Health Effects of Climate Change (IK-ADAPT). IK-ADAPT is a 3-year project that works closely with 6 communities across the Canadian Arctic (Ulukhaktok, Inuvik, Igloolik, Iqaluit, Rigolet, Nain) to identify how Inuit traditional knowledge can help enhance health in light of a rapidly changing climate. The first stage of this project was to learn from community members what ethical considerations must be taken when documenting and disseminating Inuit knowledge. In July 2012, researchers worked with a noted elder in Ulukhaktok to collect data for this first stage. Similar work will be conducted with community members in the other five communities. Taken together, this information will be used to develop protocol for documenting and disseminating Inuit traditional knowledge in the project. The project will continue for another two years and will focus on initiatives, decided in partnership with the community, to identify how Inuit traditional knowledge can help enhance health given a changing climate.

Poole, Nancy

BC Centre of Excellence for Women's Health Vancouver, BC wavelength@telus.net

Repairing the holes in the net: Responding to the mental health needs of northern homeless women

This research sought to involve decision makers in the mental health, housing and social sectors in improving the systemic response to northern women with mental health concerns who have unstable housing/are homeless, informed by research with women and service providers. The objectives were: (1) to gather the perspectives of northern women regarding the trajectory of their unstable housing/homelessness; mental health challenges and access to services; and ideas for what might make/have made a difference to prevent homelessness, mental health concerns and related issues like violence and substance use problems; (2) to involve service providers in identifying specific service improvements, including First Nations perspectives on mental health, that could remove barriers and enhance support for services for northern women with mental

health and related health, housing and social concerns, contribute to improvement in women's health, and prevent homelessness; (3) to summarize academic literature, web-based reports and existing circumpolar policy reviews which describe effective approaches to prevention and alleviation of homelessness among women with mental health, substance use and violencerelated concerns, and consider their relevance to guide approaches that would be effective in specific, northern, cultural contexts; (4) to utilize a community of practice (CoP) model to support health system decision makers in each of the three territories to collectively discuss: evidence arising from the literature search, ideas for expanding and improving policy and practice, factors affecting the implementation of such policy and knowledge translation strategies.

Sandlos, John

Memorial University of Newfoundland St. John's, NL isandlos@mun.ca

File Number: 12 410 847 **Licence No:** 15035 (Multi-year licence)

Location: Former Pine Point mine/townsite; Fort Region: SA, NS, SS

Resolution and Hay River: Giant and Con mine sites:

Yellowknife and Dettah: Déline; Port Radium mine/townsite

Abandoned mines in northern Canada: Historical consequences and mitigation of current impacts

The ongoing Abandoned Mines Project looks at how mineral development impacted the social life, economic prospects and local environments of Northern communities throughout the twentieth century, considering carefully the connections between social justice and environmental change that were engendered by historical mining practices in the region. This project is transitioning from fieldwork and data collection to the writing and production of results. The major fieldwork activity this year was conducted in the Pine Point area in July and August. Researchers conducted a mapping survey of the Pine Point area, ground truthing data from aerial photos and historical maps, while assessing current land cover in the area. A researcher also spent six weeks living in Fort Resolution, conducting interviews with elders to produce map biographies of former land use practices in the Pine Point area. Continued archival research on the history of Giant Mine at the NWT Archives at the Prince of Wales Northern Heritage Center was ongoing.

Simmons, Deborah

University of Manitoba Yellowknife, NT simmons@cc.umanitoba.ca

File Number: 12 410 678 **Licence No:** 15104 (Multi-year licence)

Region: SA Location: Community of Déline

Caribou and communities in the Sahtú Region

No summary was submitted for this licence. This project is not in compliance with licencing requirements.

Smart, Miles

Concordia University Kirkland, QB milessmrt@gmail.com

File Number: 12 410 936 Licence No: 15161 **Region:** SA **Location:** Norman Wells and Norman Wells Land
Corporation (NWLC); Yellowknife - Prince of Wales

Northern Heritage Centre

Implementing the future: Ethnography of a land corp

This research project was an anthropological analysis of the Sahtú Dene and Métis Land Claim focused on the community of Norman Wells and the Norman Wells Land Corporation. The central research themes were the negotiation and implementation of the land claim, related political changes, and resource development in the Sahtú. A focus was put on including the views and words of involved community members. The fieldwork was done over five weeks in November and December of 2012. The main fieldwork site was Norman Wells with a smaller amount done in Yellowknife. Research was based mainly on one-on-one interviews. Some further interviews have been held over the phone since the fieldwork period ended. Additionally, archival research was done relating to the research topic including publicly accessible documents, media, academic literature and a small amount of available Norman Wells Land Corporation archives. Interviewees have had the opportunity to review their responses to ensure they approve. The project is now in the writing stage with an anticipated completion of late 2013.

Svoboda, Michael

Arctic Borderlands Ecological Knowledge Co-op Whitehorse, YT michael.svoboda@ec.gc.ca

File Number: 12 410 811 Licence No: 15030 (Multi-year licence)

Region: IN, GW Location: Fort McPherson; Tsiigehtchic; Aklavik; Inuvik;

Tuktoyaktuk

Arctic borderlands ecological knowledge coop: Community-based ecological monitoring program

The Coop uses both local and scientific knowledge to monitor and assess environmental changes in the range of the Porcupine Caribou Herd and nearby coastal and marine areas. Interviews are conducted every year by community researchers. People share what they see and hear about fish, berries, caribou, unusual animal sightings, weather conditions, and other things while they are out on the land. In 2012, monitors conducted interviews with selected community members and entered data directly online. This improved the reporting time and data validation 'step,' which occurred at the March 2012. A Youth Gathering was also held at the same time in Inuvik, which focused on leadership, facilitation, and environmental project/ monitoring. The Coop website (www.taiga.net/coop) remains an important communication tool and contains past Gathering Proceedings, as well as the results from the community monitoring program.

Todd, Zoe

University of Aberdeen Edmonton, AB zoe.todd@gmail.com

File Number: 12 410 815 Licence No: 15011 (Multi-year licence)

Region: IN **Location:** Paulatuk; Yellowknife - Prince of Wales Northern
Heritage Centre; Inuvik - Inuvialuit Cultural Resource Centre

Lands, lakes and livelihoods: women's subsistence fishing in Paulatuk, NT

From January to October 2012, the researcher lived and worked in Paulatuuq in order to conduct research on people's relationships to fish, fishing and water. Throughout the spring, she

conducted a series of interviews on fishing in Paulatuuq, and also participated in a series of fishing trips with community members. Throughout June, July, August and September, the researcher was on the land fishing with community members in order to better understand the who, what, when, where, why and how of people's fishing activities. She travelled to lake, river and coastal areas with families in order to verify and contextualize the interview, oral history and archival data that were collected earlier in the year. Extensive archival research was undertaken at the Hudson's Bay Archives in Winnipeg in November 2011 and May 2012. At the current stage of analysis, this work indicates that fishing has played an important role in Paulatuuq life in the past, and continues to be an important activity that supports food security, intergenerational learning and is a crucial way in which many Paulatuuqmiut participate in land-based activities. Fish, fishing and water – particularly sensitive sites like the Hornaday River watershed – will be potentially heavily impacted by future resource development. For these reasons, it will be important for mining proponents to consider the impacts of their activities on fishing within the community.

Turcotte, Andre
Carleton University
Ottawa, ON
andre turcotte@carleton.ca

File Number: 12 410 933 **Licence No:** 15157

Region: DC, NS

Location: 3 Groups in Yellowknife - one with Dechinta

students that are attending a conference in Yellowknife, and 2 other possible groups with high school students from high

schools in Yellowknife and Fort Smith

Democracy talks

During the fall of last year researchers conducted 22 focus group conversations with young people across the country, to understand what things they were concerned about and how they felt about politics and political leaders in Canada. Four groups took place in Yellowknife and Fort Smith. Young people shared a negative view of politics; felt left out and felt that politicians didn't care about them. The level of knowledge about politics varied within each group – some understood a lot, where others knew almost nothing. The issues that concerned the participants were often local and personal, and they expressed a desire for a leader that cares about what they do.

Webb, Kernaghan

Ryerson University Toronto, ON kernaghan.webb@ryerson.ca

File Number: 12 410 905 **Licence No:** 14990

Region: NS, SS Location: Avalon Rare Metal's NECHALACHO Mine

project Thor Lake Property

Community engagement in the Canadian mining sector: Identifying best practice and testing stakeholder theory

The field research was undertaken August, 2012 to identify, explore, and examine the community engagement approach of Avalon Rare Metals, Inc. as part of the company's corporate social responsibility strategy, at their Thor Lake Nechalacho Mine project. The field research consisted of semi-structured interviews in an effort to elicit responses concerning the community engagement practices of Avalon. Work is now underway to prepare a report of this field research. When completed, the report will be provided to the community members.

Will, Alice Memorial University St. John's, NL alicewill@live.ca

File Number: 12 410 916 Licence No: 15079
Region: SS Location: Fort Smith

Integrating human dimensions research in Wood Buffalo National Park: Understanding attitudes, beliefs and values toward wood bison and bison management

The goal of this project was to gain an understanding of the perspectives of local people regarding bison disease and management in Wood Buffalo National Park (WBNP). With a social science approach, the theme of this research surrounds the human dimensions of conservation and the role that local attitudes play in the conservation of wood bison (Bison athabascae). Support and opposition for management decisions were explored using a quantitative survey conducted doorto-door. In addition, qualitative research in the form of focus groups was used to gain a deeper understanding of the various perspectives about this issue. Fieldwork was conducted in Fort Chipewyan, Fort Fitzgerald, and Fort Smith in 2012 over the course of seven weeks in June and July. The most evident conclusion from this study was that Aboriginal peoples not only wish to be seen as partners on paper, but wish to have an ongoing dialogue with WBNP resulting in a genuine sense of inclusiveness. The majority of participants held positive attitudes toward bison, despite the presence of disease. There were some differences between Aboriginal and non-Aboriginal residents, however, on average, there was agreement on how bison in the park should be managed. This study emphasizes the need for public participation beyond traditional public meetings and consultation sessions in the wildlife decision-making process in Canadian national parks. The hope of this research is to contribute knowledge that furthers work between park managers, local Aboriginal governments, and local people, while aiding in informed decisionmaking regarding the future of this threatened species. This research is the result of a collaborative effort between Memorial University, Wood Buffalo National Park, and the Aboriginal and non-Aboriginal people of Fort Chipewyan, Fort Fitzgerald, and Fort Smith.

Young, Michael

Royal Roads University Victoria, BC michael.young@royalroads.ca

File Number: 12 410 922 **Licence No:** 15126

Region: IN, GW **Location:** Inuvik and outlying communities in the Beaufort

Delta

Rural migration and homelessness in the north

The goal of this research is to assess the role that the gaps in mental health and addictions services in Inuvik and the surrounding region play in the production of homelessness. The majority of time spent on this project in 2012 involved the establishment of networks working with homeless persons with co-morbid disorders (addictions and mental health). Focus groups were held with service providers in Inuvik, Sachs Harbour, Tuktoyaktuk, Fort McPherson and Aklavik. The information collected in these focus groups was used to develop a questionnaire to be administered to homeless persons with addictions and mental health disorders at a later in the project. The "Quality of Life for Hard to House Individuals" (QoLHHI) was selected at this time to replace the SF36 as an instrument to measure quality of life for homeless participants. The QoLHHI will be administered following the focus groups with homeless persons. Next steps

involve the development of the focus group questionnaire, the identification of potential homeless participants and the scheduling of interviews.

TRADITIONAL KNOWLEDGE 2012

Alexie, Elaine University of Victoria Victoria, BC edalexie@uvic.ca

File Number: 12 410 928 **Licence No:** 15148

Region: GW **Location:** Fort McPherson

The limits of sovereignty: Practices of indigeneity among the Teetl'it Gwich'in

The research was conducted in September to October in Fort McPherson. Nine Teetl'it Gwich'in elders were interviewed as part of the project. Community members and elders interested in participating were recruited through local posters and radio announcements. Some interviews were took place at participant's bush camp and others were completed in the homes in the community. The location of the interview was based on the preference of each participant. On some occasions, with proper consent, interviews were conducted over multiple sessions. The interviews were very open ended and allowing the elders to talk freely as long as they wished. Close to 600 minutes of interview was collected. Analysis of the material is ongoing and expected to be completed in the Fall of 2013.

Balanoff, Helen

NWT Literacy Council Yellowknife, NT helen@nwtliteracy.ca

File Number: 12 410 617 Licence No: 15102 Location: Ulukhaktok

Pitquhiraluavut Puiglimiatavut (We will not forget our ways): Bringing home photographs of the Inuinnait Collection at the British Museum

The trip to the British Museum in London, England took place in April 2012 (the previously planned trip was postponed because of elder illness). Three Ulukhaktummiut, selected by the community, travelled with two elders and a researcher from Cambridge Bay, as well as the academic researcher, a museum curator and the principal investigator. The group spent four days in the British Museum stores examining the objects they had requested through the photographs. The discussions about the artifacts included: the skills involved, how they were made, and related language. The sessions were filmed and photographed. On returning to the communities, community members were invited to view the video footage and photographs and to provide

further input into the stories that were attached to the objects. As well, community members were asked which objects they would like to recreate in workshops, using the information derived from the visit. Two workshops were then held - one on making crimped drum dance slippers and the other on bow making. The bows were then used to hunt muskox. The materials (photos, film shorts, narratives) are now being uploaded on to a new website. The film footage is currently being edited (by community members) for a one hour TV documentary.

Benson, Kristi

Gwich'in Social & Cultural Institute San Clara, MB kbenson@learnnet.nt.ca

File Number: 12 410 697 Licence No: 15165 (Multi-year licence)

Region: GW **Location:** Aklavik; Fort McPherson; Tsiigehtchic; Inuvik

Building capacity and documenting traditional knowledge on species at risk in the Gwich'in Settlement Area 2012- 2014

The purpose of this ongoing research is to gather Gwich'in traditional knowledge of three species of special concern- wolverine, grizzly bear, and woodland caribou. The project team, made up by staff and contractors from the Gwich'in Social and Cultural Institute and Gwich'in Renewable Resources Board (GRRB), completed several important tasks in 2012-2013. First, a community steering committee meeting was held (November 2012). The committee was made up of a member from the Renewable Resource Council in each of Aklavik, Inuvik, Tsiigehtchic, and Fort McPherson. They reviewed the questions and other documents for the project team, and selected possible interviewees. Interviews about grizzly bears were held in December, with knowledgeable Gwich'in hunters, trappers, and elders. The interviews focussed on grizzly bears as a species-atrisk – even though grizzlies are not declining in the Gwich'in Settlement Area, they are considered at risk across Canada so having region-specific information is important for wildlife management. A report on grizzlies was prepared and community review sessions were held in Fort McPherson and Tsiigehtchic. The report had information from all the interviews, plus other information from previous studies as well. A draft final report was produced after the community review sessions. Upon review by the community steering committee in early 2014, the report will be available on the GRRB's website.

Brook, Julia

Queen's University Kingston, ON 7jeb@queensu.ca

File Number: 12 410 938 Licence No: 15171 Region: NS Location: Yellowknife

Perseverance in/through the arts: Life histories of northern indigenous artists

No summary was submitted for this licence. This project is not in compliance with licencing requirements.

Coedv. Bill

Aboriginal Affairs and Northern Development Canada Yellowknife, NT bill.coedy@aandc-aadnc.gc.ca

Region: IN Location: BAR-C Tununuk Point, Richards Island

Traditional oral history of land use at Tununuk Point

The Department of Aboriginal Affairs and Northern Development Canada and Imperial Oil have undertaken a number of environmental assessments at Tununuk/BAR C as part of their clean-up plan for the site. These studies have identified knowledge gaps in the history of human land use in the area that AANDC hopes to fill through oral history interviews. This project summarizes existing evidence from historic accounts and prior oral history interviews, and poses questions in order to learn more about how Inuvialuit used the area in the vicinity of Tununuk. Two groups of ancestral Inuvialuit hunted, fished and travelled in the Tununuk region. The territory of the Kuukpangmiut included Richards Island and the area east of the Mackenzie River further upstream into the northern part of the Mackenzie River Delta. The Kitigaaryungmiut lived directly to the east of the Kuukpangmiut. They spent most of the year hunting and fishing at the mouth of the East Channel of the Mackenzie River and along the shores of Qangmalig Bay. Artifacts found in Qangmaliq Bay archaeological sites show fishing using nets in open water and ice fishing in winter were important activities. Foreigners, mainly from Europe, began coming into the Inuvialuit area in the 1800s. Few Inuvialuit oral histories survive that tell of the coming of Europeans to this area. The collection of drawings on small pieces of wood produced by Inuvialuit who traded at Fort Anderson between 1861 and 1866 may be the earliest surviving record by Inuvialuit of Europeans. Several burials are present at Tununuk. Judging from appearances, the burials at Tununuk are quite old, and most likely from the pre-Christian period.

Fabijan, Michael

KAVIK-STANTEC Inc.

Inuvik, NT

michael.fabijan@kavik-stantec.com

File Number: 12 410 907 Licence No: 15017

Region: IN Location: Inuvik; Tuktoyaktuk

GNWT Inuvik to Tuktoyaktuk traditional knowledge / traditional land use study

No summary was submitted for this licence. This project is not in compliance with licencing requirements.

Gauthier, Brenda

University of Victoria Fort Smith, NT bgauthier@northwestel.net

File Number: 12 410 930 **Licence No:** 15152

Region: DC, SS **Location:** Fort Providence

A narrative of women's crafts, learning and cultural identity

No summary was submitted for this licence. This project is not in compliance with licencing requirements.

Gilday, Cindy

Institute for Circumpolar Health Research Yellowknife, NT cindy.gilday@ichr.ca

File Number: 12 410 925 Licence No: 15135 (Multi-year licence)

Region: NS Location: Yellowknife; Ndilo; Dettah; Cultural camp on

Mackenzie Island just past Akaitcho Bay on Great Slave

Lake

Climate change and emergency measures

Adaptation to climate change is critical for Northern peoples because it is affecting health and safety during northern travel. Unprecedented weather patterns now butt up against traditional knowledge (TK) about how to live and travel safely on the land. In this way it threatens the safety of First Nations and Inuit peoples of the Northwest Territories engaged in hunting, fishing and other land-based travel and activities informed by TK. Extreme weather variability is also proving to be a threat to those reliant on air travel. Air travel is the backbone of the emergency healthcare in the Northwest Territories (NT) as well as the only way to access some NT communities. Preventing of death and injury as a result of weather-related travel accidents is an important priority for First Nation and Inuit communities in the NT and a critical part of addressing the health impacts of climate change in Canada's north. The proposed project engaged northern youth to look at the intersection of climate change and health from the vantage point of its impact on TK and travel safety. It also promoted discussion between youth and their communities on the issues of emergency preparedness and injury prevention. Using community-based research methods and digital film as data collection tools, students discussed the themes of climate change with elders, climate change experts, and their community members to identify key lessons about the impacts of climate change on traditional lifestyles and travel in the north. As part of their investigations, students looked what the dangers/risks to health are and how they have impacted the community. Through reflection on guest lectures and class discussion, and personal interviews, students identified recommended actions to prevent and/or effectively respond to safety concerns that arise from unpredictable weather patterns caused by climate change. Key themes and lessons emerging from interviews form the basis of a short documentation film, created by the students with mentorship from local northern filmmakers.

Gordon, James Jr.

Aurora College Inuvik, NT james.gordon@myauroracollege.ca

File Number: 12 410 910 **Licence No:** 15039

Region: IN Location: Husky Lakes - all cabin owners

Husky Lakes cabin owner information survey

No summary was submitted for this licence. This project is not in compliance with licencing requirements.

Ireland, Margaret

Jean Marie River First Nation Jean Marie River, NT negotiations@jmfrn.ca

File Number: 12 410 883 **Licence No:** 15138

Region: DC Location: In and around Jean Marie River

Permafrost vulnerability assessment and landscape changes related to climate change in the Jean Marie River First Nation

No summary was submitted for this licence. This project is not in compliance with licencing requirements.

Jansen, Kelsey University of Alberta Edmonton, AB kljansen@ualberta.ca

File Number: 12 410 923 Licence No: 15130 (Multi-year licence)
Region: SS Location: Łutsel K'e; Artillery Lake

Denesoline traditional knowledge of landscape-caribou movement interactions with Łutsel K'e Dene First Nation

The purpose of this ongoing project is: (1) to collect and analyze traditional knowledge and ecological data of landscape-caribou interactions; and (2) to develop a set of community-based indicators to assist the community of Łutsel K'e in the monitoring of barren-ground caribou movements within their traditional territory. The work completed as of October 31, 2012 includes scoping and preliminary interviews in Łutsel K'e and at field sites at Artillery Lake conducted between July and September 2012. Only preliminary analysis has been conducted with no formal writing undertaken thus far. An informal update has been provided to the community in the form of a pamphlet summarizing the 2012 research season and training activities undertaken with Łutsel K'e youth at the research camp at Artillery Lake between September 5 and 11, 2012. This pamphlet is currently under review by the Łutsel K'e Dene Band and Wildlife Office; once approved copies will be disseminated within the community. The researchers planned to return to Łutsel K'e in 2013 to conduct some secondary and verification interviews and to present preliminary findings to Chief and Council and community members.

Lantz, Trevor University of Victoria Victoria, BC tlantz@uvic.ca

File Number: 12 410 906 **Licence No:** 15109 (Multi-year licence)

Region: IN Loc

Location: Husky Lakes; Hendrickson Island: Areas East and West of Tuktoyaktuk; The Mackenzie Pipeline Corridor (ISR); The Inuvik – Tuktoyaktuk road corridor (proposed); Aklavik Mountain Road (proposed); The Peel Plateau / Dempster Highway; in and around the communities of Aklavik, Inuvik, and Tuktoyaktuk, Tsiigehtchic and Fort

McPherson

Using Inuvialuit and Gwich'in observations to monitor environmental change in the Mackenzie Delta Region

The Mackenzie Delta Region (MDR) is a dynamic environment that is ecologically and culturally significant. This area is experiencing environmental changes that are expected to increase in magnitude with continued climate warming and additional anthropogenic stressors. In some areas, changes in land cover are occurring so rapidly that maintaining an accurate inventory is problematic. In this context of environmental change and uncertainty, there is critical need to draw on local knowledge and observations to inform decision-making. In the MDR, Inuvialuit hunters and trappers are in a unique position to assess ongoing changes in the regional environment and

to inventory cumulative impacts. Over the last three years, researchers at the University of Victoria and Aboriginal Affairs and Northern Development Canada (AANDC) have worked with the Hunter and Trapper Committees of Inuvik, Aklavik, and Tuktoyaktuk on several community-based monitoring initiatives. Since 2010, the research group has been developing and field testing a participatory monitoring protocol that uses participatory photography, video, and semi-structured interviews to record Inuvialuit observations. Observations of environmental conditions made during field outings with Inuvialuit experts and local youth are recorded using digital cameras, and handheld GPS units. Subsequently, digital photographs and video became the focus of photo elicitation interviews. The detailed narratives recorded in these interviews, along with geovideo entered referenced photos. and are into web-based (mapping.uvic.ca/mackenziedelta). During the winter of 2012, 12 Inuvialuit participants made observations using the participatory multimedia mapping (PMM) protocol. Monitoring activities took place on the land using snow machines and trucks. One multi-day monitoring and knowledge exchange camp was held in the Eastern Mackenzie Delta near Reindeer Station. To date, participants have focused their monitoring on environmental changes, including: shifts in wildlife and vegetation (range and distribution), drained lakes, thaw slumping, landslides, river bank erosion, increased run-off, increased overflows, changes in permafrost, and increasingly hazardous travel conditions. Observations also focused on damage to infrastructure (e.g. roads, cabins, camps, buildings), important sacred sites (e.g. traditionally used camps, travel routes, grave sites), and areas important for food harvesting. Expected impacts from proposed development (all season road to Tuktoyaktuk and the pipeline) were also discussed. This pilot project suggests that using PPM and web-based mapping to record local observations can make a contribution to local planning that will increase community resilience. Interviews with monitors and a range of potential map users suggest that the researchers' protocol and web map is an effective way to record and share observations of the regional environment. Elements of the protocol that contributed to its success include: the use of photography (the visual medium); the photo-interviews (story-telling); time spent on the land traveling and observing; and the paring of local youth and local experts. A monitoring program organized around continuous local observations that are linked to geo-referenced images (and other media) will significantly improve the capacity quickly detect environmental changes that impact northern communities. While webbased PPM should never replace direct community consultation, the research suggests that it can provide a resource that communities can use to share knowledge among themselves, across northern networks, and in meetings with researchers, regulators and decision-makers.

Moore. Kristin

Diavik Diamond Mine Inc. Yellowknife, NT kristin.moore@riotinto.com

File Number: 12 410 924 Licence No: 15131 Region: NS Location: Lac de Gras

Diavik palatability and tissue chemistry

This summary brings together results from traditional knowledge (TK) and scientific knowledge shared during a camp held near the Diavik Diamond Mine at Lac de Gras during the summer of 2012. These efforts were part of the Aquatic Effects Monitoring Program (AEMP) established by Diavik Diamond Mines Inc. with five Aboriginal parties to their Environmental Agreement: Kitikmeot Inuit Association; Łutsel K'e Dene First Nation; North Slave Métis Alliance; Tłįchǫ Government; and Yellowknives Dene First Nation. The primary objective of the 2012 program was to facilitate a two-way flow of information, resources, and knowledge between TK holders and scientists regarding the health of fish and water in Lac de Gras. Four key elements were the

focus of the AEMP: communications and engagement; fish palatability and texture studies; water quality and quantity studies; and video documentaries. Elders, youth and scientists collaborated to set nets and inspect overall fish health. Elders tasted a total of four fish that they baked, boiled, fried, and grilled. There were mostly positive descriptions based on the taste test of each fish. From this holistic, interconnected perspective, camp participants deduced that water quality was good by virtue of observing the health of surrounding or submerged vegetation, birds, wildlife, and fish; the shoreline; the presence/absence of surface foam and/or vegetation; clarity; movement; temperature; and taste. A 'tea test' was carried out whereby water samples were taken from Lac de Gras, boiled and then made into tea to evaluate the taste. In all cases, the taste of the water was said to be good. Water quality results from scientific results and TK support the same general conclusion that the water is still good in Lac de Gras. A video-documentary entitled "Five Ways, Two Days, One Camp" which was filmed and produced through a partnership of participating youth and a production crew.

Nash, Tyler Queen's University Kingston, ON tinash@hotmail.com

File Number: 12 402 872 **Licence No:** 15055

Region: NS Location: Along Baker Creek near Giant Mine

An investigation of arsenic speciation and toxicity in Baker Creek sediments from Giant Mine in the Northwest Territories, Canada

No summary was submitted for this licence. This project is not in compliance with licencing requirements.

Rice, Keren

University of Toronto Toronto, ON rice@chass.utoronto.ca

File Number: 12 410 678 Licence No: 15124 (Multi-year licence)

Region: SA Location: In and around Déline

Mapping, language and stories in Déline

This three year collaborative program develops an interdisciplinary approach to language documentation. As the community of Déline makes a transition to self-government, there has been increased interest in stories, song, and concepts of place in order to better understand what government means. Governance thus is one focal point of this research. Complementing this, the project involves development of an indigenous research methodology with respect to language research. The research explores variation, change and continuity in language, stories, song, and concepts of place as they relate to governance and land stewardship. The approach involves documentation with three groups of families from distinct traditional land use areas across generations, including archival and new materials, as well as dialogue with relatives from neighbouring communities with distinct dialects in order to understand the role of place of origin in variability. 2012 has been a time for transcription of existing narratives and development of the archiving system. Work will continue in 2013, with researchers staying in Déline for an extended period with community researchers and Dene language speakers.

Robinson, Andrew

Rescan Environmental Services Ltd.

Vancouver, BC arobinson@rescan.com

File Number: 12 410 912 **Licence No:** 15047

Region: SS **Location:** Hay River; Fort Resolution

Pine Point Socioeconomics and Traditional Use/Knowledge Study

No research was conducted under this licence in 2012.

Van Wyck, Peter Concordia University Montreal, QB pvanwyck@gmail.com

File Number: 12 410 939 **Licence No:** 15173 (Multi-year licence)

Region: IN, GW **Location:** Inuvik

Shifting stories, changing places: Transformational narratives of climate change in northwestern Canada and Alaska

Gwich'in and Inuvialuit roles in caribou stewardship go back thousands of years, and traditional values and laws shape how Gwich'in and Inuvialuit work to protect the caribou. At the same time, especially in Canada, gains made through land claims negotiations have helped Gwich'in and Inuvialuit have more say not just in managing the herd, but in educating the public and moving the indigenous perspective into more prominence in official channels, such as into the agenda of the Government of Canada. To keep the calving grounds campaign strong, it is important to celebrate victories, to maintain strong personal connections between people working on the campaign, and to renew involvement from new generations of people. Films, slideshows and other arts-based outreach projects have an important role to play in engaging people and giving a public face to the campaign. Arts-outreach projects on the calving grounds and other northern issues are more effective campaigning tools when they are made while consulting with northern partners from the ground up. New communications tools, such as social media, can be very helpful in increasing campaign outreach - but they are only effective if combined with a "human touch" where people are drawn into actively taking part in campaigns and having personal connections to them. "Journey North" is an educational website used in the Being Caribou project. School children can follow animal migrations in real time as information is posted to the web.

von Kuster, Jenica

MWH Canada, Inc. Calgary, AB jenica.vonkuster@mwhglobal.com

File Number: 12 410 929 **Licence No:** 15149

Region: SA, DC **Location:** Norman Wells; Tulít'a, Wrigley; Fort Simpson;

Jean Marie River; Trout Lake

Enbridge Pipelines (NW) Inc. Traditional Knowledge Study

No research was conducted under this licence in 2012. The research was transferred to ARI Licence 15243 for continuation into 2013.

Wesche, Sonia University of Ottawa Ottawa, ON

swesche@uottawa.ca

File Number: 12 410 934 **Licence No:** 15158

Region: DC **Location:** Fort Providence

Landscape scale flooding in the Great Slave Lake Plain: Expansion of lakes, flooding of wetlands and implications for bison habitat and local land users (Traditional knowledge study component)

This project is a partnership with the community of Fort Providence. Twelve interviews and a community workshop were carried out with local residents this past year. Local land users perceive a combination of different causes to be impacting water levels in the Great Slave Lake Plain, including climate change, increased beaver activity (due to reduced trapping), and the disruption of natural drainage patterns due to the Mackenzie Highway infrastructure. Impacts of changing water levels were noted in relation to vegetation and wildlife habitat, with knock-on effects for bison and moose. Snow and ice conditions are changing with warmer winters. These environmental impacts have implications for local residents, particularly in relation to traditional land use. Bison movement out of the sanctuary are impacting the big game bison hunt, which locals rely on for employment, and also highway safety (e.g. increased numbers of vehicle-bison collisions). Residents also experience significant limitations in terms of travel on the land (e.g. by snowmobile), where they must often delay or change the routing of harvesting trips due to unfavourable conditions. Various community-based adaptations are under discussion.

ARCHAEOLOGY 2012

Andrews, Tom

Prince of Wales Northern Heritage Centre

Region: SA Location: Tulít'a District

NWT ice patch monitoring project (2012)

Research in the high alpine was severely restricted this summer due to a record snowfall the previous winter. Even by mid-August, all locations the researchers regularly inspect were still deeply buried under a thick bed of winter snow. As a result, the regular survey was abandoned and the focus shift to helping Todd Kristensen with his excavations at nearby O'Grady Lake. The abundance of snow provided a new source of possible site locations. Specifically, a central ice patch site (KhTe-2) - that has produced a complete arrow dating to 400 ± 90 cal. Yr BP (see Andrews *et al.* 2012) - had completely melted out by 2011. Researchers removed an ice core from the site in 2007 that exhibited several stratified layers of caribou dung, the earliest dating to 3500 ± 110 cal. Yr BP, suggesting that it had been relatively stable for nearly four millennia (see Meulendyk *et al.* 2012). In August 2012, not only was it completely covered in snow again, two other patches, not visible at this time of year in previous years, were noted on lower slopes north and south of it, suggesting that these might be fossil patches exploited sometime in the ancient past. Researchers hope to explore these locations in a future year.

References: Andrews TD, MacKay G, and Andrew L. 2012. Archaeological Investigations of Alpine Ice Patches in the Selwyn Mountains, Northwest Territories. *Arctic* 65(5):1-21.

Meulendyk T, Moorman BJ, Andrews TD, and MacKay G. 2012. Morphology and Development of Ice Patches in Northwest Territories, Canada. *Arctic* 65(5):43-58.

Bussey, Jean

Points West Heritage Consulting Ltd.

Permit No: 2012-004 **Class:** 2

Region: NS Location: Akaitcho Region

Tibbitt to Contwoyto winter road project

In 2011, Jean Bussey of Points West Heritage Consulting Ltd. conducted archaeological investigations for the Joint Venture that operates the Tibbitt to Contwoyto Winter Road. This work was conducted through EBA Engineering Consultants Ltd. under Northwest Territories Archaeological Permit 2012-004. The Tibbitt to Contwoyto winter road runs from the south end of Tibbitt Lake near Yellowknife to almost the north end of Contwoyto Lake in Nunavut. Until about six years ago, the full length of this ice road was utilized every winter, but most years it now only extends as far as Lac de Gras due a lack of mining activity further north. In previous years, a

number of archaeological sites located near the winter road or its associated developments (e.g. gravel pits and camps) were marked by stakes to ensure avoidance during winter activities. Monitoring of the protected archaeological sites is undertaken every year or two. No monitoring was conducted in 2012. Instead an archaeological site discovered within a proposed gravel source in 2011 was tested and collected. Investigations in 2012 indicated that KkNx-16 consisted of three small localities containing relatively sparse archaeological material. The site is located on esker deposits on the north side of a lake locally known as Sandridge Lake. This lake is part of the winter road route and there are numerous sites recorded on this well-defined esker. KkNx-16 is on south facing slope on the north side of Sandridge Lake. It consists of three localities with quartz flakes visible on the surface. Locality 1 was characterized by a surface scattering of approximately 40 specimens of quartz including white, grey and clear. They extended over an area about 10 m by 10 m; the specimens were sparsely scattered, likely as a result of slope wash. No artifacts were recovered during subsurface testing at Locality 1. All surface specimens were collected using a 2 m by 2 m grid. Locality 2 at KkNx-16 was also characterized by a surface scattering of about 40 specimen of quartz, but the artifacts were limited to white and grey coloured materials. The majority of the artifacts were scattered across an area approximately 6 m by 8 m and were collected using a 2 m by 2 m grid consisting of 12 units. These artifacts are more likely in their original provenience since they are on level terrain characterized by exposed rock. No artifacts were recovered during subsurface testing. Locality 3, situated between the other two localities, contained fewer than 20 specimens of white quartz, most of which are chunky. The white quartz specimens were found on gentle slope in an area about 3 m by 3 m and were collected by measuring from a central datum. No artifacts were encountered during subsurface testing at Locality 3. The lack of artifacts beneath the surface suggests that KkNx-16 consists of three small surface scatters. White quartz is the dominant material, but grey and clear quartz are also present. In addition, in 2011, three specimens of a dark grey siltstone were collected downslope from localities 1 and 2.

Bussey, Jean

Points West Heritage Consulting Ltd.

Region: NS Location: Akaitcho Region

Gahcho Kué project

Points West Heritage Consulting Ltd. (Points West) conducted archaeological investigations for De Beers Canada Inc. (De Beers) at Kennady Lake, the location of the proposed Gahcho Kué Project. Kennady Lake is situated approximately 280 km northeast of Yellowknife and 140 km north of Łutsel K'e. Jean Bussey directed the investigations under Class 2 Northwest Territories Archaeological Permit 2012-003. She was assisted by Brenda Michel of the ŁutselK'e Dene First Nation. Points West previously conducted work at Kennady Lake for De Beers in 2004, 2005, 2006, 2007 and 2010. The objectives of the 2012 field investigation included archaeological potential assessment and limited ground reconnaissance. These investigations were prompted by minor revisions to the original Project footprint – the identification of a pipeline that would divert water from Kennady Lake to a small lake to the east and the identification of possible dykes. No new archaeological sites were discovered and low archaeological potential is identified in association with the pipeline and the small eastern lake. No new archaeological sites were found near the proposed dykes, but previously recorded sites are located nearby. In addition, there is archaeological potential at landforms near some of the potential dyke locations. If construction is proposed, locations with archaeological sites or with archaeological potential will require additional investigation - either site survey or site protection/mitigation. Using techniques employed along the Tibbitt to Contwoyto Winter Road, three sites along the winter access

between Mackay and Kennady lakes were marked to assist in protecting them during winter use. Wooden survey markers were installed at KkNq-6, KkNq-10 and KkNq-28. The markers were sprayed with fluorescent orange paint to make them more visible when there is snow cover. These locations will be monitored in the summer after each winter of use. De Beers sponsored workshops that were held at Kennady Lake during the summer and early fall of 2012; one was an archaeological workshop. Representatives of six First Nations groups attended: Deninu Kue First Nation, Łutsel K'e Dene First Nation, North Slave Métis Alliance, Northwest Territories Métis Nation, Tłycho Government and Yellowknives Dene First Nation. The archaeology workshop included a demonstration of how a lithic scatter is created, as well as a display of some of the artifacts collected from the Project area. Other workshops included tours of the project footprint via boat, a wildlife monitoring station, and aerial reconnaissance of the Kennady Lake watershed and specific proposed developments. An archaeological management plan was prepared, reviewed by the Prince of Wales Northern Heritage Centre, revised, and submitted in October 2012. The plan includes archaeological measures that will be undertaken if the Gahcho Kué Project is approved. These measures include surface collection, subsurface excavation, monitoring, and site protection and will be updated whenever there are changes in the development plans.

de Guzman, Margarita Altamira Consulting Ltd.

Permit No: 2012-017 **Class:** 2

Region: NS Location: Akaitcho Region

Archaeological overview Giant Mine remediation

In October of 2012, Altamira Consulting Ltd. conducted an initial archaeological field survey of the Giant Mine Remediation Project area. The survey is part of an Archaeological overview of the Giant Mine area and was conducted in advance of the Giant Mine Remediation Project. Giant Mine is an abandoned gold mine located within the City of Yellowknife. The lease boundary covers 872 hectares and encompasses a number of ponds and small lakes, including Baker Creek, Pocket Lake, Trapper Lake and a portion of Yellowknife Bay (Great Slave Lake). Close to 60 years of gold mining has resulted in a massive environmental liability. The proposed remediation plan involves reclamation of the abandoned gold mine and the containment and immobilization of 237,000 tonnes of arsenic trioxide, a by-product of the gold production process. Included in the contamination is an estimated 325,000 m³ of soils, as well as various buildings and mine facilities. The property is now Commissioner's Land and is administered by the Department of Municipal & Community Affairs. The archaeological overview assessment, which consisted solely of pedestrian reconnaissance, was directed at assessing lands within the Giant Mine Lease Boundary for archaeological potential. This included identifying areas of disturbance and determining areas of potential use for remediation, as well as determining if any previously recorded sites remained in the area. During the overview, three previously recorded heritage sites were revisited and eight areas were identified as having heritage potential; several new sites were also recorded. The results of these investigations underline the need, and provide justification, for a full Heritage Resource Impact Assessment of the project area prior to the initiation of field remediation operations.

Jankuta, Kimberly Altamira Consulting Ltd.

Region: NS Location: Akaitcho Region

Ingraham Trail realignment

In July of 2012, Altamira Consulting Ltd. conducted an archaeological field survey of the Ingraham Trail located near Yellowknife, NT. The survey is part of an Archaeological Impact Assessment for the Ingraham Trial Realignment Project. The project personnel included Kimberly Jankuta and Jode MacKay. Originally constructed in the mid-1960s the Ingraham Road is part of Highway 4 that extends from Yellowknife approximately 70 km east to Tibbit Lake. The proposed realignment moves the southernmost leg of the road to the west to detour around the Giant Mine area. The proposed realignment will move the southern access point to the west from 48th street onto Highway 3. The realignment will connect with the existing road several kilometers to the north, near the turn off towards Vee Lake. The realignment will detour traffic around the Giant Mine area in anticipation of the remediation of the abandoned gold mine. The archaeological survey, which consisted of pedestrian reconnaissance and sub-surface shovel tests, was directed at assessing archaeological potential within the proposed Ingraham Trail Realignment right-of-way. This included shovel testing in areas thought to have potential and identifying areas of disturbance, both modern and historic. No archaeological sites were found during the survey. Two rectangular shaped rock features were noted, which may represent the use of rocks to hold down a tent on the rocky landscape. However, no hearths or cultural materials were found in association. A known historic site was revisited during the survey; the historic site is located adjacent to the project area. It represents the location of a 1935 Geological Survey of Canada field crew's discovery of gold. The team led by Norman Jennejohn discovered a quartz vein with visible gold. The site itself consists of a large trench cut through the bedrock.

Kristensen, Todd University of Alberta

Permit No: 2012-007 **Class:** 2

Region: SA Location: Tulít'a District

O'Grady Lake archaeology and ice patch monitoring project

A collaborative team from the University of Alberta, the Prince of Wales Northern Heritage Centre, and the Tulit'a Dene Band, visited O'Grady Lake and several ice patches in the Selwyn Mountains of the Northwest Territories from late July to mid-August. Crew members included Glen MacKay, Leon Andrew, Mike Donnelly, Tom Andrews, and Todd Kristensen. The goals were to monitor ice patches where ancient artifacts have been found and to find new archaeological sites around O'Grady Lake. In past years, ice patches have yielded well-preserved weapons and technologies left behind by people hunting caribou in high alpine areas. It is hoped that archaeological excavations around neighbouring O'Grady Lake will reveal more about the relationship between this alpine caribou hunting and lowland camps. Eight new archaeological sites were discovered around O'Grady Lake during 2012 fieldwork. Most of these sites consist of stone tools and debris from tool production. No new artifacts were discovered during ice patch monitoring due to heavy winter snows that expanded the extent of many patches. Small scale excavation units were dug at two of the O'Grady Lake archaeology sites in order to learn about possible dwellings and activity areas. One site produced fire cracked rock from boiling food while another yielded a deep cultural occupation below a layer of volcanic ash deposited 1250 years ago. Radiocarbon collected from this deep deposit will be tested to determine when O'Grady Lake was first used by pre-contact people. Both sites will be returned to in 2013 for more excavations.

MacKay, Glen

Prince of Wales Northern Heritage Centre

Region: NS Location: Akaitcho Region

Yellowknife Bay archaeology project

In 2012, archaeologists from the Prince of Wales Northern Heritage Centre began an archaeological survey of the Yellowknife Bay area in collaboration with the Yellowknives Dene First Nation. The goal of the project is to record archaeological sites in and around Yellowknife Bay, which will facilitate their protection when land use activities are proposed in the area. This summer researchers focused their survey efforts between the mouth of the Yellowknife River and Tartan Rapids. They recorded archaeological evidence of past land uses in this area ranging from pre-contact stone tool scatters to campsites associated with historic mineral exploration activities. The continued survey of the Yellowknife area is planned for future summers.

MacKay, Glen

Prince of Wales Northern Heritage Centre

Region: DC Location: Dehcho Region

Five Lakes archaeology project

The Prince of Wales Northern Heritage Centre (PWNHC) continued a community archaeology project in partnership with the Jean Marie River First Nation in 2012. The goal of the project was to document cultural values in the Łue Túé Sųlái Candidate Cultural Conservation Area, which is being considered for protection through the NWT Protected Areas Strategy. According to the oral traditions of the people of Jean Marie River, these small fish lakes were important winter harvesting areas, where fish caught through the ice and small game provided important staples for the winter months. In 2012, archaeologists from the PWNHC visited all five lakes in the Łue Túé Sųlái Candidate Cultural Conservation Area with elders from Jean Marie River, and recorded 14 new archaeological sites. In September, archaeologists participated in the community culture camp at Tthets'éhk'e (McGill Lake), and were able to involve students from Henry Ekali school in the work. Most of the archaeological sites recorded this summer consist of pre contact lithic scatters. The most significant sites were found at the outlet of Tthets'éhk'e, where archaeologists found evidence of multiple pre contact occupations. They hope to return to these sites and others in the Łue Túé Sųlái Candidate Cultural Conservation Area in future years to conduct more detailed excavations.

Murphy, Brent

Prince of Wales Northern Heritage Centre

Region: DC **Location:** Dehcho Settlement Area

Archaeological impact assessment - Canada Zinc Corporation proposed access road alignment changes, Nahanni Butte, NT

During September of 2012, Golder Associates Ltd. conducted an Archaeological Impact Assessment under NWT Permit 2012-016 on behalf of Canadian Zinc Corporation of changes to their Prairie Creek Mine Access Road Alignment near Nahanni Butte, NT. The study included the assessment of the proposed Nahanni Range Alternative (56.2 km) winter road. The alternative road travels from just northeast of the community of Nahanni Butte north along the Nahanni Front Range to Grainger Gap where it meets up with the existing winter road. The current winter road was used by the past owner of the mine in the 1980s and was subject to an archaeological assessment in 2009. The objectives of the Archaeological Impact Assessment were to identify, record and assess heritage resources that might be impacted by the proposed winter road and to devise appropriate mitigation strategies should any be found in conflict with the proposed winter road alignment. The archaeological sites may include previously unrecorded sites within or

adjacent to the proposed right of way, temporary workspace and/or borrow areas, if relevant. The field assessment was planned in conjunction with Elders and community members in Nahanni Butte prior to the field studies. Although the meetings were informal, advice and information from several community members and Elders was obtained that aided in the design of the archaeological field program. The field studies included the participation of Wilbert Antoine from Canadian Zinc Corporation and Peter Marcellais and Elder Leon Konisenta from the community of Nahanni Butte who assisted during the field program and provided advice on the cultural significance of the landscape traversed during the investigation. The field studies also included low and slow helicopter overflight and some pedestrian survey. The entire project right-of-way was examined from the air and pedestrian survey was focused on the proposed crossing of the Liard River. The results of the assessment were that no new archaeological sites were recorded or revisited; however, two traditional land use locations, both trails, were noted but not officially recorded as they do not meet some or all of the criteria required to be designated as an archaeological site under the Northwest Territories Archaeological Sites Regulations.

Ng, Tommy

Bison Historical Services Ltd.

Region: SA Location: Tulít'a District

Slater River winter 2012-2013

On behalf of MWH Canada, Inc., acting as agent for Husky Energy, Bison Historical Services Ltd. conducted a heritage resource survey for the proposed Slater River Winter 2012 - 2013 Program within the exploration licence area EL462 and 463 within the Tulít'a District of the Sahtú Region. The exploration licence area is located within the Mackenzie Plain, which is southeast of the Town of Norman Wells and on the south side of the Mackenzie River. The proposed Slater River Winter 2012 - 2013 Program includes an upgrade to an existing eight metre wide winter road and a four metre wide source 126 seismic line into 20 m wide all-weather road. This includes four areas of new road cut along the source 126 seismic line. Associated with the road construction are a base camp/storage area, an airstrip and 13 quarry sites. Additionally, the proponent requested a heritage assessment of 32 future developments, which included eight petroleum wellsites (four horizontal and four vertical) and 24 groundwater monitoring pads. Most of these developments will be located along the all-weather road. This heritage resource survey is an assessment of 49 petroleum associated developments and includes eight petroleum well sites, 24 monitoring water wellsites, 13 quarries, an airstrip, a base camp and two all-weather roads (19 specific landforms along the two roads were assess for heritage resource potential). A total of 66 locales were assessed for heritage resources. Personnel of Bison Historical Services Ltd., based in Calgary, Alberta, assisted by a wildlife monitor and local advisor from the Tulít'a Renewable Resources Council, conducted the heritage resources survey from September 5 to 16, 2012. The survey was based out of Norman Wells and included helicopter overflight and pedestrian reconnaissance accompanied by the excavation of shovel tests within the proposed project area. Pedestrian investigations focused on areas that were assessed to have high to moderate potential for new heritage resources within the confines of the proposed project area. A total of 432 shovel tests were conducted and all yielded negative results. It is recommended that the proposed Slater River Winter 2012 - 2013 Program be granted approval to proceed with development. This recommendation is subject to the approval from the Government of Northwest Territories through the Prince of Wales Northern Heritage Centre.

Ng, Tommy

Bison Historical Services Ltd.

Region: SA Location: Tulít'a District

Chinook project

On behalf of MWH Canada, Inc., acting as agent for ConocoPhillips Canada (CPC) Resources Corporation, Bison Historical Services Ltd. conducted a heritage resource survey for the proposed ConocoPhillips Chinook Drilling Program within the exploration licence area of EL470 within the Tulít'a District of the Sahtú Region. The exploration licence area is located within the Mackenzie Plain, which is southeast of the Town of Norman Wells and on the south side of the Mackenzie River. During the 2012/2013 winter season, CPC plans to use an ice bridge access owned by Husky Energy in early winter, drill two vertical petroleum wells, develop access to a storage and staging area on the shores of the Mackenzie River, and construct an ice bridge across to Norman Wells for light truck traffic. The Norman Wells ice bridge access would be CPC's main access for subsequent years. Additionally, CPC wants to assess two petroleum well locations planned for the 2014 winter season. The heritage resource survey for ConocoPhillips was conducted at 14 locales, which included two 2012/2013 vertical petroleum wells, the two 2014 petroleum wells, base camp, construction camp, staging area, storage area and five ground water wells. All of these developments are located next to an existing access route. Also included is a heritage resource survey of the proposed new cut (D - E) that will be connecting two segments of the existing access route. Personnel of Bison Historical Services Ltd., Calgary, Alberta, assisted by a wildlife monitor and local advisor from the Tulít'a Renewable Resources Council. The heritage resource survey was conducted from August 8 to 11, 2012 and included a helicopter overflight to assess the archaeological potential of the entire proposed project area. Additionally, pedestrian reconnaissance accompanied by shovel tests were conducted in areas deemed to have moderate to high archaeological potential within the proposed project area. A total of 159 shovel tests were conducted and no artifacts were found. No new heritage resource sites were identify and there are no previously documented heritage resource sites within the general vicinity of the proposed project area. It is recommended that the proposed ConocoPhillips Chinook Drilling Program be granted approval to proceed with development. This recommendation is subject to the approval of the Government of Northwest Territories through the Prince of Wales Northern Heritage Centre.

Ng, Tommy

Bison Historical Services Ltd.

Region: SA Location: Tulít'a District

MGM East Mackay project

On behalf of MGM Energy Corp., Bison Historical Services Ltd. conducted a heritage resource survey for the proposed MGM East Mackay Two Well Horizontal Project within the exploration licence area of EL466 within the Tulít'a District of the Sahtú Region. The exploration licence area is located within the Mackenzie Plain, which is south of the Hamlet of Tulít'a and on the south side of the Mackenzie River. During the 2012/2013 winter season, MGM Energy Corp. plans to drill one vertical wellsite and one horizontal wellsite, both of which are 300 m apart from each other. Additionally, MGM plan to construct two future petroleum horizontal wellsites in the general area. Despite what is shown in Figure 1, the proposed project does not include the staging area and construction campsite; they will be constructed on the north side of the Mackenzie River. The heritage resource survey for the MGM East Mackay Two Well Horizontal Project was conducted at eight locales, which included the two 2012/2013 petroleum (vertical and horizontal) wellsites and the general locales for two additional future (2014) horizontal well sites. Also included is a heritage resource survey of four new access cuts crossing bodies of water and connecting

separate segments of the existing access route. All of these developments are located along an existing access route. Personnel of Bison Historical Services Ltd., based in Calgary, Alberta, assisted by a wildlife monitor and local advisor from the Tulít'a Renewable Resources Council, conducted the heritage resources survey from August 20 to 22, 2012. The heritage resource survey was based out of Norman Wells and included a helicopter overflight and a pedestrian reconnaissance accompanied by the excavation of shovel tests within the proposed project area. Pedestrian investigations focused on areas that were appraised to have high to moderate potential for new heritage resources within the confines of the proposed project area. A total of 105 shovel tests were conducted and all were negative of artifacts. No new heritage resource sites were identify and there are no previously documented heritage resource sites within the general vicinity of the proposed project area. It is recommended that the proposed MGM East Mackay Two Well Horizontal Project be granted approval to proceed with development. Regarding the North and South Areas of Interest, the proponent assured the permit holder that the two future horizontal petroleum wellsites will be placed 200 m away from the drainages and lakes. Landforms that are considered to be locales of high heritage resource potential within these areas of interest. This recommendation is subject to the approval from the Government of Northwest Territories through the Prince of Wales Northern Heritage Centre.

Prager, Gabriella

Points West Heritage Consulting Ltd.

Region: NS Location: Akaitcho Region

Nechalacho Rare Earth Element project

The Nechalacho Rare Earth Metals project is located on the north side of the east arm of Great Slave Lake, approximately 95 km southeast of Yellowknife. The mine development is focused around Thor Lake, about 4 km due north of Great Slave Lake, with a dock on the GSL shore; a marshalling yard is proposed at Pine Point on the south side of Great Slave Lake. The 2012 Points West Heritage Consulting Ltd. archaeological team consisted of Gabriella Prager (Project Director), Carol Rushworth of Points West, and a local person from each of the three closest communities: they were Fred Sangris from Dettah, Gabriel Enzoe from Łutsel K'e and Victor Mandeville from Deninu Kue (Fort Resolution). For the Pine Point work, Wilfred Beaulieu represented the Fort Resolution Métis Council. The 2012 archaeological inventory survey of the mine area covered gaps that remained after the initial 2011 surveys, that is, where project components were revised or boundaries were not accurately identifiable. All mine related facilities proposed on the north side of Great Slave Lake, as well as the proposed marshalling yard at Pine Point were examined by pedestrian transects sufficient to provide good visual coverage and subsurface testing in selected areas. No archaeological remains was found in the Pine Point marshalling area due to extensive past disturbance and ongoing use, but an interesting structure of driftwood logs was found on the shore just outside the identified yard that could have been a hunting blind. Seven previously recorded sites (KaPb-4, KaPb-6 to KaPb-11, inclusive) in the north project area were subjected to systematic data recovery comprising detailed plan mapping, careful surface inspection of surrounding area, extensive photography, and subsurface testing of at least two tests units at each site where there was soil. No artifacts or additional features were uncovered during these mitigation actions. Because several finished tools had been found at KaPb-4 in 1988, this year archaeologists conducted a very careful surface inspection of the beach and all surface exposures, and extensive shovel testing. Although no additional artifacts were found, this is a very large site area and the vegetation is thick; therefore, artifacts could still be present. Two new sites were recorded at the Great Slave Lake north shore. One is a historic tipi style camp site that contains several prepared poles, a concentration of cut spruce boughs, and

a hearth. It is adjacent to a fairly fresh looking skid trail that extends from the lakeshore to the road. This site was thoroughly recorded and three units were excavated. Nothing was found in the units except a metal snap which may be intrusive because it looks quite new. The second site contains stone features situated on the southwest end of the lake point a short distance south of the existing road. These features consist of a rock pile that may have been used as a cache, a propped large, flat rock that could represent a possible trap or a platform for some purpose such as a table, and a hearth, all on bare bedrock. The site terrain and features were mapped to scale and extensively photographed. All known sites are now considered fully recorded.

Ross, Julie

Prince of Wales Northern Heritage Centre

Region: SA Location: Tłįcho Settlement Area

Archaeological survey - Indore, Hottah Mines and surrounding areas

In June of 2012, Julie Ross of Golder Associates Ltd. and Dolphous Apples from Gamètì, conducted an Archaeological Impact Assessment (AIA) in the vicinity of the Indore and Hottah mine sites along the shores of Beaverlodge and Hottah Lake north of Gamètì for Aboriginal Affairs Northern Development Canada (AANDC). The Euro-Canadian sites consist of uranium exploration camps and the two mine sites. Both site types have resulted in contaminated waste being distributed along the landscape and AANDC's intention is to remediate these sites. Eleven previously unrecorded archaeological sites were recorded and six previously recorded sites were revisited. Many of the Euro-Canadian sites were used by Tłįcho after their abandonment. The Euro-Canadian sites were mapped and photographic documentation was conducted. The Tłįcho sites that were recorded include 3 fish caches, 2 hunting blinds and 2 camp sites.

Seip, Lisa

Rescan Environmental Services Ltd.

Region: NS **Location:** Akaitcho Region, North Slave

Courageous Lake

In June and September of 2012, Rescan Environmental Services Ltd. conducted archaeological baseline studies for Seabridge Gold Inc.'s Courageous Lake Project under Northwest Territories Class #2 Archaeologist's Permit 2012-002. These investigations were a continuation of baseline studies conducted in 2010 and 2011, under Northwest Territories Class #2 Archaeologist's Permits 2010-015 and 2011-006, respectively. Lisa Seip directed the fieldwork and was assisted by archaeologists Daniel Walker, Vanessa Neuman, and Michael Campbell, also of Rescan Environmental Services Ltd., and by First Nations assistants Ernie Sangris of the Yellowknives Dene, and Darcy Zoe and Charlie Tatzia of the Tłjcho First Nation. Investigations included the assessment of proposed drill pad locations to the north of Courageous Lake, and surrounding Walsh and Saucer Lakes, and proposed project infrastructure to the south of Courageous Lake. The objective of the investigation was to identify sites that would potentially be impacted by newly proposed infrastructure and drill pad locations. Pedestrian surveys were conducted, focusing on areas considered to have high archaeological potential; subsurface testing was conducted in areas with adequate soil deposition. Examinations resulted in the identification of 54 archaeological sites, including 44 lithic sites, 6 rock feature sites, and 4 historical sites. Twelve archaeological sites with diagnostic artifacts were identified, including two Shield Archaic tradition sites, five Arctic Small Tool tradition sites, and five Taltheilei tradition sites; all of the attributed cultural affiliations are tentative. One previously recorded site, LaNv-20, was revisited. Avoidance

is the preferred management recommendation for all sites. If avoidance is not possible, then systematic data recovery is recommended. As the project is currently in the design phase, no impacts are anticipated this year. Additional archaeological studies are planned for 2013.

Thomson, J. Callum

Thompson Heritage Consultants

Region: NS Location: North Slave Region

Preliminary archaeological inventory and assessment of Peregrine Diamonds Lac de Gras West and East Claim Blocks

No summary was submitted for this permit.

Walker. Daniel

Rescan Environmental Services Ltd.

Region: SS Location: Akaitcho Region

Pine Point Project

No summary was submitted for this permit.

Youell. Alan

Stantec Consulting Ltd.

Region: IN Location: Inuvialuit Settlement Area

Inuvik to Tuktoyaktuk Highway borrow source investigations program

On behalf of the Department of Transport, Government of the Northwest Territories, Kavik-Stantec Inc. conducted an archaeological impact assessment of the Inuvik to Tuktoyaktuk Highway Borrow Source. The specific purpose of the archaeological component of the Inuvik to Tuktovaktuk Highway Borrow Source Investigations Program was to identify archaeological. historical, palaeontological and traditional land use sites at the proposed gravel borrow source locations. These borrow source locations are situated within the Inuvialuit Settlement Region east of the east channel of the Mackenzie River and west of Eskimo (Husky) Lakes. Investigation of the developments was conducted under Northwest Territories Class 2 Archaeologists Permit #2012-012. To conduct the assessment, archaeologist Alan Youell and wildlife monitor Tommy Chicksi of Inuvik conducted a field reconnaissance of the proposed development areas. The field reconnaissance consisted of a pedestrian traverse and intensive surface examination to determine the presence of unrecorded archaeological or cultural sites. Shovel tests were excavated in areas with the potential for buried cultural materials. The areas investigated during the archaeological assessment of Tuktoyaktuk Highway Borrow Source Investigations Program included the assessment of borrow sources 2.45, 170, 172, 173/305, 307, 314/325 and 312, no archaeological, historical or palaeontological sites were located and no previously recorded sites were revisited. However, two land use sites (modern campsites) and a section of the Jimmy Lake to Eskimo (Husky) Lakes trail were recorded. Based on the results of this assessment, there are no outstanding conflicts between archaeological, historical or palaeontological sites and the potential gravel borrow sources 2.45, 170, 172, 173/305, 307, 314/325 and 312. It is recommended that any impact to the land use sites should be mitigated through consultation with the communities involved.

WILDLIFE 2012

Armstrong, Terry

Environment and Natural Resources South Slave terry armstrong@gov.nt.ca

Permit No: 5770 Species Studied: Wood bison

Region: SS, NS **Location:** North of the Mackenzie River to Behchokò; Great

Slave Lake to the Horn Plateau

Mackenzie wood bison population census - 2012

The objective of this study was to estimate the size of the Mackenzie wood bison population with a coefficient of variation of 20% or less.

Barclay, Robert

University of Calgary barclay@ucalgary.ca

Permit No: 500046 Species Studied: Bats

Region: SS Location: In and around Fort Smith and Kakisa

Bat surveys and investigation of northern adaptations of little brown bats

The objectives of this study were: 1) to gather baseline information for northern bat species' presence and little brown bat population structure and health, to assist in monitoring for white nose syndrome in Wood Buffalo National Park and the Northwest Territories; 2) to describe the annual cycle of little brown bats in the South Slave region, including timing of reproduction and hibernation; 3) to investigate foraging behaviours of little brown bats in response to short summer nights at northern latitudes; and 4) to compare the diet of two sympatric *Myotis* species at northern latitudes.

Bidwell, Mark

Canadian Wildlife Service mark.bidwell@ec.gc.ca

Permit No: 500051 **Species Studied:** Whooping cranes

Region: SS Location: In and around Wood Buffalo National Park

Whooping crane ecology and rehabilitation

The objective of this population monitoring study was to monitor and understand the breeding ecology of whooping cranes in Wood Buffalo National Park and the surrounding area. The

objective of the three-year banding program was to gain a better understanding of whooping crane ecology and behaviour during the annual cycle with the use of GPS and radio-telemetry technology. Specifically to: a) gain a better understanding of stopover areas, habitat use patterns, and factors influencing habitat use at different spatial and temporal scales, b) define a current migratory route to compare to previous route models and determine environmental and anthropogenic factors that influence migratory behavior, c) identify causes, locations, and conditions of actual or potential mortality, and, d) expand current knowledge of winter and breeding ground use through high resolution GPS technology.

Bourn, Stephen

Diavik Dimond Mines Inc.

Permit No: 6889 Species Studied: Wildlife

Region: NS Location: Diavik wildlife study area, centered on Lac de

Gras

Wildlife monitoring program for the Diavik Diamond Mine

The objectives of this study were: 1) to verify the accuracy of the predicted effects determined in the Environment Effects Report (Wildlife 1998) and the Comprehensive Study Report (June 1998); and 2) ensure that management and mitigation measures for wildlife and wildlife habitat are effective in preventing significant adverse impacts to wildlife.

Carrière, Suzanne

Environment and Natural Resources - Wildlife suzanne carriere@gov.nt.ca

Permit No: 500063 **Species Studied:** Mice, voles, lemmings, shrews,

snowshoe hares

Region: IN, GW, SA, NS, SS, DC Location: Across the NWT

Northwest Territories Small Mammal and Hare survey

The NWT Small Mammal Survey monitors changes in density of voles, mice, lemmings, and shrews across five ecozones in the territory. The hare transect survey monitors snowshoe hare density across all forested ecozones, and an abundance index for Arctic hare at the tundra site.

Cluff, Dean

Environment and Natural Resources - North Slave dean cluff@gov.nt.ca

Permit No: 500092 Species Studied: Moose

Region: NS Location: Taiga Shield and Plains in the North Slave

Region

Moose abundance in the North Slave Region

The objectives of this study were: 1) to estimate the number and density of moose in the North Slave Region; and 2) to estimate the bull:cow and calf:cow ratios for population monitoring.

Cluff, Dean

Environment and Natural Resources - North Slave dean_cluff@gov.nt.ca

Permit No: 6886 Species Studied: Wolves

Region: NS Location: Central tundra region of the NWT

Aerial wolf den surveys and monitoring on the central tundra

The objectives of this study were: 1) to establish an annual relative abundance index for tundra wolves; 2) to investigate wolf population response to changing caribou abundance; and 3) to quantify frequency of den site usage.

Cluff, Dean

Environment and Natural Resources - North Slave dean cluff@gov.nt.ca

Permit No: 500070 **Species Studied:** Black bears **Location:** Along Highway 3 and 4

Baseline study on Black Bear Movements in the North Slave Region

The objectives of this study were: 1) to collect baseline information of black bear ecology, specifically on their movements and behaviour; 2) to document the movements and range sizes of black bears in the northern boreal forest; 3) to identify den site locations and black bear fidelity to them; 4) to document movements and behaviour of black bears around highways and food attractants; and 5) to provide management recommendations for relocating bears in the North Slave region.

Condon, Wayne

Aurora Research Institute

Permit No: 5414 **Species Studied:** Breeding birds

Region: GW **Location:** Around Fort McPherson, Inuvik, and Tsiigehtchic

Breeding bird surveys in the Gwich'in Settlement Region

The objectives of this study were: 1) collect information about species at risk in the Gwich'in Settlement Area and provide the data to the Species at Risk Stewardship Program and the local Renewable Resource Councils; 2) to collect information about breeding birds in the region on breeding bird survey routes 043-001, 002, 003 and 004 that were vacant (i.e. no surveyors were scheduled to complete the routes in 2012); 3) to provide the data collected to the Canadian Wildlife Service for inclusion in the North American-wide breeding bird surveys program to determine long-term population trends; and 4) to provide training for local field assistants so they may survey the routes in future years.

Coulton, Daniel

Golder Associates Limited daniel_coulton@golder.com

Permit No: 500013 **Species Studied:** Unqulates, raptors, water birds

Region: NS

Location: 15 km radius from NICO Project basecamp with

a 5 km buffer on either side of the proposed all-weather

access road

Baseline wildlife studies for the Fortune Minerals NICO Project

The objectives of this study were: 1) to further describe the occurrence, relative abundance, distribution and habitat use of wildlife in the study area; 2) to predict effects to the environment and wildlife from project development; 3) to provide baseline data for testing environmental effects

predictions and the effectiveness of mitigation; and 4) to guide further mitigation and adaptive management for reducing unexpected effects.

Coulton, Daniel

Golder Associates Limited daniel_coulton@golder.com

Permit No: 500011 **Species Studied:** Caribou, grizzly and black bears,

wolverines, wolves, raptors, water birds

Region: NS, SS **Location:** 70 x 80 kilometre rectangle centered on Kennady

Lake

Baseline wildlife studies for the De Beers Gahcho Kué Project

The objectives of this study were: 1) to further describe the occurrence, relative abundance, distribution, and habitat use of wildlife in the study area; 2) to predict effects to the environment and wildlife from project development; 3) to provide baseline data for testing environmental effects predictions and the effectiveness of mitigation; and 4) to guide further mitigation and adaptive management for reducing unexpected effects.

Cox, Karl

Environment and Natural Resources South Slave karl_cox@gov.nt.ca

Permit No: 4816 Species Studied: Wood bison

Region: SS Location: The known bison range in the Slave River

Lowlands between Fort Smith and Fort Resolution

Slave River Lowlands Bison population studies

The objectives of this study were: 1) to measure calf, yearling, and bull to cow ratios during the post-calving period for bison in the Slave River Lowlands; and 2) to monitor the Slave River Lowlands herd for the occurrence of anthrax related mortalities. If any suspected anthrax mortalities are detected, the Anthrax Emergency Response Plan would be activated and carcasses would be dealt with according to this plan. An increase in aerial surveillance would also occur if anthrax is suspected.

Cox, Karl

Environment and Natural Resources South Slave karl_cox@gov.nt.ca

Permit No: 4815 Species Studied: Wood bison

Region: DC, SS **Location:** The range of the Mackenzie and Nahanni-Liard

herds

Bison control area

The goal of the bison control program in the Northwest Territories is to reduce the risk of infection of the Mackenzie and Nahanni-Liard herds with tuberculosis and brucellosis. Objectives of the program were: 1) continue surveillance of the Bison Control Area; 2) maintain the Bison Control Area free of bison and prevent the establishment of any herds within its boundaries; and 3) increase public awareness of the bison control program.

Davison, Tracy

Environment and Natural Resources - Sahtú Region

tracy_davison@gov.nt.ca

Permit No: 500009 Species Studied: Caribou

Region: IN, SA **Location:** Late winter and calving/post-calving ranges of

the Bluenose-West barren-ground caribou herd

Radio collar deployment and a post-calving survey to estimate the number of caribou in the Bluenose-West herd in 2012

The objectives of this study were: 1) to deploy 40 satellite tracked and 20 GPS tracked radio collars on 60 adult caribou from the Bluenose-West herd on the late winter range. It is anticipated that 25-35% of collars will be deployed on bulls and the remainder on cows; 2) to obtain an accurate and precise estimate of herd size using aerial photographic survey on the post-calving range; and 6) to monitor movements and range use by remote tracking of satellite and GPS collars throughout the year.

Davison, Tracy

Environment and Natural Resources - Inuvik Region tracy_davison@gov.nt.ca

Permit No: 5415 Species Studied: Muskox and caribou

Region: IN Location: Melville Island; Prince Patrick Island; Byam

Martin Island; Eglinton Island; Emerald Island

Arctic Island caribou and muskox population survey

The objective of this study was to update the population estimate for Peary caribou and muskoxen. The last survey to estimate the population size for this region was conducted in 1997.

Davison, Tracy

Environment and Natural Resources - Inuvik Region tracy davison@gov.nt.ca

Permit No: 7424 Species Studied: Caribou

Region: IN **Location:** Late winter range of the Tuktovaktuk Peninsula.

Cape Bathurst, and Bluenose-West barren-ground caribou

herds

Late winter recruitment of the Tuktoyaktuk Peninsula, Cape Bathurst, and Bluenose-West barren-ground caribou herds

The objective of this study was to obtain a current estimate of late winter recruitment for the Tuktoyaktuk Peninsula, Cape Bathurst, and Bluenose-West barren-ground caribou herds.

Davison, Tracy

Environment and Natural Resources - Inuvik Region tracy_davison@gov.nt.ca

Permit No: 7423 **Species Studied:** Barren-ground caribou

Region: IN, GW Location: Inuvik region

Caribou sampling initiative: Caribou body condition and health monitoring

The objectives of this study were: 1) to monitor the estimated body weight, body fat and body protein of adult cow caribou over the winter and monitor trends over time; 2) to monitor selected fat depots of adult bull caribou over the winter to document trends over time; 3) to compare body condition to other herds being monitored using the same standardized system across the north; 4) to monitor levels of heavy metal contaminants in submitted caribou kidney samples; and 5) in the future the relationship of these trends to other indicators, such as pregnancy rate, calf survival, herd size, timing of spring thaw, fall storm patterns, and winter range snow depth will be investigated.

Elkin, Brett

Environment and Natural Resources - Wildlife brett elkin@gov.nt.ca

Permit No: 5768 Species Studied: Wildlife Region: NS, SS, IN, GW, DC, SA Location: NWT wide

Wildlife health, condition and genetic monitoring

The objectives of this study were: 1) to determine the cause of sick or dead wildlife found, harvested or handled by hunters, trappers, biologists, wildlife researchers, Renewable Resource Officers, or the general public; 2) to assist hunters and trappers by testing samples from harvested wildlife to determine what diseases or parasites are present, and the implications for consumption of the carcass; 3) to work cooperatively with hunters, trappers, biologists, Renewable Resource Officers and members of the general public to monitor the health and condition of wildlife on an on-going basis; 4) to identify the types, relative levels and geographical distribution of diseases, parasites and contaminants found in wildlife across the Northwest Territories; 5) to increase community awareness of diseases and parasites; and 6) to collect genetic information that will contribute to the understanding and management of wildlife populations.

Hood, Alex

De Beers Canada Inc. alexandra.hood@debeerscanada.com

Permit No: 500012 Species Studied: Caribou, grizzly bears, wolverines,

falcons, waterfowl, upland birds

Region: NS Location: In and around Snap Lake Mine

De Beers Snap Lake Mine: Wildlife effects monitoring program (WEMP)

Objectives of the monitoring studies were to determine if the mine influences: a) the density, distribution, group composition and behaviour of caribou; b) the relative activity (presence) and distribution of grizzly and black bears; and c) the relative activity (presence) and distribution of wolverine. Though not part of the wildlife effects monitoring program, wolf den activity was monitored to provide regional information to the Department of Environment and Natural Resources.

Kelly, Allicia

Environment and Natural Resources - South Slave allicia kelly@gov.nt.ca

Permit No: 4817 Species Studied: Muskox

Region: SS Location: Around Rennie Lake; Northeast of Fort

Resolution

Muskoxen abundance and distribution survey

The objective of this study was to estimate the abundance and distribution of muskoxen in each of the survey blocks. This information will be used to inform management decisions related to quotas and tag allocations for muskox in the South Slave region.

Kelly, Allicia

Environment and Natural Resources - South Slave allicia_kelly@gov.nt.ca

Permit No: 4818 Species Studied: Mink

Region: SS Location: Around Fort Smith and Kakisa in the Slave River

and Cameron River/Tathlina Lake riverine ecosystems

Contaminants in mink

The objectives of this study were: 1) to examine levels of metals and contaminants in mink along the Slave River and at a control site near Kakisa, and compare to levels documented in the same area almost 20 years previously; 2) to examine levels of contaminants in several key prey species, including muskrat, snowshoe hare and northern red-backed vole, and ensure safe levels of subsistence species (muskrat and hares); and 3) to look at patterns of contaminant input and accumulation in mink through a combination of existing data from water, sediment, plankton, amphipods and other prey species including fish along the Slave River.

Kelly, Allicia

Environment and Natural Resources - South Slave allicia_kelly@gov.nt.ca

Permit No: 4819 **Species Studied:** Boreal woodland caribou **Region:** SS **Location:** Hay River Lowlands, Cameron Hills

Boreal caribou monitoring - Hay River Lowlands (Ka'a'gee Tu Candidate Protected Area) and Cameron Hills

The objectives of this study were: 1) to monitor population demographics: adult female survival, calf production, ten-month calf recruitment, and finite rate of population increase (the relative change in size of population from one year to the next); 2) to document seasonal range use, annual home ranges and fidelity to calving areas (whether cows use the same area to calve year after year); and 3) to examine boreal caribou habitat use and selection in relation to natural and human caused disturbance (e.g. wildfire, development) and landscape features (e.g. forest type).

Kelly, Allicia

Environment and Natural Resources - South Slave allicia_kelly@gov.nt.ca

Permit No: 500045 **Species Studied:** Boreal caribou

Region: SS **Location:** Hay River Lowlands; Cameron Hills

Boreal caribou population trends and habitat use in the Hay River Lowlands and Cameron Hills

The objectives of this study were: 1) to monitor population demographics: adult female survival, calf production, ten-month calf recruitment, and finite rate of population increase (the relative change in size of population from one year to the next); 2) to document seasonal range use, annual home ranges and fidelity to calving areas (whether cows use the same area to calve year

after year); and 3) to examine boreal caribou habitat use and selection in relation to natural and human caused disturbance (e.g. wildfire, development) and landscape features (e.g. forest type).

Kelly, Allicia

Environment and Natural Resources - South Slave allicia kelly@gov.nt.ca

Permit No: 500043 **Species Studied:** Wood bison **Location:** Slave River Lowlands

Slave River Lowland's Bison Population Studies

The objectives of this study were: 1) to measure calf, yearling, and bull to cow ratios during the post-calving period for bison in the Slave River Lowlands; and 2) to monitor the Slave River Lowlands herd for the occurrence of anthrax related mortalities. If any suspected anthrax mortalities are detected, the Anthrax Emergency Response Plan is activated and carcasses are dealt with according to this plan. An increase in aerial surveillance also occurs if anthrax is suspected.

Kent, Christine

Rescan Environmental Services ckent@rescan.com

Permit No: 500082 Species Studied: Waterbirds and species identified within

the taiga shield ecozone

Region: NS **Location:** Tundra mine site

Scientific study at the Tundra Mine Site

The objectives of this study were: 1) to conduct a survey for waterfowl and shorebird broods at lakes and wetlands downstream of the water discharge route (Pathway 1); and 2) to record incidental sightings of wildlife (non-target species) observed at or between survey sites.

Klimstra, Jon

US Fish & Wildlife jon_klimstra@fws.gov

Permit No: 500060 Species Studied: Waterfowl **Region:** NS Location: Stagg River Delta

Western Canada cooperative waterfowl banding program

The objective of this study was preseason banding of 2,000 mallards, 1,500 northern pintails and 1,000 of all other waterfowl species at each of the approximately 20 banding stations in western Canada.

Larter. Nic

Environment and Natural Resources - Dehcho nic_larter@gov.nt.ca

Permit No: 5035 Species Studied: Wood bison

Region: DC

Location: The range of the Nahanni Wood Bison
(Northeastern British Columbia, southeastern Yukon
Territory and the southwestern NWT with the majority of the

range along and adjacent to the Liard and lower reaches of the South Nahanni River Valleys)

Monitoring of the Nahanni Wood Bison population

The objectives of this study were: 1) to measure calf, yearling, and bull:cow ratios during the post-calving period; 2) to monitor annual calf production and estimate overwinter survival of calf bison; 3) to collect biological samples as and when available from harvested animals or those involved in motor vehicle collisions; 4) to document seasonal movement patterns and range use of male and female bison throughout the range; 5) to delineate the area used by the population and document animal movement into new areas of range; 6) to document the frequency of river crossings by collared animals; 7) to identify and monitor the presence, movements, and behaviour of wood bison in communities; 8) to document the year round diet of Nahanni wood bison; and 9) to monitor the Nahanni wood bison population for the presence of brucellosis, tuberculosis and Johne's disease.

Larter, Nic

Environment and Natural Resources - Dehcho nic_Larter@gov.nt.ca

Permit No: 50377 **Species Studied:** Boreal caribou

Region: DC **Location:** East of the Mackenzie Mountains

Dehcho boreal caribou collar deployment

The objectives of this study were: 1) to monitor annual calf production, calf survival, and adult survival in order to make annual estimates of the rate of population change; 2) to ensure that the distribution of collared boreal caribou covers key areas throughout the range of boreal caribou in the Dehcho region; 3) to determine the calving period and the degree of fidelity of female caribou to calving areas over multiple years in areas with a range of seismic and fire disturbance history; 4) to use location data of female boreal caribou over multiple years overlaid with the current human footprint and wildfires to determine areas of high use and areas of avoidance by female boreal caribou in the landscape, and whether there is a seasonal component; 5) to provide empirical data to determine areas of secure boreal caribou habitat, given the current human footprint, and to compare this to the predictions and robustness of the study completed to predict high value boreal caribou habitats in the Dehcho; 6) to provide current knowledge of boreal caribou ecology for use with evaluating land use applications made in the Dehcho; 7) to provide empirical data for RSF modeling to assist with assessing important habitat types/areas; 8) as development occurs, to be able to assess responses of female caribou in relation to their use of space in the landscape; and 9) to continue to document and assess disease and parasites in boreal caribou.

Mougeot, Charlotte

MMG Minerals and Metals Group charlotte.mougeot@mmg.com

Permit No: 6891 **Species Studied:** Ungulates, wolverines, grizzly bears,

raptors, tundra breeding birds

Region: NS Location: Izok Lake area, including 30 km buffer,

preliminary rd alignment from Izok to Lupin

Baseline wildlife studies MMG's Izok Project 2012

The objectives of this study were: 1) to gather environmental baseline data to support environmental assessment of the Izok project; and 2) to provide the information needed to

develop a project proposal and environment impact statement for review by the Nunavut Impact Review Board and the Mackenzie Valley Environmental Impact Review Board.

Obst, Joachim jobst@ssimicro.com

Region: NS Location: Daring Lake Tundra Ecosystem Research

Station

Breeding densities and population trends for tundra birds at Daring Lake, NWT

The objectives of this study were: 1) to record breeding densities, habitats and population trends of tundra birds; 2) to use the data for monitoring the state of the environment and for conservation efforts; 3) to provide the data as an educational and research tool online for public use; and 4) to collect moulted and beached feathers of loons for contaminants analysis.

O'Keefe, Harry

Dominion Diamond Ekati Corporation harry.o'keefe@ekati.ddcorp.ca

Permit No: 6882 Species Studied: Caribou, grizzly bears, wolves,

wolverine, upland breeding birds, falcons

Region: NS Location: Ekati Diamond Mine property

Wildlife effects monitoring program (WEMP)

The objectives of this study was to test impact predictions and efficacy of mitigation measures for the following species: Caribou: to monitor the potential effects of the following mine activities: potential collisions with vehicles, incidents involving aircraft, general disturbance from the mine, roads as potential barriers, incidents at pits and the Long Lake containment facility. Grizzly bears: to monitor the potential effects of the following mine activities: potential collisions with vehicles, disturbance possibly affecting bear activity level, and the mine possibly attracting bears. Wolves: to monitor the potential effects of the following mine activities: potential collisions with vehicles, disturbance possibly affecting den use and the mine possibly attracting wolves. Wolverine: to monitor the potential effects of the following mine activities: potential collisions with vehicles, disturbance possibly affecting their presence near the mine and the mine possibly attracting wolverine. Upland breeding birds: to obtain a species count by conducting the North American breeding bird survey and maintaining incidental records of upland breeding birds, shorebirds and waterfowl. Falcons: to monitor the potential effects of the following mine activities: disturbance possibly affecting occupancy and productivity.

Pongracz, Jodie

Environment and Natural Resources - Inuvik Region jodie_pongracz@gov.nt.ca

Permit No: 5410 **Species Studied:** Polar bears

Region: IN **Location:** The Northern Beaufort polar bear subpopulation area: north of Norway Island and Victoria Island to the

area. Horur di Norway Island and Victoria Islan

mainland (southeast corner of the area)

Examining the boundary between the Northern Beaufort and Viscount Melville polar bear subpopulations

The objectives of this study were: 1) to assess the current boundaries of the Viscount Melville and Northern Beaufort polar bear subpopulations; and 2) to assess polar bear habitat use and movement patterns during changing sea ice conditions in the area of the Northern Beaufort Sea subpopulation north of Norway Island.

Pongracz, Jodie

Environment and Natural Resources - Inuvik Region jodie_pongracz@gov.nt.ca

Permit No: 5411 Species Studied: Polar bear Region: IN Location: Viscount Melville Sound

Viscount Melville Sound polar bear subpopulation survey

The objectives of this study were: 1) to conduct mark-recapture to estimate the current population size and demographic parameters of the Viscount Melville polar bear subpopulation; 2) to assess the current boundaries of the Viscount Melville polar bear subpopulation; and 3) to assess polar bear habitat use of changing sea ice habitat in the area of Viscount Melville Sound.

Pongracz, Jodie

Environment and Natural Resources - Inuvik Region jodie_pongracz@gov.nt.ca

Permit No: 7421 **Species Studied:** Wolverines **Region:** IN, GW **Location:** Inuvik region

Wolverine carcass collection

The objectives of this study were: 1) to document the distribution and level of wolverine harvest in the Inuvik Region; 2) to assess the age, sex ratios, condition, seasonal diet, and reproductive parameters of the wolverines harvested within the Inuvik region and compare to similar work in other regions of the Northwest Territories; and 3) to collect genetic material for later analysis to compare wolverine populations across the Northwest Territories.

Pongracz, Jodie

Environment and Natural Resources - Inuvik Region jodie_pongracz@gov.nt.ca

Permit No: 7422 **Species Studied:** Wolves **Region:** IN, GW **Location:** Inuvik region

Wolf sample collection

The objectives of this study were: 1) to document age, sex, condition and location of wolf harvests within the range of the Cape Bathurst and Bluenose-West caribou herds; 2) to document pack size and incidental observations of wolves and wolf kills; 3) to document diet of mainland wolves to determine what portion of the diet is caribou; and 4) to archive samples to look at genetic diversity of Arctic wolves.

Ramona, Maraj

Yukon Government ramona.maraj@gov.yk.ca

Region: IN Location: Southern Beaufort Sea polar bear subpopulation

range from the Alaska/Yukon border to Tuktoyaktuk

Southern Beaufort Sea polar bear project

The objectives of this study were: 1) to evaluate and further develop a non-invasive technique as a method that will allow estimation of numbers of bears; and 2) to provide additional information about the numbers of bears in the distant offshore and in particular over the Chevron offshore lease to complement the Beaufort Regional Environmental Assessment proposal.

Rausch, Jennie

Canadian Wildlife Service jennie.rausch@ec.gc.ca

Permit No: 5413 **Species Studied:** Arctic breeding shorebirds

Region: IN Location: Banks Island; Western Victoria Island;

Mackenzie Delta; Kendall island; Fish Island

Arctic shorebird monitoring program

The objectives of this study were: 1) to generate population estimates for all Arctic breeding shorebirds; 2) to produce maps showing shorebird distribution and abundance across the North American Arctic; 3) to identify highest-quality habitats for each shorebird species; 4) to provide shorebird densities and breeding ecology information at each survey site; and 5) to assist local managers in meeting their conservation goals.

Richardson, Evan

Environment Canada evan.richardson@ec.gc.ca

Region: IN **Location:** Aerial line-transect surveys for polar bears were

conducted in the Northern Beaufort Sea in areas up to 300

nautical miles offshore

Aerial survey for polar bears in the far offshore Beaufort Sea

The objective of this study was to assess the distribution and abundance of polar bears in areas of the offshore Beaufort Sea.

Robertson, Myra

Canadian Wildlife Service myra.robertson@ec.gc.ca

Permit No: 500069 Species Studied: Gulls

Region: NS **Location:** Frame Lake in Yellowknife

Gull surveys on Frame Lake, Yellowknife

The objective of this study was gather basic information on the species, numbers, and timing of gulls breeding on Frame Lake during 2012 with specific focus on two islands in the southern part of Frame Lake where gulls had nested in previous years

Robertson, Myra

Canadian Wildlife Service myra.robertson@ec.gc.ca

Permit No: 5412 Species Studied: Geese, swans

Region: IN Location: Mackenzie Delta; Tuktoyaktuk Pennisula;

Anderson River Delta; Kendall Island

Population management of geese and swans in the Inuvialuit Settlement Region using aerial surveys and banding studies

The objectives of this study were: 1) to monitor waterfowl populations in the ISR; 2) to evaluate the effect of harvest and other stressors on western Arctic populations of waterfowl; and 3) to monitor migration routes, harvest rates, and survival rates of the mainland goose populations.

Robertson, Myra

Canadian Wildlife Service myra.robertson@ec.gc.ca

Region: NS Location: Along Highway 3 near Yellowknife

Abundance and productivity of waterfowl and other aquatic birds breeding in the boreal forest

The objectives of this study were: 1) to monitor population trends and productivity of boreal waterfowl and aquatic bird populations near Yellowknife; and 2) to determine factors that limit the size, composition, and productivity of the breeding populations of aquatic birds near Yellowknife.

Robertson, Myra

Canadian Wildlife Service myra.robertson@ec.gc.ca

Permit No: 500037 **Species Studied:** Herring gulls and gull species

Region: NS Location: North Arm of Great Slave Lake

Chemicals management plan wildlife national monitoring program

The objective of this study was to assess the toxicological characteristics of local gulls in relation to national data.

Selwyn Chihong Mining Ltd.

Permit No: 5766 **Species Studied:** Caribou, moose **Region:** SA, DC **Location:** Howard Pass Access Road

Selwyn Project baseline studies

The objectives of this study were: 1) to support permit applications and the environmental review process; 2) to predict and mitigate effects to the environment and wildlife that may result from project development; 3) to provide pre-development information in support of any future environmental affects monitoring program; and 4) to contribute to regional studies for assessing and managing potential cumulative effects.

Sharam, Greq

Rescan Environmental Services

gsharam@rescan.com

Permit No: 6885 **Species Studied:** Ungulates, carnivores, waterbirds, loons

and grebes, shorebirds, upland breeding birds, raptors

Region: NS Location: Courageous Lake project

Wildlife baseline program at the Courageous Lake Study Area

The objective of this study was to characterize baseline conditions and ecology of the area early in the process of any potential development.

Sharam, Greg

Rescan Environmental Services

Permit No: 4832 **Species Studied:** Ungulates, carnivores, wetland birds,

terrestrial breeding water birds

Region: DC **Location:** Tamerlane's Pine Point Property

Pine Point Project wildlife baseline program

The objective of this study was to collect data on wildlife habitat features and evaluate selected wildlife populations and landscape use within the five deposits LSA and RSA and N-204 LSA.

Tout, Ann Marie

Enbridge Pipelines (NW) Inc. annmarie.tout@enbridge.com

Permit No: 5767 Species Studied: Wildlife

Region: SA **Location:** Enbridge Right-of-Way

Monitoring wildlife along the Enbridge Right-of-Way

The objective of this study was support and encourage community-based programs to document wildlife sightings and wildlife tracks along the Enbridge Right-of-Way.

Wells, David & Claudine Lee

Dominion Diamond Ekati Corporation & Diavik Diamond Mine Inc. claudine.a.lee@bhpbilliton.com

Permit No: 500068 Species Studied: Grizzly bears

Region: NS Location: Lac de Gras Region including Ekati and Diavik

mine sites

2012 Joint regional grizzly bear DNA hair snagging program

The objective of this study was to determine if mine-related activities influence the relative abundance and distribution of grizzly bears over time

Wortham, Jim

United States Fish and Wildlife Service jim wortham@fws.gov

Permit No: 500058 Species Studied: Waterfowl

Region: NS, SA, GW, IN **Location:** Mackenzie River drainage

Cooperative waterfowl population surveys in the Northwest Territories

The objective of this study was to determine the size and species composition of the breeding populations of ducks and other waterfowl in Mackenzie River drainage.

FISHERIES PERMITS 2012

Bisaillon, Jean-Francois

Parks Canada - Western Arctic Field Unit jean-francois.bisaillon@pc.gc.ca

Licence Number: S-12/13-3017-YK

Species: Arctic char, lake trout Location: Cache Lake

Uyarsivik Monitoring Program for Tuktut Nogait National Park

The objectives of this study were: 1) to document and assess the density and abundance of Arctic char and Lake trout fish habitat; and 2) to evaluate the biological integrity of the fish population.

Blanchfield, Paul

Fisheries and Oceans Canada paul.blanchfield@dfo-mpo.gc.ca

Licence Number: S-12/13-3012-YK

Species: Lake trout Location: Alexie Lake

Determination of lake trout habitat use in an NWT lake, with a focus on spawning and overwintering

The objective of this study was to quantify lake trout (Salvelinus namaycush) habitat use in a northern lake, with a special focus on critical habitat, such as spawning and over-wintering areas. This information will provide the foundation for more investigations on physical lake trout habitat in the future and is critical if we are to better understand, predict, and mitigate the potential impacts of industry to fish and fish habitat. The research team will also provide a rich data set to hypothesize and model on the effects of climate change to cold-water fish species such as lake trout in northern lakes, which are predicted to be most severely affected by climate change.

Clark, Robert (Bob)

Environment Canada bob.clark@ec.gc.ca

Licence Number: S-12/13-3011-YK

Species: Amphipod (gammarus lacustris) **Location:** Rae Lake

Effect of water temperature on breeding phenology and seasonal abundance of amphipod Gammarus lacust

The objectives of this study were: 1) to determine how water temperature affects breeding phenology; and 2) to study seasonal abundance of the amphipod Gammarus lacustris, in a laboratory experiment.

Cott, Peter

Fisheries and Oceans Canada pete.cott@dfo-mpo.gc.ca

Licence Number: S-12/13-1053-NU-A1

Species: All fish species Location: Yellowknife Bay

YK Bay PCB and stable isotope fish tissue analysis

The objectives of this study were: 1) to investigate the PCB levels in Great Slave Lake burbot, define PCB levels in fish tissues, make comparisons between sexes, and southern Great Lakes; and 2) to define the trophic position and food web dynamics of northern fish communities in boreal lake ecosystems to enable comparisons among lakes, among apex predators, and between climatic regions.

Dunmall, Karen

Fisheries and Oceans Canada karen.dunmall@dfo-mpo.gc.ca

Licence Number: S-12/13-3025-YK

Species: Pacific salmon Location: Mackenzie River, Mackenzie

Delta Area, Beaufort Sea

Pacific salmon distribution in the Western Arctic

The objective of this study was to collect information on the distribution of Pacific salmon in the western Arctic. Once the study establishes a basic understanding of the distribution for each species we can monitor annual catches to track dispersal. A major shift in these distributions could serve as an indicator of environmental change.

Evans. Marlene

Environment Canada marlene.evans@ec.gc.ca

Licence Number: S-12/13-3033-YK

Species: Lake trout, burbot, northern pike **Location:** Great Slave Lake Area, Stark

Lake

Spatial and long-term trends in persistent organic contaminants and metals in fish from the NWT

The objective of this study was to investigate whether contaminant levels are changing in fish in the Northwest Territories with a focus on Great Slave Lake which we have been studying since the early 1990s. The study plan was to collect lake trout from Great Slave Lake (Hay River area and ŁutselK'e area. The study plan was also to collect burbot from the ŁutselK'e and Fort Resolution areas of Great Slave Lake, and northern pike from the Fort Resolution area of Great Slave Lake.

Gantner, Nikolaus

University of Victoria gantnern@uvic.ca

Licence Number: S-12/13-3001-YK

Species: All fish species **Location:** Noell Lake, Big Lake (Ilkaasuat),

Ya Ya Lake

Evaluation of hydro-climatic drivers of contaminant transfer in aquatic food webs in Husky Lakes area

The objectives of this study were: 1) to characterize the lake-ice conditions and associated lake productivity and aquatic food webs along a hydroclimatic/salinity gradient within the Husky Lakes Watershed (HLW); 2) to develop and validate a process-based hydro-ecological food-web subcomponent coupling changes in hydro-climatology and ice conditions to water column productivity and food webs; 3) to explain uptake of Hg including isotopes as tracer/marker through a comparison of: i) spatial comparison of Hg bioaccumulation in food webs; ii) Hg stable isotope ratios in biota along a salinity gradient in the HLW and with lakes outside the HLW. 4) to review of existing and new documentation of Traditional Knowledge (TK) on historical and present ice and climate conditions and fish biology and subsistence fisheries; and 5) to combine both knowledge bases to help develop future strategic monitoring of locally relevant sites.

Harris. Katherine

Golder Associates Ltd. katherine_harris@golder.com

Licence Number: S-12/13-3023-YK

Species: All fish species **Location:** Burke Lake and Area

Fortune Minerals - 2012 open-water plankton baseline program for the NICO Project

The objective of this study was to assess baseline seasonality in the phytoplankton and zooplankton communities in three lakes expected to be directly influenced by the NICO Project, as well as from a reference lake, and to collect sufficient baseline data to allow for comparison with potential future monitoring data. Additional plankton sampling will occur in a wetland area that will be permanently covered by Project-related activities to assess and document the seasonal variation in a wetland plankton community with elevated arsenic levels during baseline conditions.

Harris, Katherine

Golder Associates Ltd. katherine_harris@golder.com

Licence Number: S-12/13-3030-YK

Species: Slimy sculpin, ninespine stickleback **Location:** Baker Creek

Giant Mine Phase 4 Environmental Effects Monitoring

Giant Mine is in their fourth phase of the Environmental Effects Monitoring (EEM) program under the Metal Mining Effluent Regulations (MMER). A lethal Slimy Sculpin survey and a non-lethal Ninespine Stickleback survey are required as part of the EEM to determine if treated mine effluent is affecting fish. The specific objective of the study was to examine the health of two small-bodied fish species (Slimy Sculpin and Ninespine Stickleback) exposed to treated effluent from Giant Mine. Additional sampling for Arctic Grayling may also be required as part of an on-going sediment study in Baker Creek.

Harwood, Lois

Fisheries and Oceans Canada lois.harwood@dfo-mpo.gc.ca

Licence Number: S-12/13-3015-YK **Species:** Bearded seal, ringed seal

Location: Ulukhaktok Area - Coastal Marine

Waters

Ulukhaktok barvest Based monitoring of ringed and bearded seals

The objectives of this study were: 1) to assess of reproduction, condition disease and contaminants of ringed seals and bearded seals through harvest-based monitoring at Ulukhaktok, NT, 2012; 2) to sample and measure ringed seals taken in the annual harvest in the Ulukhaktok (n=100) area, using reproductive status and body condition as indicators of ecosystem productivity and fluctuations in the seal population; 3) to examine the aspects in objective 1 in the context of regional ice conditions; 4) to co-ordinate with, and provide samples for, diet and stock health studies, such as disease, particularly relevant given the UME in Alaska, NWT, Nunavut and Russia; and 5) to sample and measure any bearded seals that happen to be taken in the annual harvest in the Ulukhaktok (n=5) areas, to examine reproductive rates, growth, condition and prey preferences.

Hoos, Richard

EBA Engineering Consultants rhoos@eba.ca

Licence Number: S-12/13-3009-YK

Species: Slimy sculpin, aquatic invertebrates **Location:** Flat River

Aquatic Environmental Effects Monitoring - Flat River at Cantung Mine

The objective of this study was to determine length, weight, and general condition of 100 fish and tissue analysis of not more than 10 fish (Slimy Sculpin – Cottus cognatus) per site at five site locations in the Flat River: two locations upstream of the CanTung mine site, one adjacent to the mine site and at two locations downstream of the mine site. Also, to determine sediment and water quality (nutrients and metals) as well as benthic invertebrate (10 sampler per site) abundance and taconomic composition at the same sites. Statistical methods will be used to determine whether differences between sample sites (exposure and control) are of statistical significance.

Howland, Kimberly

Fisheries and Oceans Canada kimberly.howland@dfo-mpo.gc.ca

Licence Number: S-12/13-3007-YK-A1

Species: Lake trout, cisco **Location:** Great Bear Lake (Keith Arm)

Monitoring of lake trout stocks and cisco diversity in Great Bear Lake

The objective of this study was to monitor size and age structure, fecundity (egg number per female), growth and mortality of lake trout populations from Dareli (Keith), Turili (McVicar), Kwit tla (McTavish), Tugacho (Dease) and Tirato (Smith) Arms of Sahtú (Great Bear Lake). These data will be used for stock assessment purposes and to follow changes in the biological characteristics of lake trout stocks over time: 1) to determine the extent of movements (if any) by lake trout in Great Bear Lake through molecular genetics; 2) to monitor species composition and if sufficient data are available, presence, size structure and other biological characteristics of by catch and invertebrate species; 3) to examine the morphological, meristic and life history characteristics of archived ciscos collected from Great Bear Lake over the past 7 years to test the hypothesis that there are multiple forms/species including shortjaw; 4) to conduct targeted sampling and examination of characteristics for cisco from deeper regions of Great Bear Lake (>50 m) to

increase sample size and increase the range of surveyed habitat; and 5) to compare Great Bear Lake cisco with shortjaw cisco identified in other lakes to verify species identification and provide information that will feed into broader questions regarding the taxonomy of shortjaw cisco.

Howland, Kimberly

Fisheries and Oceans Canada kimberly.howland@dfo-mpo.gc.ca

Licence Number: S-12/13-3016-YK

Species: Arctic char Location: Hornaday River

Arctic char monitoring at Hornaday and Brock rivers, NT, 2012

The objectives of this study were: 1) to maintain char monitoring project and continue to provide information on status and life history of Arctic char captured at the mouth of the Hornaday and Brock rivers; 2) to confirm the identification of 'blue char' captured near Tippi (weatern Darnley Bay); and 3) to provide important support information for the formulation, delivery and compliance of the Paulatuk Char Management Plan.

Howland, Kimberly

Fisheries and Oceans Canada kimberly.howland@dfo-mpo.gc.ca

Licence Number: S-12/13-3020-YK

Species: Dolly varden **Location:** Babbage River System

Population studies on Dolly Varden from the Northwest Territories and Yukon North Slope

The objectives of this study were: 1) to investigate the presence of anadromous Dolly Varden in the Big Fish River that may not migrate to sea annually after smoltification; 2) to conduct mark recapture studies: recapture tagged Dolly Varden in the Big Fish and Babbage rivers, and then tag Dolly Varden from the Big Fish River, Babbage River, Firth River and Joe Creek; 3) to investigate the presence of Dolly Varden in Fish Creek and possibly seine and tag 100 Dolly Varden from each river system for mark-recapture; and 4) to collect 20 resident Dolly Varden from the Big Fish and Babbage rivers during the fall in order to obtain biological information such as length, weight, age, sex and maturity and diet, and tissue samples for mercury analysis. Sample subsistence catch of Dolly Varden for biological information at Herschel Island and Ptarmigan Bay, Yukon.

Howland, Kimberly

Fisheries and Oceans Canada kimberly.howland@dfo-mpo.gc.ca

Licence Number: S-12/13-3024-YK

Species: Plankton Location: Rat River Area

Biological investigation of Dolly Varden and arctic grayling from the Rat River

The objectives of this study were: 1) to conduct a harvest-based monitoring program to obtain tag returns, and collect catch-effort and biological information; 2) to obtain biological data of Dolly Varden captured at the spawning/overwintering area of the Rat River (Fish Creek); and 3) to collect dead-samples of Arctic grayling in order to obtain biological information such as length, weight, age, sex and maturity, diet information and tissue samples.

Howland, Kimberly

Fisheries and Oceans Canada kimberly.howland@dfo-mpo.gc.ca

Licence Number: S-12/13-3027-YK

Species: Arctic char **Location:** Fish Lake

Assessment of arctic char stock from Fish Lake

The objective of this study was to conduct a harvest-based assessment program involves enumerating and measuring Arctic char taken in the annual subsistence harvest at Fish Lake each October-November. Indicators of stock status such as CPUE, age, length, weight, sex and maturity are used to evaluate the impact of the fishery on the stock, and to provide information on status and life history of the char stock. The study has been done annually since 1992, and is one of the longest char monitoring studies in place in the ISR. It has provided important support for formulation, delivery and compliance of the Holman Char Fishing Plan.

Johnson, Jim

Fisheries and Oceans Canada jim.johnson@dfo-mpo.gc.ca

Licence Number: S-12/13-3021-YK

Species: All species (except marine mammals) **Location**: Darnley Bay

Darnley Bay nearshore fish survey

The objective of this study was to conduct nearshore fish survey along the west side of Darnley Bay; the targeted sampling sites are Brown's Harbour and Bennett Point, weather and ice permitting; an alternate site would be the Argo Bay area. The objective of the study is to determine what species of fish are found in the area, to collect basic biological information from a relatively small number of these fish, and to collect tissue samples from an even smaller number of dead-sampled fish for follow-on laboratory analyses. These data and biological samples will be utilized by a number of linked projects and initiatives including: 1) the BREA offshore fish survey; 2) the beluga monitoring program in developing a better understanding of their potential diet in the area; 3) development of an understanding of the trophic structure in this area in support of the establishment of an MPA in Darnley Bay; and 4) the development of a community-based ecosystem monitoring program based out of Paulatuk. It is not the purpose of this study to collect large numbers of fish to determine detailed population parameters for the fish community inhabiting these waters.

Lea. Ellen

Fisheries and Oceans Canada ellen.lea@dfo-mpo.gc.ca

Licence Number: S-12/13-3018-YK

Species: Arctic char Location: Ulukhaktok Area - Coastal Marine

Waters

Ulukhaktok summer coastal harvest monitoring 2012

The objective of this study was to collect harvest and biological information from the summer coastal subsistence Arctic char harvest by the community of Ulukhaktok. The information that will be collected by the monitor is an integral part of the community fishing management plans that

are established between the Olokhaktomiut Hunters and Trappers Committee, Fisheries Joint Management Committee, and the Department of Fisheries and Oceans Canada.

Lea, Ellen

Fisheries and Oceans Canada ellen.lea@dfo-mpo.gc.ca

Licence Number: S-12/13-3029-YK

Species: Dolly varden (searun) Location: Mackenzie Delta Area

Big Fish River Dolly Varden Harvest monitoring program

The objective of this study was to collect harvest and biological information from the subsistence harvest of 150 Big Fish River Dolly Varden Char. The monitor collected biological information and samples including length, weight, sex and maturity, otoliths, gonads for fecundity, a small piece of muscle tissue for mercury analysis, and liver and fin clips for genetic analysis for as many fish as possible while respecting harvester's wishes.

Leonard, Deanna

Fisheries and Oceans Canada deanna.leonard@dfo-mpo.gc.ca

Licence Number: S-12/13-3036-YK

Species: All fish species Location: Snare Lake

Snare Lake Fish Camp

The objectives of this were: 1) to document and combine traditional knowledge of fish in the Wek'weeti area with conventional fish sampling methods; 2) to obtain baseline data on mercury levels in fish consumed by Wek'weeti residents (Snare Lake area); 3) to evaluate mercury levels found in fish in terms of species and size/age relationships, habitat use, trophic status; and 4) to establish a monitoring program to identify contaminant levels and changes in levels through time for fish in the Tłicho region as part of a larger Aquatic Ecosystem monitoring program.

Loseto, Lisa

Fisheries and Oceans Canada lisa.loseto@dfo-mpo.gc.ca

Licence Number: S-12/13-2001-HR

Species: Beluga whales Location: Kendall Island, Hendrickson

Island

Assessment of beluga whales through harvest-based monitoring

The objective of this study was to maintain long term sampling of beluga in the Tarium Niryutait Marine Protected Area. Community beluga samplers will record morphological measurements and collect tissue samples (blubber, liver, kidney, muscle/meat, skin, blood, lower jaw) taken during the regular subsistence hunt. The hunters will permit access to their landed whales for sampling (aging structures, tissues for contaminants, disease and diet testing) and recording measurements (girth, length, fatness). Samples collected are frozen for preservation. Kendall Island is also a fishing area we may sub sample fish taken during the subsistence harvest as part of the community monitoring program (supported by FJMC). Samples will be sent to Fisheries and Oceans Canada (DFO) labs in Winnipeq.

Loseto, Lisa

Fisheries and Oceans Canada lisa.loseto@dfo-mpo.gc.ca

Licence Number: S-12/13-3019-YK

Species: Dolly varden Location: Shingle Point

Community based monitoring of coastal fish ecology using biomarkers

The objective of this study was to expand existing knowledge of ecosystem structure and function within the TNMPA by collecting fish samples for stable isotopes and fatty acids analyses from fish caught by local fishermen at the Shingle Point. Additionally, collect catch and biological information (length, weight, sex/maturity and ageing structure) and tissue samples (fin clips for genetic analysis and stomachs) of fish species from the subsistence harvest with an emphasis on obtaining tag returns from and determining the total catch of Dolly Varden. Results of analyses will provide information on inter-annual variability in baseline signatures and information on harvest levels and data useful to evaluate the Dolly Varden mixed stock fishery at Shingle Point. The program will provide training to support: 1) long-term, community-based, coastal monitoring; and 2) an evaluation of indicators for cumulative impact monitoring.

Low, George Dehcho AAROM geobarbgeo@hotmail.com

Licence Number: S-12/13-3034-YK

Species: All fish species

Location: Bluefish Lake, Gargan Lake, Deep Lake, Ekali Lake, Cli Lake, Little Doctor Lake, Yendo Lake, Greasy Lake, Long Lake, Mackenzie River, South Nahannie River,

Liard River

Updating data on mercury levels in food fish species in lakes used by Dehcho communities.

The objective of this study was to update data on mercury levels of certain fish in Dehcho lakes.

The objective of this study was to update data on mercury levels of certain fish in Dehcho lakes that have shown elevated levels in previous studies. This project will also determine if any other traditional fishing lakes are affected. The data from this study will also be used by scientists to help determine which factors are leading to increases of mercury in fish. It is also an objective of this study to provide traditional knowledge data on which lakes and fish species are being used for subsistence by the first nations.

Machtans, Hilary

Golder Associates Ltd. hilary_machtans@golder.com

Licence Number: S-12/13-2002-HR-A1

Species: Lake trout **Location:** Snap Lake

De Beers Snap Lake Mine AEMP

The objective of this study was to undertake a series of open-water programs to fulfill the requirements of the Water License and Fisheries Authorization for the Snap Lake Mine. The studies are designed to determine fish health, to conduct a reconnaissance program and monitor lake trout populations with the use of PIT tags.

Machtans, Hilary

Golder Associates Ltd. hilary_machtans@golder.com

Licence Number: S-12/13-2003-HR-A1

Species: All fish species Location: Yellowknife Bay

Con Mine Environmental Effects Monitoring Program fourth phase

The objective of this study was to complete the 4th phase of the Environmental Effects Monitoring Program at Con Mine to investigate the health of fish from an area exposed to mine effluent.

Mason, Kristine

Golder Associates Ltd. kristine_mason@golder.com

Licence Number: S-12/13-3013-YK

Species: All fish species Location: Kennady Lake watershed (A3, A1,

D3, D2, E1)

Gahcho Kué Project

De Beers Canada Inc. (De Beers) has been conducting fish and fish habitat baseline studies for a number of years in the area of the Gahcho Kué Project. The overall objective of this year's program is to collect fish in the lakes west of Kennady Lake identified as potential habitat compensation options in spring and summer.

McPherson, Morag

Fisheries and Oceans Canada morag.mcpherson@dfo-mpo.gc.ca

Licence Number: S-12/13-3005-YK

Species: Arctic grayling Location: Yellowknife River

Arctic grayling stream habitat use in the Yellowknife Area

The objective of the study was to document the baseline characteristics of select Arctic grayling spawning streams and rivers in the northern region of Great Slave Lake.

Mochnacz. Neil

Fisheries and Oceans Canada mochnacznj@dfo-mpo.gc.ca

Licence Number: S-12/13-3031-YK

Species: Bull trout Location: Prairie Creek

Bull Trout Ecology and Metals in the South Nahanni Watershed

The objective of this study was to improve our understanding of metal contamination and ecology of Bull Trout and associated species in the South Nahanni watershed. Specific objectives are: 1) document spawning and winter habitat use; 2) develop a non-lethal technique for monitoring mercury levels; 3) investigate spatial relationships and connectivity among critical habitats; 4) develop a standardized stream sampling protocol for monitoring juvenile habitat.

Moore, Kristine

Diavik Diamond Mines Inc. kristin.moore@riotinto.com

Licence Number: S-12/13-3028-YK

Species: Round whitefish, longnose sucker, **Location:** Lac de Gras Area (Lake E17)

lake trout, lake whitefish

Diavik fish palatability and tissue chemistry study

The objective of this study was to catch fish in Lac de Gras for tasting by local Aboriginal groups to determine texture and taste of fish in the receiving lake for the mine effluent. Obtain aging structures and tissue samples for mercury analysis of 20 lake trout in Lac de Gras to contribute to an increased knowledge base of age to mercury concentration ratios in local trout populations. Monitor mercury levels in Lac de Gras trout to determine if concentrations are increasing over time.

Morinville, Genevieve

Rescan Environmental Services gmorinville@rescan.com

Licence Number: S-12/13-3032-YK

Species: All fish species Location: Lakes at EKATI Diamond Mine

2012 EKATI AEMP

Fish monitoring was conducted as part of the Aquatic Effects Monitoring Program at EKATI every six (previously five) years. Monitoring of impacted and reference lakes was conducted through the use of gillnetting and electrofishing. Monitoring was conducted to ensure there are no impacts to the environment due to the operation of the EKATI Diamond Mine.

Muir, Andrew

Great lakes Fishery Commission amuir@glfc.org

Licence Number: S-12/13-3022-YK

Species: Lake trout Location: Great Bear Lake

Lake trout diversity in Great Bear Lake: Do deep-water forms exist?

The objective of this study was to identify and describe lake trout forms (e.g., deep versus shallow dwelling trout) in North America, and to use this information to understand potential impediments to, and strategies, for re-establishing extirpated lake trout forms in the Laurentian Great Lakes. Our research will also provide information on how deepwater food webs are structured and identify evolutionary patterns and processes. The objective of our Great Bear Lake project is to sample known and newly discovered deepwater (>50 m) habitats southwest of the Narakay Islands and within the McTavish Arm to determine if a deepwater humper or siscowet lake trout form occurs. Data on the occurrence and characteristics of deepwater lake trout forms from Great Bear Lake will fill in gaps in our North American lake trout dataset. Currently it is unknown if deepdwelling lake trout forms occur in Great Bear Lake. Our sampling efforts will be concentrated in deep (>50m), offshore waters that are not targeted by subsistence fisheries or by recreational anglers.

Olenick, Adrian

MWH Canada Inc.

adrian.olenick@mwhglobal.com

Licence Number: S-12/13-2004-HR

Species: All fish species

Location: Streams: (65°00'59" 126°50'45"), (65°03'45" 126°56'17"), (65°01'45" 127°07'54"), (65°08'36" 126°57'11"). Lakes: (64°59'18" 126°46'26"), (65°03'46" 126°57'33"), (65°01'07" 127°00'18"), (65°00'58" 126°50'07")

Fish population study in the Sahtú Settlement Area, NT

The objective of this study was to determine presence/absence of fish species and gather baseline data on species found.

Olenick, Adrian

MWH Canada Inc

adrian.olenick@mwhglobal.com

Licence Number: S-12/13-3035-YK

Species: All species (excludes marine mammals) **Location:**

Location: Streams: 1-N645432.719W 261933.277, 2-N65036.484 W 1262721.267, 3-N65234.639 W 1261650.219, 4-N645436.880 W 126655.535

N645136.889 W 126655.535

Fish population study of Four Streams in the Sahtú Region NT

The objective of the study was to identify the fish species present, and to gather baseline population data on said species, in waterbodies in the area of a proposed Husky drilling project. This data will allow mitigative measures that benefit the specific species encountered to be employed during and following the program. Four streams are to be assessed.

Panayi, Damian

Golder Associates Ltd damian panayi@golder.com

Licence Number: S-12/13-3014-YK-A1

Species: All fish species Location: Yellowknife River

NTPC Bluefish Hydro fish salvage and fisheries survey

The objectives of this study were: 1) to conduct a spring fish salvage below the dam construction site on the Yellowknife River; 2) to monitor fish habitat compensation as required by Fisheries Authorization 09-HCAA-CA-00079-A; and 3) to monitor changes to mercury in fish community as per water licnce MV2004 L4-0004.

Paradis, Adrian

Aboriginal Affairs and Northern Development Canada adrian.paradis@aandc-aandc.qc.ca

Licence Number: S-12/13-3037-YK

Species: All fish species Location: Yellowknife Area

Yellowknife Bay fish collection and tissue sampling - Giant Mine Remediation Project

The objective of this study was to create an updated preliminary dataset for fish tissue chemistry in Back Bay, Yellowknife Bay and one reference area for the Giant Mine Remediation Project in Yellowknife, NT. This program represents a first step in building a 'pre-remediation' dataset on fish tissue contaminant concentrations. The data gathered from the program will be used to support the development of a 'Life of Project' Environmental Monitoring Program, and provide information to support the detailed design of infrastructure associated with a proposed effluent diffuser in Yellowknife Bay. The primary objective of the aquatics field program is to collect baseline data on water chemistry, sediment chemistry, plankton and benthic invertebrates in Yellowknife Bay.

Pink, Melissa

Rescan Environmental Services Ltd.

Licence Number: S-12/13-3002-YK

Species: All fish species Location: Courageous Lake Project Area

Courageous Lake Project 2012

Rescan Environmental Services Ltd. (Rescan) was retained by Seabridge Gold Inc. to complete the 2012 Courageous Lake fish and fish habitat environmental program. The objectives for the summer of 2012 will be to: 1) conduct further detailed assessment of fish and fish habitat of Matthews Creek; 2) increase baseline sampling of fish and fish habitat in selected areas within the Project area; 3) increase sampling of fish and fish habitat at potential fish habitat compensation sites, including previously unstudied sites; and 4) determine the presence or absence of fish in waterbodies within the proposed airstrip expansion area.

Pink, Melissa

Rescan Environmental Services Ltd.

Licence Number: S-12/13-3003-YK

Species: All fish species Location: Unnamed Lake at Pine Point

Baseline fisheries surveys for the Pine Point Project

Rescan Environmental Services Ltd. (Rescan) was retained by Tamerlane Ventures Inc. to conduct the 2012 year-two summer baseline fish and fish habitat surveys for the Pine Point Project approximately 42 km east of Hay River, Northwest Territories. The information collected during year-two will build upon current understanding of fish and fish habitat within the study area and assist in preparing a Developers Assessment Report (DAR). The objectives for the 2012 year-two baseline study are to: 1) identify whether waterbodies near the deposits or the footprint of the project infrastructure are fish-bearing; 2) to collect biological data on fish found in those waterbodies; 3) prepare a baseline report that describes the fish habitat and fish population of the study area; and 4) collect baseline tissue data to inform the Country Foods Study.

Reist. Jim

Fisheries and Oceans Canada jim.reist@dfo-mpo.gc.ca

Licence Number: S-12/13-3026-YK

Species: All species (excludes marine mammals) **Location**: Beaufort Sea (Offshore)

Fishes, habitats and ecosystem linkages to oil and das Development in the Canadian Beaufort Sea

The objective of this study was to address gaps regarding offshore marine fishes and supporting ecosystem components, thereby 1) increasing knowledge of the Beaufort Sea marine ecosystem, the fishes therein, and the structural and functional relationships to key biota harvested by Inuvialuit; 2) establishing pre-development baselines within which developmental stressors may be assessed; and 3) providing a benchmark within which anticipated future effects of climate change may be assessed in the context of developmental impacts.

Renaud, Claude

The Canadian Museum of Nature crenaud@mus-nature.ca

Licence Number: S-12/13-3004-YK

Species: Arctic lamprey Location: Hay River

Evolution of arctic lampreys

The objective of this study was to learn more about the lampreys of NWT. It is part of a larger scale study of lampreys of the genus Lethenteron across the northern hemisphere (Eurasia and North America). We have four learning objectives: 1) describe the gross morphology and pigmentation of the larvae of the two lamprey species that occur in the NWT; 2) compare their DNA (mitochondrial and nuclear markers); 3) compare their gonadal histology; and 4) compare their gene expression.

Savage, Pascale-Laure

RESCAN Environmental Services Ltd. psavage@rescan.com

Licence Number: S-12/13-3008-YK

Species: All fish species Location: Sandy Lake

Tundra Mine construction activity monitoring program

Rescan Environmental Services Ltd. (Rescan) was retained by Contaminants and Remediation Directorate (CARD), Aboriginal Affairs and Northern Development Canada (AANDC), NT Region to conduct fish larval surveys to detect the potential impacts from the discharge of treated effluent in the project area and its targeted pathway. The 2012 field program's objective will be to complete year-three of the Construction Activity Monitoring Program by conducting fish habitat and larval surveys along six potentially effluent impacted tributaries.

Sawdon, Lorraine

Fisheries and Oceans Canada lorraine.sawdon@dfo-mpo.gc.ca

Licence Number: S-12/13-3006-YK

Species: Northern pike, lake trout **Location:** Alexie Lake

Improving population estimates for lake trout and northern pike in Alexie Lak

The objective of the study was to conduct a recapture event (angling) for a modified Petersen's population estimate on large-bodied fish species (Lake Trout and Northern Pike), in Alexie Lake in the NWT, to improve the accuracy of the 2011 population estimates.

Tonn, William

University of Alberta bill.tonn@ualberta.ca

Licence Number: S-12/13-3000-YK

Species: All fish species Location: Lac de Gras Area Lakes

Improving habitat connectivity to enhance productive capacity of arctic freshwater ecosystems

The objective of this study was to evaluate the effectiveness of the habitat manipulations and assess the biotic and abiotic responses to these treatments. In 2012, the study initiated the first year of post-treatment monitoring of the habitat and biota in the study systems, following 3 years of pre-treatment monitoring.

Tonn. William

University of Alberta bill.tonn@ualberta.ca

Licence Number: S-12/13-3038-YK

Species: All fish species Location: Lac de Gras Area Lakes

Improving habitat connectivity to enhance productive capacity of arctic freshwater ecosystems

The objective of this study was to enhance accessibility of the lakes to fish and provide in-stream habitat enhancement to promote spawning and rearing. This study is to evaluate the effectiveness of the habitat manipulations and assess the biotic and abiotic responses to these treatments. The team initiated post-treatment monitoring of the habitat and biota in the W1 system, and 2nd-year post-treatment monitoring in the M-lakes system, following 3-4 years of pre-treatment monitoring.

Wilcockson, John

Hatfield Consultants jwilcockson@hatfieldgroup.com

Licence Number: S-12/13-3010-YK

Species: All fish species Location: Prairie Creek

Fish usage and habitat assessment - Small lakes along proposed Winter Road

The objective of this study was to assess fish utilization and habitat of small lakes along a proposed winter road. Lakes would be used as a source of water to produce ice on the road.

Zhu, Xinhua

Fisheries and Oceans Canada xinhua.zhu@dfo-mpo.gc.ca

Licence Number: S-12/13-2000-HR

Species: All fish species Location: Great Slave Lake

Monitor and assess cummulative impacts on Important fish population productivity and community community diversity the Great Slave Lake

This objective of this study was to develop a standard monitoring framework addressing cumulative impacts on fish population productivity and integrity and compare mesh-specific gear selectivity.

GLOSSARY

Abiotic - Not living

Active layer -The area where the soil continually freezes and thaws above the permafrost

Adaptation - A process by which a living organism (human, animal or plant) changes to become better suited to a new environment. This generally on an evolutionary timescale however, in the human context, it may be over a short period.

 $\ensuremath{\textit{Adipose}}$ - Of, relating to, or composed of animal fat; fatty

Aerial - In the air

Aeromagnetic survey - Surveys from aircraft that make use of the magnetic field caused by magnetized rocks in the Earth's crust to make estimates about underlying geology of a given area such as distribution of potential resources

Algae - Simple living aquatic single or multi celled plant organisms that contains chlorophyll

Algorithm - A procedure or formula for solving a problem

Alkali - A basic substance that can range in strength

Analytical - A detailed examination of the structure or some other parameter of a substance or thing

Anoxic - A situation where oxygen is present in very low amounts or not at all, common in water

Annual - Occurs every year

Anthropogenic - Caused by a human action

Anthropology - The study of the human beings including their origins, cultures, evolution

Aquatic - Of water

Aquatic Biota - All living organisms in the aquatic environment

Arable - Land fit to be cultivated

Archaeology - The study of past human life and culture by looking at remains and artifacts like tools

Archean - A period of geologic time from about 3.9 billion years to 2.5 billion years ago

Archival - Pertaining to a collection of documents, normal over long periods of time

Arsenic - A chemical element that is gray in color and that is highly poisonous with no taste

Artifact - A historical tool, weapon or other humanmade object that can be studied

Asexual - An organism that reproduces without the aid of a partner and who passes on all of its genetic information

Atmosphere - The layers of gases that surround and protect the Earth

Attributed - To explain by indicating a cause

Avifauna - the birds of a particular region or period

Bacteria - A large and varied group of single-celled microorganisms

Baseline - A set of information and data serving as a basis for comparison into the future

Bathymetry - Underwater topography. Mapping the underwater contours of the bottoms of water bodies

Beaufort Gyre - The major ice and ocean current circulation of the Arctic Ocean

Benthos - The bottom of the ocean or body of water

Biochemistry - The study of chemical processes in living organisms

Biodiversity - Pertaining to the variety of species in an area

Biogenic - Produced by living organisms or biological processes

Biogeography – The study of the geographical distribution of organisms

Biomass - The total amount of all living material within a specific volume of the environment

Biomes - Distinct areas of the Earth that are common in climate conditions, life forms and physical features like the tundra or woodland

Biostratigraphy - Identification and differentiation of rocks based on the types of fossils they contain

Biotic - Having to do with living organisms

Boreal - Relating to the forest areas of the Northern Temperate Zone that are dominated by coniferous trees such as spruce, fir and pine

Brachiopods - Any of various marine invertebrates of the phylum Brachiopoda, having bivalve dorsal and ventral shells enclosing a pair of tentacled, armlike structures that are used to sweep minute food particles into the mouth. Also called *lampshell*.

Breccia - Rock composed of sharp-angled fragments embedded in a fine-grained matrix

Brunisol Soil - soil type that is associated with forest vegetation. It is usually poorly developed and immature

Carbon¹⁴ – A radioactive isotope of carbon used to date ancient rocks and artifacts

Carnivore - A flesh/meat eating animal

Characterized - To describe an object or idea

Chlorophyll A - A pigment in plants that give them their green color and which absorb energy from the sun. Plants use Chlorophyll to change carbon dioxide and water into food and oxygen

Classification - Organize into groups or categories

Climate – Typical weather patterns of a region over long time periods

Community - All organisms in a particular environment

Comprehend - Being able to understand

Comprehensive - Conveying or including everything or almost everything

Coniferous woodland - A wooded area that is dominated by evergreen trees

Conifers - A group of woody plant commonly known as evergreen trees such as pine, spruce or fir that bears cones

Connectivity - As something is able to connect or relate with another thing

Core - A part removed from the interior of a mass especially to determine the interior composition

Correlated - A mutual relation between two comparable things

Cretaceous - Of or belonging to the geologic time, system of rocks and sedimentary deposits of the third and last period of the Mesozoic Era, characterized by the development of flowering plants and ending with the sudden extinction of the dinosaurs and many other forms of life

Crustacean - any mainly aquatic arthropod usually having a segmented body and chitinous exoskeleton

Cryosols - Cryosols are characterized by frozen soil within 1 metre (39 inches) of the land surface and by waterlogging during periods of thaw. They often show disrupted soil layers, cracks, or patterned surface features such as frost mounds, caused by the physical actions of ice formation and melting. Cryosols may be either mineral soils or humus-rich materials

Cryosphere - frozen water in the form of snow, permanently frozen ground (permafrost), floating ice and glaciers

Cumulative - Objects or ideas that add together

Cyanobacteria - predominantly photosynthetic prokaryotic organisms containing a blue pigment in addition to chlorophyll; occur singly or in colonies in diverse habitats; important as phytoplankton

Deciduous – A plant that lose their leaves during one season, usually winter

Deducing – To draw a conclusion

Deformation - A measurable change in structure, normally for the worse

Degradation - To reduce something or to place something at a lower level

Delta – The land formed where a river deposited silt as it enters into a larger water body, classic example, the Mackenzie Delta

Dendrochronology - A system of dating wooden objects by studying the tree growth rings

Density - A quantity of mass per unit volume

Devonian - Of or belonging to the geologic time, system of rocks, or sedimentary deposits of the fourth period of the Paleozoic Era, characterized by the development of lobe-finned fishes, the appearance of amphibians and insects and the first forests

Discontinuous – Not continuing or linked

Diurnal - Relating to or occurring in a 24-hour period; daily. Occurring or active during the daytime rather than at night

Diversion - A changing of the direction an object is going

Ecology - The science that deals with how living organisms live in relation to each other and their environment

Ecological integrity - Ensuring the relationship in plant and animal communities remains healthy

Ecosystem – The organisms present in a defined area and how they interact with the non-living surrounding (the biotic and the abiotic)

Effluent - A pollutant that flows out from a main source, such as sewage or waste matter

Ekman Grab - A box core type of sediment sampling device.

ELC data - Ecological Land Classification data

Electrofishing - Using electricity to stun and kill fish, usually used during scientific scenarios

Electromagnetic - Magnetism that is caused by electricity

Emissions - A water product that is radiated outward or discharged from a source

Endocrine – 1) designating or of any gland producing one or more hormones 2) designating or of such a hormone

Endophyte - An organism, especially a fungus or microorganism, that lives inside a plant, in a parasitic or mutualistic relationship

Environment – An organism's physical surroundings

Epoch - A period of time during which something important developed or happened

Erosion - Group of natural processes (weathering, disintegration, abrasion, corrosion, transportation) where the Earth's surface is worn away and removed

Eskers - A long, narrow ridge of coarse gravel deposited by a stream flowing under a decaying glacial sheet of ice

Estuary - A place where coastal seawater comes into contact with the current of a freshwater stream

Eukaryote - any member of the *Eukarya*, a domain of organisms having cells each with a distinct nucleus within which the genetic material is contained. Eukaryotes include protoctists, fungi, plants and animals

Eutrophication – The enrichment of aquatic systems, promoting dense algal and plant growth in a body of water, depriving the water of oxygen and forcing change in species composition

Evaporites A sedimentary deposit that results from the evaporation of seawater

Evolution - A process where different species come into existence by differentiation and genetic mutations from common ancestors over a long period of time.

Excavated - Extracting or revealing something by removal of the surrounding earth

Fauna - Animal life of a particular region, environment, or geological period

Fault - A fracture in a rock along which the rocks move; the place of origination of seismic activity; types include: strike-slip and thrust

Fecundity - Ability to reproduce

Fen - Low, flat, swampy land; a bog or marsh

Flora - The plants of a particular region, environment or geological region

Fluvial - Pertaining to something's existence or growth around a stream or river

Fossil - Trace of an organism of a past age, embedded and preserved in the Earth's crust

Fry - Infant fish

Fungi - A kingdom of heterotrophic organisms that produce spores

Fyke - A long, bag-shaped fishing net held open by hoops

Gas hydrates (clathrates) – Crystalline water based solids physically resembling ice, in which small non polar molecules (typically gases) are trapped inside "cages" of hydrogen bonded water molecules

Gender - One's characteristics or traits determined socially as a result of one's sex

Genetic - Pertaining to an organism's traits or characters being linked to genes

Genera - A group of organisms that share common characteristics

Geochemistry - The science that deals with the chemical composition of and chemical changes in the solid matter of the Earth

Geochronological - The chronology of the earth's history as determined by geologic events and not by human history

Geomorphologic - Pertaining to the physical features of the Earth's surface

Glauconite - A greenish mineral of the mica group, a hydrous silicate of potassium, iron, aluminum, or magnesium

Gonad - a gland in which gametes (sex cells) are produced

Grams (g) - A unit of measurement for mass

Habitat - A place where organisms live

Hepatic – (Anatomy) of or relating to the liver; (Botany) botany of or relating to the liverworts

Heterogeneous - A situation where something is in a mixed composition

Holocene - The most recent 11,000 years of the Earth's history starting at the end of the last major iceage, which has been relatively warm

Hydraulic - Pertaining to movement caused by water

Hydroacoustic survey - An echo-sounding (SONAR) survey used for measuring such things as fish stocks, water velocity, etc.

Hydrocarbon – A molecule containing hydrogen and carbon, often petroleum, natural gas and coal

Hydrograph - A graph showing the water level, discharge, or other property of river volume with respect to time

Hydrology - Science dealing with the properties, distribution and circulation of water

Isotope - Atoms that have nuclei with the same number of protons (as the atomic number) but different numbers of neutrons

Igneous - A rock or mineral that solidified from molten or partly molten material, i.e. from magma; one of three rock types with metamorphic and sedimentary

Implement - To put into effect

Iron - A metallic element used for making tools and essential for all living organisms' survival

Jarosite - a yellow to brown secondary mineral consisting of basic hydrated sulphate of iron and potassium in masses or hexagonal crystals

 $\begin{tabular}{ll} \textbf{\textit{Kimberlite}} - An igneous that forms in volcanic pipe, an indicator of diamond deposits \end{tabular}$

Larvae - A premature stage for an insect where it feeds before becoming a pupa

Latitude - A measurement of the from the equator to a given point on the Earth's surface in the north and south direction

Laurentide Ice Sheet - Principal glacial cover of North America during the Pleistocene Epoch (2.6 million - 11,700 years ago). At its maximum extent it spread as far south as latitude 37° N and covered an area of more than 5 million sq mi (13 million sq km). In some areas its thickness reached 8,000 - 10,000 ft (2,400 - 3,000 m) or more

Ligotrophic (oligotrophic) - The opposite of eutrophic. Waters having very low levels of primary productivity and (usually) low concentrations of nutrients; good, clear water quality

Limestone - A sedimentary rock that contains mostly calcium carbonate and can be formed by either inorganic or organic processes

Limnology - The scientific study of the life and phenomena of fresh water, especially lakes and ponds

Lithic - Of, like, or made of stone. Archaeological artifacts made of stone

Meristic - Having or composed of segments; segmented

Mesic - Of, characterized by, or adapted to a moderately moist habitat

Metabolism - The chemical processes occurring within a living cell or organism that are necessary for the maintenance of life. In metabolism some substances are broken down to yield energy for vital processes while other substances, necessary for life, are synthesized

Metamorphic rock - Any rock derived from preexisting rocks by changes in response to environmental factors such as temperature and pressure over a long period of time; one of three types of rocks with igneous and sedimentary

Methane - The simplest hydrocarbon that is the main ingredient in natural gas (CH₄)

Microclimate - The climate of a small area that is different due to changes in geography

Microorganisms - Organisms that must be viewed under a microscope, such as bacteria or a virus

Migration - The long range movement of a group of animals based on the seasons

Molecular analysis - A detailed look at the chemical structure and properties of a molecule

Moraine - A mound of rock debris carried and deposited by a glacier

Multicellular - Composed of more than one cell

Nutrient – Any chemical that an organism removes from the environment to aid with growth and development; common nutrients include nitrogen and phosphorus

 ${\it Otolith}$ – A part of a fish's inner ear, often used to determine the age fish

Organic - Material pertaining to plants or animals

Outcrop - A portion of bedrock or other stratum protruding through the soil level

Overlie - Sedimentary or volcanic rock that lies on top of older rock

Paleoecological - A relationship or study of ancient organisms and how they related to their ancient environment

Paleoenvironmental - An environment that existed in the past

Parr - a juvenile fish

Parameter - One set of measurable factors, such as the temperature and pressure that define a system and determine its behavior and are varied in an experiment

Pelagic - Relating to or living in or on oceanic waters. The pelagic zone of the ocean begins at the low tide mark and includes the entire oceanic water column

Permafrost – The permanently frozen layer of soil that characterizes the Arctic's ground; there are two various types: continuous and discontinuous

Pertinent – An object, idea or concept that is relevant to the topic

Phylogeography - the study of the historical processes that may be responsible for the contemporary geographic distributions of individuals

Phylum – (Biology) a major taxonomic division of living organisms that contain one or more classes. An example is the phylum *Arthropoda* (insects, crustaceans, arachnids, etc., and myriapods)

Physiological - Pertaining to the physical structures and functions of living organisms

Phytoplankton - A group of plant-like plankton that all sea animals depend on either directly or indirectly

Pingo – A large frozen mound covered with vegetation in permafrost areas

Pleistocene - An age of notable ice ages and development of humans between 2,000,000 and 10,000 years ago

Postglacial - Relating to or occurring during the time following a glacial period

ppm - An abbreviation of parts per million

Precipitation – Water (in the form of rain, snow hail, etc.) falling from the atmosphere

Prokaryote - An organism of the kingdom Monera (or Prokaryotae), comprising the bacteria and cyanobacteria, characterized by the absence of a distinct, membrane-bound nucleus or membrane-bound organelles, and by DNA that is not organized into chromosomes. Also called *moneran*

Qualitative – A complete detailed descriptions usually taken from a small sample that allows for distinctions to be drawn from the data

Quantitative - Use of large amounts of data where statistics can be applied to interpret the data

Quaternary - Of or belonging to the geologic time, system of rocks, or sedimentary deposits of the second period of the Cenozoic Era, from the end of the Tertiary

Period through the present, characterized by the appearance and development of humans and including the Pleistocene and Holocene epochs

Qiviuq - The soft downy undercoat of muskoxen

Radiocarbon dating - The determination of the approximate age of an ancient object, such as an archaeological specimen, by measuring the amount of carbon¹⁴ it contains

Raptor - A bird of prey such as an eagle, falcon or osprey

Regolith - The layer of loose rock resting on bedrock, constituting the surface of most land. Also called *mantle rock*

Regosol - a type of azonal soil consisting of unconsolidated material derived from freshly deposited alluvium or sands

Remote Sensing – A technique used to study locations using technology that does not require the researcher to be in the field

Revitalization - To give new life or vitality to something

Riffle – a) A rocky shoal or sandbar lying just below the surface of a waterway b) A stretch of choppy water caused by such a shoal or sandbar; a rapid

Satellite imagery - Computer images generated by a satellite which allow researchers to look at a specific area and monitor surface features such as vegetation

Sediment - Solid fragment material that occurs from the weathering of rocks. In water it is material that has settled from a state of suspension

Sedimentary rock - Rock derived from loose particles that have accumulated over time

Sedimentation - The process where small particles are moved and deposited to accumulate into layers

Seine - A large fishing net made to hang vertically in the water by weights at the lower edge and floats at the top

Seismic - Pertaining to vibrations in the Earth, both natural and induced

Shovel testing - A simple test where a sample of ground is taken by use of a shovel and examined

Species - A group of organisms that share common characteristics that group them together and also distinguish them from others

 $\ensuremath{\textit{Stone flakes/chards}}$ - debris left over from a rock while making tools

Stratified - A system that is set up in layers or strata

Stratigraphic - Formation of rock where different layers can be picked out based on type and age of the rock

Subsidence - The shifting of the Earth's surface downwards (compared normally to sea-level)

Succession - A progressive change in the biological community as a result of a response from species to the changing environment

Surficial - Pertaining to something that is on the surface

Suspension - A situation where the medium is able to support the weight of the particles trapped inside it, example: silt in a river.

Symbioses – An interaction between two or more organisms that usually benefits both

Sympatric - Occupying the same or overlapping geographic areas without interbreeding. Used of populations of closely related species

Systematic - Done according to a plan

Taxonomy - The classification of organisms in an ordered system that indicates natural relationships

Thermokarst - Sinking holes, caves and underground drainage that are produced in regions with permafrost from melting of ground ice and settling of the remaining ground

Theodolite - a surveying instrument for measuring vertical and horizontal angles. Also called (in the US and Canada) **transit**

Thermocline - Layer in a large body of water that sharply separates regions differing in temperature. An abrupt temperature gradient in a lake

Topography - A description of the surface of a given area

Trace metals - A metal that is not essential in the sample but is found in small quantities

Transect - An imaginary line across a surface where observations are made

Tributary - A stream or river which feeds into a larger body of water

Turbid - Stirred up material suspended in a medium leaving it unclear and opaque

Ungulate - Hoofed animals

Velocity - Rate of change of position; quickness of motion

Volatile - Unstable; a substance that easily vapourizes

Watershed - A region draining into a river, river system, or other body of water

Weather – Daily variable changes in temperature, precipitation, wind and other atmospheric conditions

Zooplankton - Microscopic animal organisms floating in water

210-Pb Method - is used to determine the accumulation rate of sediments in lakes, oceans and other water bodies. It is used for over a period of 100 - 200 years

AUTHOR INDEX

Biology 2011
Blaschuk, Katherine 1
Buddle, Chris1
Budziak, Jerry2
Carriere, Suzanne 2
Cote, Jason3
Guthrie, Glen3
Guthrie, Glen4
Johnson, Mary Ann 4
Jones, Paul4
Krizan, Julia5
Langhorne, Amy6
Lantz, Trevor6
Lawson, Nick7
Lennie-Misgeld, Peter . 7
Leski, Michael8
Low, George8
Maier, Kris9
Panayi, Damian9
Robb, Tonia10
Tonn, William11
Trimble, Annika 12
Turetsky, Merritt 12
Venables, Chandra 12
Wilcockson, John 13
Contaminants 2011
Blowes, David14
Evans, Marlene 14
Gantner, Nikolaus
(Klaus)15
Geddes, Robert 15
Krizan, Julia16
McLachlan, Stephane 16
Wiatzka, Gerd17
Wolfe, Brent17
Engineering 2011 Patterson, R. Tim 18
Patterson, R. Tim 18
Trimble, Annika 18
Trimble Annika 19

i rimbie, Annika	. 19
Health 2011	
Badry, Dorothy	. 20
Brennan, Jodi	. 20
Goodman, Karen	
Hammond, Merryl	
Hannon, Judy	. 22
Kuhn, Karen	. 22
Martin, Jim	. 22
Mitton, Craig	. 23
Physical Sciences 201	1
Arrigo, Kevin	. 24
Aubet, Natalie	
Bédard, Jean	
Bhatti, Jagtar	
Bird, Sam	
Bottenheim, Jan	
Burgess, David	
Burn, Chris	
Clark, Ian	
Corriveau, Louise	. 28
Coulton, Daniel	. 29
Dallimore, Scott	
Duffe, Jason	. 30
Duthie, Andrew	
Fortier, Martin	
Grogan, Paul	
Haas, Claudia	
Halverson, Galen	. 33
Herber, Andreas	
Hicks, Faye	. 34
Hilton, Robert	. 34
Holmes, Robert	
Kanigan, Julian	
Kors-Olthof, Rita	
Lafleur, Peter	
Landry, Francois	
Langhorne, Amy	
Lawson, Nick	. 39

Lennie-Misgeld, Peter Lintern, Gwyn	40 40 41 41 42 42 ra
Manua Kristin	40
Moore, Kristin	
Mumford, Thomas	44
O'Neill, Norm	
Paradis, Suzanne	
Pickart, Robert	
Pisaric, Michael	
Quinton, William	46
Reford, Stephen	47
Rose, Rachael	
Russell, Hazen	
Skeries, Kristina	48
Smith, Sharon	
Snyder, David	
Spence, Christopher	
Steele, Michael	51
Turetsky, Merritt	51
Urbanic, Jane Challen	151
Vonk, Jorien	52
Wang, Zhaohui (Aleck	()
Wolfe, Stephen	53
Worthy, Douglas	53
Wrona, Frederick	54
Wrona, Frederick	
Wrona, Frederick	55
ocial Sciences 2011	
Abele, Frances	56
Baumann, Britt	
Cash, Penny	
Delormier, Treena	

Denning, Bryany58	Clarke, Grant80	Reford, Stephen	.95
Duran, Nelida58	Clarke, Grant81	Robertson, Myra	
Duthie, Andrew58	Harris, Ryan81	Sharam, Greg	
Fresque, Jennifer 59	Lobb, Wayne Murray .82	Wood, Cindy	
Gagnon, Graham 59	MacKay, Glen82	Wortham, Jim	
Hampton, Mary60	MacKay, Glen83	Fisheries Permits	
Hodgkins, Andrew 60	Prager, Gabriella 83	2011	
Levitan, Tyler61	Seip, Lisa84	Ash, Jason	.97
Little, Lois61	Walker, Daniel84	Bisaillon, Jean-Franc	
Loovers, Jan Peter 62	Wickham, Michelle84		.97
Luig, Thea62	Wildlife 2011	Bisaillon, Jean-Franc	
Mair, Heather63	Abernathy, David86	,, , , , , , , , , , , , , , , ,	
Noble, Bram63	Bidwell, Mark86	Blais, Jules	
Oehler, Alexander 63	Carriere, Suzanne 86	Bohnet, Seth	
O'Keefe, Jeffrey64	Cluff, Dean86	Bourke, Robin	
Parker, Aliana64	Cluff, Dean87	Cote, Jason	
Pearce, Tristan65	Coulton, Dan87	Fortier, Martin	
Rawluk, Andrea66	Croft, Bruno87	Gantner, Nikolaus	
Reinfort, Breanne 66	Croft, Bruno87	Guthrie, Glen	
Sabin, Jerald67	Davidson, Tracy88	Hanna, Bruce	
Saxon, Leslie67	Davidson, Tracy88	Hanna, Bruce	
Schurr, Theodore 67	Decker, Robert88	Harwood, Lois	
Schurr, Theodore 68	Derocher, Andrew88	Hayward, David	
Schurr, Theodore 69	Elkin, Brett89	Hayward, David	
Simmons, Deborah 69	Enbridge Pipelines89	Howland, Kimberly	
Southcott, Chris70	English, Colleen89	Howland, Kimberly	
Stirling, Mark70	Fast, Marie89	Howland, Kimberly	
Stuhl, Andrew70	Fronczak, David90	Howland, Kimberly	
Todd, Zoe71	Green, David90	Jones, Paul	
Vannini, Phillip71	Groves, Debbie90	Knopp, Jennie	
Wood, Mellissa72	Harpley, David90	Landry, Francois	
Traditional Knowledge	Hegel, Troy91	Landry, Francois	
2011	Kardynal, Kevin91	Leonard, Deanna	
Borowitz, Michelle 73	Kelly, Allicia91	Low, George	
Chavarie, Louise 73	Kelly, Allicia91	Machtans, Hilary	
Davison, Tracy74	Kelly, Allicia92	Maier, Kris	
Lantz, Trevor74	Kutz, Susan92	Mason, Kristine	
Maraj, Ramona74	Larter, Nic92	McPherson, Morag	
Nesbitt, Lorien 75	Larter, Nic92	McPherson, Morag	
Parlee, Brenda75	Larter, Nic93	McPherson, Morag	
Sandlos, John75	Larter, Nic93	Mochnacz, Neil	
Simmons, Deborah 76	Larter, Nic93	Mochnacz, Neil	
	Lausen, Cori93		
Snortland, Jody76 Svoboda, Michael77	Machtans, Craig 93	Nicol, Sandra	
		Panayi, Damian	
Archaeology 2011	Madsen, Erik94	Parent, Bruce	
Andrews, Tom 78	McCallum, Dee94	Pink, Melissa	
Bussey, Jean78	Mulders, Robert94	Pink, Melissa	
Cary, Henry79	Panayi, Damian94	Pink, Melissa	
Cary, Henry79	Popko, Richard 95	Robertson, Dale	
Clarke, Grant80	Rausch, Jennie95	Sawdon, Lorraine	ΙIU

Tonn, William110	Mitton, Craig131	MacNaughton, Robert
Toyne, Melanie 111	Scott, Shannon132	154
Ware, Chaz 111	Young, Barbara 132	Marsh, Philip154
Wiatzka, Gerd 111	Physical Sciences 2012	Messmer, David155
Wilcockson, John 111	Anderson, Natalie 133	Miles, Warner 155
Zhu, Xinhua 112	Armstrong, Terry 133	Moore, Kristin156
Biology 2012	Barber, David134	Narbonne, Guy 156
		O'Neill, Brendan157
Buddle, Chris113	Bartlett, Mike135	Osinski, Gordon157
Hansen, Ken113	Bierwirth, Eike135	Panayi, Damian 158
Harris, Allan114	Bishop, Nicole136	Paradis, Suzanne 158
Jones, Paul114	Bottenheim, Jan 136	Pehrsson, Sally158
Krizan, Julia	Budziak, Jerry137	Pickart, Robert159
Marken, Sandra 115	Burgess, David 137	
Muir, Andrew 115	Burn, Chris138	Pisaric, Michael 159
Renaud, Claude 115	Coulton, Daniel 138	Pratt, Brian160
Robb, Tonia116	Dahl, Mark139	Prinsenberg, Simon . 160
Sibbald, Carey117	Dallimore, Scott 139	Quinton, William 161
Tonn, William117	Derksen, Chris139	Reimink, Jesse 161
Trimble, Annika 118	Desrosiers, Sarah 140	Sachs, Torsten 162
Contaminants 2012	Duffe, Jason140	Schroder-Adams,
Blais, Jules 119	England, John141	Claudia162
Blowes, David 119	English, Michael 141	Smith, Sharon163
Dewar, David120	Fortier, Martin 142	Snyder, David163
Evans, Marlene 120	Grogan, Paul 143	Sofko, George164
Farrell, Rory121	Guest, Bernard 143	Spence, Christopher 164
Gantner, Nikolaus	Haas, Claudia143	Tank, Suzanne165
	Hansen, Ken144	Trimble, Annika165
(Klaus)121	Hansen, Ken144	Trimble, Annika165
Geddes, Robert 121	Harris, Katherine 145	Trimble, Annika166
Livingstone, Steve 122	Haugaard, Rasmus 145	Trusler, Scott166
Low, George	Hicks, Faye146	Turner, Elizabeth 166
Macdonald, Colin 122	Holmes, R. Max146	Urbanic, Jane Challen
McLachlan, Stephane	Hood, Alexandra147	167
123		van der Sanden,
Nash, Tyler123	Hoos, Rick147	Josephus167
Osawa, Akira 124	Kanigan, Julian 147	Vavrek, Matthew168
Purdy, Colin124	Kerr, Jeremy	Wang, Zhaohui (Aleck)
Reimer, Kenneth 125	Kershaw, G. Peter 148	168
Reinfort, Breanne 125	Kjarsgaard, Bruce 148	Whalen, Dustin 168
Stern, Gary126	Kokelj, Steven149	Wilcockson, John169
Wiatzka, Gerd126	Kokelj, Steve149	Williams, Mathew169
Engineering 2012	Krizan, Julia150	Wolfe, Stephen 169
Patterson, R. Tim 127	Lacelle, Denis150	•
Trimble, Annika 127	Lafleur, Peter151	Wrona, Frederick 170
Health 2012	Laidlaw, Shawn 151	Social Sciences 2012
	Lamoureux, Scott 151	Anakin, Megan171
Goodman, Karen 129	Landry, Francois152	Bathe, Adam171
Hammond, Merryl 129	Langhorne, Amy 152	Baumann, Britt172
Hannon, Judy 130	Lantz, Trevor153	Bishop, Nicole172
Janssen, Patricia 130	Machtans, Hilary153	Borowitz, Michelle173
Kuhn, Karen 131	•	
MacLeod, Martha 131		

Callingham, Christina	Van Wyck, Peter 194	Mougeot, Charlotte214
173	von Kuster, Jenica194	Obst, Joachim215
Cash, Penny173	Wesche, Sonia 194	O'Keefe, Harry215
Chugh, Pawan 173	Archaeology 2012	Pongracz, Jodie215
Daitch, Sarah174	Andrews, Tom 196	Pongracz, Jodie216
Douglas, Vasiliki 174	Bussey, Jean196	Pongracz, Jodie216
Fraser, Gail175	Bussey, Jean197	Pongracz, Jodie216
Gordon, Dylan 175	de Guzman, Margarita	Ramona, Maraj216
Hall, Karen175	198	Rausch, Jennie217
Hampton, Mary 176	Jankuta, Kimberly 198	Richardson, Evan217
Hanson, Dorie 176	Kristensen, Todd 199	Robertson, Myra217
Hay, Amie177	MacKay, Glen 199	Robertson, Myra218
Henderson, Joanna. 177	MacKay, Glen200	Robertson, Myra218
Hodgkins, Andrew 177	Murphy, Brent200	Robertson, Myra218
Horowitz, Wayne 178	Ng, Tommy201	Sharam, Greg219
Jaker, Alessandro 178	Ng, Tommy201	Sharam, Greg219
Jardine, Cindy 178	Prager, Gabriella 203	Tout, Ann Marie219
Lizotte, Amy179	Ross, Julie204	Wells, David & Claudine
Luig, Thea179	Seip, Lisa204	Lee219
Mair, Heather180	Thomson, J. Callum. 205	Wortham, Jim219
Mitrovic, Inya180	Walker, Daniel 205	Fisheries Permits 2012
Pearce, Tristan 181	Wildlife 2012	Bisaillon, Jean-Francois
Poole, Nancy181	Armstrong, Terry206	221
Sandlos, John182	Barclay, Robert206	Blanchfield, Paul221
Simmons, Deborah 182	Bidwell, Mark206	Clark, Robert (Bob)221
Smart, Miles 182	Bourn, Stephen207	Cott, Peter222
Svoboda, Michael 183	Carrière, Suzanne 207	Dunmall, Karen222
Todd, Zoe 183	Cluff, Dean207	Evans, Marlene222
Turcotte, Andre 184	Cluff, Dean207	Gantner, Nikolaus222
Webb, Kernaghan 184	Cluff, Dean208	Harris, Katherine223
Will, Alice185	Condon, Wayne208	Harris, Katherine223
Young, Michael 185	Coulton, Daniel 208	Harwood, Lois223
Traditional Knowledge	Coulton, Daniel 209	Hoos, Richard224
2012	Cox, Karl209	Howland, Kimberly224
Alexie, Elaine187	Cox, Karl209	Howland, Kimberly225
Balanoff, Helen 187	Davison, Tracy210	Howland, Kimberly225
Benson, Kristi 188	Davison, Tracy210	Howland, Kimberly225
Brook, Julia188	Davison, Tracy210	Howland, Kimberly226
Coedy, Bill188	Davison, Tracy210	Johnson, Jim226
Fabijan, Michael 189	Elkin, Brett211	Lea, Ellen226
Gauthier, Brenda 189	Hood, Alex211	Lea, Ellen227
Gilday, Cindy189	Kelly, Allicia211	Leonard, Deanna227
Gordon, James Jr 190	Kelly, Allicia212	Loseto, Lisa227
Ireland, Margaret 190	Kelly, Allicia212	Loseto, Lisa228
Jansen, Kelsey 191	Kelly, Allicia212	Low, George228
Lantz, Trevor 191	Kelly, Allicia213	Machtans, Hilary228
Moore, Kristin 192	Kent, Christine213	Machtans, Hilary229
Nash, Tyler193	Klimstra, Jon213	Mason, Kristine229
Rice, Keren193	Larter, Nic213	McPherson, Morag 229
Robinson, Andrew 193	Larter, Nic214	Mochnacz, Neil229

Moore, Kristine 230	Paradis, Adrian 231	Sawdon, Lorraine 233
Morinville, Genevieve	Pink, Melissa232	Tonn, William234
230	Pink, Melissa232	Tonn, William234
Muir, Andrew 230	Reist, Jim232	Wilcockson, John234
Olenick, Adrian 231	Renaud, Claude233	Zhu, Xinhua234
Olenick, Adrian 231	Savage, Pascale-Laure	
Panayi, Damian 231	233	

A	E
Aboriginal Affairs and Northern Development Canada, 6, 37, 103, 114, 138, 142, 178, 179, 182, 194 Aklavik, 6, 22, 38, 62, 67, 68, 72, 73, 76, 79, 118, 119, 142, 164, 165, 167, 173, 176, 178, 181, 182 Alaska, 13, 31, 35, 49, 70, 72, 92, 103, 130, 152, 158, 169, 184 Arctic Ocean, 25, 28, 32, 35, 36, 49, 52, 54, 125, 126, 136 Currents, 148 Arctic Red River, 156 Arthropods, 1, 100 Atmosphere, 28, 125 Clouds, 80, 124 Methane, 152	Education, 2, 4, 17, 24, 26, 59, 63, 64, 65, 66, 67, 72, 161, 164, 167, 168, 169, 174, 190 Elders, 8, 21, 24, 62, 64, 68, 77, 78, 81, 82, 85, 162, 169, 171, 173, 177, 178, 180, 190 Environmental Assessments, 6, 16, 30, 65, 102, 110, 128, 134, 143, 165, 179 Aquatic, 136 Cumulative Impacts, 6, 38, 65, 97, 140, 142, 182 Monitoring, 145 F Fish, 1, 3, 4, 5, 7, 8, 9, 10, 11, 13, 15, 32, 34, 38, 40, 43, 56, 59,
В	64, 73, 78, 79, 85, 100, 101, 102, 103, 105, 108, 109, 110, 111, 112, 114, 125, 134, 136, 142, 143, 145, 147, 159, 168,
Banks Island, 1, 47, 51, 92, 123, 130, 153 Beaufort Sea, 2, 28, 31, 32, 41, 52, 65, 66, 67, 72, 76, 90, 123, 124, 130, 131, 132, 138, 150, 158, 167, 176 Behchoko, 120, 122, 164, 165, 166 Benthic Community, 7, 11, 43, 45, 103, 104, 105, 128, 136, 145 Birds, 88, 89, 91, 97, 183 Ducks, 92, 97, 98 Raptors, 91, 136, 141 Shorebirds, 97 Songbirds, 96 Swan and Geese, 98 Waterfowl, 91, 92, 97, 98 Boreal Forest, 2, 10, 12, 37, 95, 112, 131 Boreholes, 14, 108, 110	174, 179, 180, 183, 190, 194, 201, 202 Angling, 9, 12, 142, 159 Electrofishing, 12, 100, 105, 125, 142, 159 Gill Nets, 8, 102, 111 Fish Species, 9, 109, 110, 111, 134 Burbot, 3, 9, 11, 14, 15, 101, 105, 108, 134 Char, 9 Cisco, 60 Dolly Varden, 9 Grayling, 3, 9, 12, 60, 105, 125, 136 Loche, 60 Long sucker, 12, 105, 125 Pike, 3, 7, 9, 15, 108, 125 Sculpin, 3, 11, 13, 105, 134, 136
	Stickleback, 3, 56, 134, 143 Trout, 3, 9, 12, 13, 14, 15, 45, 59, 75, 102, 105, 108, 111, 114,
С	125, 136
Climate Change, 4, 20, 28, 29, 33, 36, 47, 48, 51, 54, 55, 56, 57, 67, 71, 77, 90, 100, 107, 108, 112, 114, 127, 129, 132, 136, 147, 149, 150, 153, 159, 160, 171, 180, 181, 184, 185 Adaptation, 55, 159 Albedo, 80 Emissions, 28, 31 Community-based methods, 6, 22, 23, 77, 110, 114, 119, 129, 169, 180, 181, 182, 185 Capacity, 11, 64, 105, 126, 178, 182 Contaminant, 53 Arsenic, 112, 113, 145, 183, 188 DDT, 15, 108 Lead, 26, 111, 118, 131, 161 Mercury, 4, 8, 9, 15, 45, 59, 108, 110, 111, 114, 123, 147, 151 Organic, 14, 98, 108	Whitefish, 9, 12, 101, 105, 111 Food Security, 68, 165, 169, 174 Fort Good Hope, 4, 50, 77, 84, 152 Fort Liard, 94, 95, 96, 164, 167 Fort McPherson, 8, 9, 29, 36, 38, 47, 48, 64, 76, 79, 81, 118, 146, 149, 164, 167, 170, 173, 176, 177, 178, 181 Fort Providence, 20, 25, 53, 103, 123, 155, 157, 164, 167, 179, 185 Fort Resolution, 5, 14, 15, 16, 17, 61, 75, 77, 78, 82, 85, 86, 94, 101, 108, 111, 163, 164, 172, 173, 184, 193 Fort Simpson, 8, 23, 27, 48, 50, 53, 94, 95, 103, 120, 122, 152, 164, 167, 170, 184 Fort Smith, 5, 10, 12, 16, 25, 58, 66, 93, 94, 95, 101, 103, 111, 112, 119, 122, 123, 161, 164, 174, 175, 179
Remediation, 2, 12, 16, 17, 105, 114, 115, 126, 188, 189	G
D Daring Lake Tundra Ecosystem Research Station, 12, 13, 33, 39,	Gamètì, 30, 194 Geotechnical Investigation, 16, 38, 55, 102, 159 Glaciers, 34, 127, 130, 131, 138
96, 129, 132, 140	Goyatiko, 77
Dehcho, 8, 9, 40, 50, 66, 67, 71, 84, 85, 91, 93, 94, 95, 110, 153, 190 Déline, 7, 8, 71, 75, 77, 78, 81	Great Bear Lake, 7, 15, 30, 51, 75, 77, 81, 102, 110, 111 Great Slave Lake, 5, 14, 15, 17, 25, 30, 39, 43, 46, 51, 61, 70, 82, 85, 86, 98, 102, 103, 108, 114, 122, 135, 143, 180, 185, 188,
Dené, 82, 83	193 Gwich'in, 4, 9, 29, 38, 50, 64, 68, 69, 70, 76, 81, 90, 153, 170,
Dettah, 23, 60, 63, 77, 85, 89, 120, 168, 172, 180, 193 Distance Early Warning (DEW) Line, 179	177, 178, 181, 184

Н	Kendall Island Bird Sanctuary, 47
Hay River, 8, 9, 14, 15, 25, 35, 40, 65, 77, 86, 90, 103, 108, 110,	L
135, 161, 164, 166, 169, 170, 172, 184	_
Health	Lakes
Addiction, 60, 150, 167, 172	Water Level, 40, 61
Environmental Factors, 16, 111	Lichen, 6
H. pylori, 22, 118	LiDAR, 129
Mental Health, 167, 172, 176	Linguistics, 168
Pregnancy, 119	Lithic scatters, 85, 190
Primary Helath Care, 119	
Sexual Health, 23, 60	M
Smoking and Tobacco, 22, 118, 119, 169	141
Health Care, 24, 119, 120, 121 Highways and Roads	Mackenzie Delta, 5, 6, 17, 30, 31, 37, 41, 42, 47, 54, 56, 57, 61,
Dempster Highway, 38, 76, 100, 112, 137, 138, 139, 142, 146,	66, 67, 72, 73, 76, 90, 98, 101, 107, 122, 127, 128, 130, 138,
181	142, 144, 146, 149, 150, 151, 152, 154, 155, 157, 158, 167,
Highway Development, 139, 195	176, 179, 181
Homelessness, 21, 58, 172, 176	Mackenzie Gas Project, 37, 171
Hospital	Mackenzie River, 7, 25, 36, 54, 70, 94, 102, 103, 130, 134, 135,
Stanton Territorial Hospital, 23, 120, 121	155, 157, 158, 179, 191, 192, 193, 196
Hunters and Trappers, 6, 15, 76, 142	Mammals, 33, 88, 131, 141, 145, 149
	Beaver, 185
1	Caribou, 38, 71, 78, 88, 89, 90, 91, 92, 94, 95, 96, 98, 132,
I	173, 174, 184
Ice, 7, 11, 13, 15, 19, 28, 29, 32, 35, 36, 37, 52, 54, 55, 56, 57,	Grizzly Bear, 178
73, 80, 85, 104, 114, 116, 123, 124, 126, 127, 128, 131, 132,	Moose, 77, 90, 93, 94, 95, 141, 185
134, 135, 139, 140, 150, 157, 159, 160, 179, 185, 186, 189,	Muskrat, 169 Polar Bears, 76, 90
190, 192, 201	Wolf, 100
Breakup, 17, 35, 54, 56, 135, 157, 160	Wolverine, 96, 136, 178
Ice jams, 35	Wood Bison, 95, 175, 185
Industrial development, 93	Melville Ice Cap, 28, 126, 127, 141
Mine development, 17, 61, 77, 85, 111, 112, 114, 125, 134,	Métis, 17, 40, 60, 61, 82, 86, 111, 145, 162, 173, 182, 188
156, 162, 173, 183, 188, 189, 193, 194	North Slave Métis Alliance, 86, 162, 182, 188
Industrial Development, 30, 43, 44, 82, 98, 145, 190	Mines
Diamond Mines, 6, 11, 14, 40, 42, 43, 45, 49, 63, 91, 96, 103,	Con Mine, 143
105, 107, 108, 136, 142, 145, 153, 182, 187	EKATI, 11, 103
Oil & Gas, 31, 128	Giant Mine, 77, 111, 112, 134, 173, 183, 188, 189
Rare Mineral Mines, 7, 19, 40, 46, 104, 175	Snap Lake, 42, 43, 96, 136
Winter Roads, 19, 55, 80, 96, 116, 159, 186, 187	Mushrooms and Fungi, 113, 165
Ingraham Trail, 55, 159, 189	
International Polar Year, 28	N
Inuit, 70, 71, 165, 171, 180, 182	IN
Inuvialuit, 5, 15, 22, 26, 27, 33, 50, 65, 66, 67, 68, 69, 70, 72, 76,	National Parks
83, 84, 98, 109, 113, 118, 127, 139, 153, 174, 179, 181, 184,	Nahanni, 27, 93, 94, 95, 101, 150, 159, 190
195	Wood Buffalo, 10, 95, 96, 112, 175
Inuvik, 5, 7, 8, 9, 10, 12, 15, 16, 19, 20, 27, 28, 29, 35, 36, 42, 47,	Ndilo, 60, 77, 113, 164, 168, 169, 180
48, 50, 51, 53, 54, 58, 63, 64, 65, 66, 72, 73, 76, 79, 81, 83, 90, 92, 101, 105, 109, 112, 116, 124, 127, 128, 130, 132, 136,	Norman Wells, 1, 2, 3, 4, 27, 41, 42, 50, 84, 87, 91, 97, 100, 102,
137, 138, 139, 144, 151, 153, 154, 155, 158, 162, 164, 165,	134, 143, 152, 153, 157, 173, 184, 191, 192, 193
167, 168, 169, 171, 173, 174, 176, 178, 179, 180, 181, 182,	North Slave, 55, 69, 80, 82, 83, 85, 86, 88, 89, 96, 159, 162, 182,
184, 195	188, 194, 195
104, 100	Nunavik, 120
	Nunavut, 26, 31, 51, 80, 121, 129, 130, 139, 164, 186
J	Nursing, 24, 59
Jean Marie River, 8, 9, 20, 34, 37, 50, 85, 110, 152, 155, 180,	Nutrition, 59
181, 184, 190	Traditional Foods, 60, 165
101, 104, 100	_
1/	0
K	Old Craw Flata 70
	Old Crow Flats, 79

Katlodeeche First Nation, 40, 78, 164

P

Palaeontological, 157, 195 Palentology, 20, 34, 84, 130, 146, 150, 152, 157, 186 Paulatuk, 29, 49, 53, 67, 73, 76, 97, 127, 157, 163, 165, 174 Peel River, 140 Permafrost, 6, 9, 14, 29, 31, 32, 37, 38, 48, 50, 53, 55, 57, 107, 112, 127, 128, 134, 137, 138, 139, 140, 142, 146, 149, 150, 151, 153, 154, 159, 181, 182 Active layer, 6, 37, 50, 55, 110, 142, 153, 160 Petroleum, 16, 110, 191, 192, 193 Phytoplankton, 7, 11, 43, 45, 103, 104, 145 Pingo, 29 Plant Surveys Seed Collection, 105 Transects, 13, 45, 47, 82, 129, 144, 149, 193 Port Radium, 77, 111, 172 Pregnancy, 22, 111 Prince Patrick Island, 16

R

RADAR, 31, 51, 124, 129, 134, 153, 157 Renewable Energy Wind Energy, 20, 117, 155, 156 Residential schools, 164

S

Sachs Harbour, 35, 68, 76, 90, 113, 118, 123, 128, 130, 176
Sahtú, 3, 4, 50, 66, 67, 69, 71, 78, 80, 81, 84, 87, 91, 93, 94, 97, 111, 153, 173, 191, 192, 193
Shovel tests, 82, 83, 87, 189, 191, 192, 193
Snow, 5, 7, 28, 32, 35, 42, 49, 52, 57, 73, 82, 86, 93, 100, 104, 127, 129, 131, 132, 135, 138, 143, 144, 146, 154, 157, 182, 186, 188
Social Justice, 172
South Slave, 61, 66, 67, 71, 86, 88, 90, 93, 94, 96, 145, 161, 163
Streams, 11, 13, 26, 29, 35, 37, 38, 41, 42, 43, 48, 52, 104, 105, 125, 128, 133, 138, 139, 140, 142, 144, 154, 201
Flow, 35, 41, 42, 48, 52, 133, 140
Sumps, 37, 138

Τ

Tł_{Ic}ho, 80
Tł_{Ic}ho, 3, 23, 30, 63, 66, 67, 69, 70, 80, 86
Tobacco, 118, 169
Traditional Knowledge, 7, 8, 34, 40, 75, 76, 77, 109, 161, 162, 171, 178, 179, 180, 181, 182, 185
Trees, 6, 10, 25, 29, 55, 112, 130, 137, 142, 150
Treeline, 37, 137, 138
Tsiigehtchic, 36, 50, 53, 76, 79, 135, 136, 144, 167, 173, 178, 181
Tuktoyaktuk, 5, 15, 22, 27, 42, 50, 53, 58, 65, 72, 73, 76, 79, 83, 90, 92, 109, 118, 128, 130, 132, 137, 139, 152, 157, 158, 162, 164, 165, 167, 168, 173, 176, 179, 181, 182, 195

U

Ulukhaktuk, 62

٧

Victoria Island, 26, 51, 92, 147

W

Weather Station, 28, 40, 41, 43, 127, 139, 142, 149 Wekweètì, 155 Wekweètì, 20, 23, 89, 96, 131 Wetlands, 48, 53 Whatì, 163 Whitehorse, 76, 79, 93, 173

Υ

Yellowknife, 1, 2, 3, 7, 9, 10, 12, 14, 21, 23, 26, 30, 31, 33, 34, 36, 37, 38, 40, 41, 42, 43, 44, 45, 46, 48, 55, 58, 59, 60, 61, 63, 66, 67, 69, 71, 72, 73, 74, 77, 78, 80, 83, 85, 86, 88, 89, 91, 93, 96, 97, 98, 103, 104, 107, 109, 111, 113, 120, 121, 123, 125, 127, 133, 134, 135, 136, 138, 143, 144, 145, 147, 157, 159, 162, 163, 164, 165, 166, 168, 169, 172, 173, 174, 175, 177, 178, 179, 180, 182, 186, 187, 188, 189, 190, 193

Youth, 4, 24, 60, 65, 68, 78, 81, 82, 142, 163, 164, 167, 169, 180, 181, 182, 183

Yukon, 8, 9, 31, 34, 36, 46, 72, 76, 93, 100, 127, 130, 136, 137, 146, 148, 158





